

Pledge of Honesty

I affirm that I will not give or receive any unauthorized help on this exam and that all work will be my own.

Signature: _____

SCAN THIS PAGE WITH THE HONOR PLEDGE SIGNED AND UPLOAD IT WITH YOUR EXAM

Use a separate page (or pages) for each problem. Show all of your work.

1. **Fermat curve question.** (20 POINTS) Consider the subset V_d of the complex projective plane \mathbf{CP}^2 defined by the equation

$$x^d + y^d + z^d = 0 \quad \text{for a positive integer } d.$$

It is known as the *Fermat curve* of degree d . Define a map $f : V_d \rightarrow \mathbf{CP}^1$ by

$$[x, y, z] \mapsto [x, y].$$

A map of this type is called a *branched covering*. It does *not* extend to all of \mathbf{CP}^2 because it is not defined on the point $[0, 0, 1]$.

- (a) Find and count the points in the target whose preimage is *not* a set of d points in V_d . Let $K \subseteq \mathbf{CP}^1$ denote the set of these points. They are called **BRANCH POINTS**.
- (b) You may assume that the restriction of f to the preimage of $\mathbf{CP}^1 - K$ is a d -fold covering of $\mathbf{CP}^1 - K$. Use this fact to find the Euler characteristic of V_d . You may also use the fact that under suitable hypotheses, $\chi(A \cup B) = \chi(A) + \chi(B) - \chi(A \cap B)$.
2. **Complete bipartite graph question.** 20 POINTS A *bipartite graph* is one in which the vertices fall into two disjoint sets, say red and blue vertices, and each edge connects a red vertex to a blue one. It is *complete* if there is a unique edge connecting each red vertex to each blue one.

Let $K_{m,n}$ denote the complete bipartite graph with m red vertices and n blue ones. Hence it has mn edges. For example, $K_{3,3}$ is the houses and utilities graph, which is known to be nonplanar.

Show that if $K_{m,n}$ can be embedded in a closed oriented surface of genus g , then

$$g \geq \frac{(m-2)(n-2)}{4}.$$

3. **Five lemma question.** (20 POINTS) The 5-lemma says that given a commutative diagram of abelian groups with exact rows,

$$\begin{array}{ccccccccc}
 A & \xrightarrow{i} & B & \xrightarrow{j} & C & \xrightarrow{k} & D & \xrightarrow{\ell} & E \\
 \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \delta & & \downarrow \epsilon \\
 A' & \xrightarrow{i'} & B' & \xrightarrow{j'} & C' & \xrightarrow{k'} & D' & \xrightarrow{\ell'} & E'
 \end{array}$$

if α , β , δ and ϵ are isomorphisms, then so is γ . Show by counterexample that the triviality of α , β , δ and ϵ does *not* imply the triviality of γ .

4. **Brouwer Fixed Point question.** (20 POINTS) Prove the 2-dimensional case of the Brouwer Fixed Point Theorem, i.e., that any continuous map of the 2-dimensional disk D^2 to itself has a fixed point. You may assume $\pi_1 S^1 = \mathbf{Z}$.
5. **James reduced product question.** (20 POINTS) Let the 2-sphere S^2 have base point x_0 . Define an equivalence relation on the n -fold Cartesian product $(S^2)^{\times n}$ by saying that when a coordinate in a point

$$(x_1, x_2, \dots, x_n) \in (S^2)^{\times n}$$

is the base point x_0 , it may be transposed with either the coordinate on its left or the one its right. For example the points

$$(x_0, x_1, x_2), (x_1, x_0, x_2) \text{ and } (x_1, x_2, x_0) \in (S^2)^{\times 3}$$

are all equivalent for arbitrary points $x_1, x_2 \in S^2$. The space $J_n S^2$ of equivalence classes in $(S^2)^{\times n}$ is called the *n*th James reduced product of S^2 , having first been studied by Ioan James in the 1950s. One could make a similar definition with S^2 replaced by any pointed space. Thus there is a surjective map $f_n : (S^2)^{\times n} \rightarrow J_n S^2$.

We know that $(S^2)^n$ has a CW-structure with $\binom{n}{k}$ cells in dimension $2k$ for $0 \leq k \leq n$. We also know that as a ring under cup product,

$$H^*((S^2)^{\times n}; \mathbf{Z}) \cong \mathbf{Z}[y_i : 1 \leq i \leq n]/(y_i^2),$$

with $y_i \in H^2$ being the generator associated with the i th factor of the Cartesian product.

It can be shown that $J_n S^2$ has a CW-structure with a single cell in every even dimension up to $2n$. For $1 \leq k \leq n$, the group $H^{2k}(J_n S^2; \mathbf{Z}) \cong \mathbf{Z}$ has a generator u_k whose image under f_n^* is the sum of all square free k -fold products of the y_i s.

Use this information to determine the cup product structure of $H^*(J_n S^2; \mathbf{Z})$. Give a formula for $u_k u_\ell$ as a multiple of $u_{k+\ell}$ for $k + \ell \leq n$. In particular f_n induces a monomorphism in cohomology.

HINT: Try doing this first for small values of n such as 2, 3 and 4.