




ELSEVIER

Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



Hurewicz images of real bordism theory and real Johnson–Wilson theories



Guchuan Li^a, XiaoLin Danny Shi^{b,*}, Guozhen Wang^c,
Zhouli Xu^d

^a Department of Mathematics, Northwestern University, Evanston, IL 60208,
United States of America

^b Department of Mathematics, Harvard University, Cambridge, MA 02138,
United States of America

^c Shanghai Center for Mathematical Sciences, Fudan University, Shanghai, China,
200433

^d Department of Mathematics, Massachusetts Institute of Technology, Cambridge,
MA 02142, United States of America

ARTICLE INFO

Article history:

Received 4 November 2017

Received in revised form 17 October
2018

Accepted 21 October 2018

Available online 16 November 2018

Communicated by A. Blumberg

Keywords:

Hurewicz image

Real bordism theory

Real Johnson–Wilson theory

Slice spectral sequence

C_2 -equivariant May spectral
sequence

ABSTRACT

We show that the Hopf elements, the Kervaire classes, and the $\bar{\kappa}$ -family in the stable homotopy groups of spheres are detected by the Hurewicz map from the sphere spectrum to the C_2 -fixed points of the Real bordism spectrum. A subset of these families is detected by the C_2 -fixed points of Real Johnson–Wilson theory $ER(n)$, depending on n . In the proof, we establish an isomorphism between the slice spectral sequence and the C_2 -equivariant May spectral sequence of $BP_{\mathbb{R}}$.

© 2018 Elsevier Inc. All rights reserved.

* Corresponding author.

E-mail addresses: guchuanli2013@u.northwestern.edu (G. Li), dannyshi@math.harvard.edu (X.D. Shi), wangguozhen@fudan.edu.cn (G. Wang), xuzhouli@mit.edu (Z. Xu).

Contents

| | |
|--|-----|
| 1. Introduction | 68 |
| 2. The equivariant dual Steenrod algebra and Adams spectral sequence | 75 |
| 3. The slice spectral sequence of $BP_{\mathbb{R}}$ | 87 |
| 4. The equivariant May spectral sequence | 93 |
| 5. Map of May spectral sequences | 100 |
| 6. Detection theorem | 104 |
| References | 114 |

1. Introduction

1.1. Motivation and main results

In 2009, Hill, Hopkins, and Ravenel resolved a longstanding open problem in algebraic topology. In their seminal paper [18], they showed that the Kervaire invariant elements θ_j do not exist for $j \geq 7$ (see also [32,16,17] for surveys on the result). The crux of their proof relies on a detecting spectrum Ω , which detects the Kervaire invariant elements.

Theorem 1.1 (*Hill–Hopkins–Ravenel detection theorem*). *If $\theta_j \in \pi_{2j+1-2}\mathbb{S}$ is an element of Kervaire invariant 1, and $j > 2$, then the Hurewicz image of θ_j under the map $\pi_*\mathbb{S} \rightarrow \pi_*\Omega$ is nonzero.*

The detecting spectrum Ω is constructed as the C_8 -fixed point of a genuine C_8 -equivariant spectrum $\Omega_{\mathbb{O}}$, which is an equivariant localization of $MU^{((C_8))} := N_{C_2}^{C_8}MU_{\mathbb{R}}$. Here, $MU_{\mathbb{R}}$ is the Real cobordism spectrum of Landweber, Fujii, and Araki [27,9,2] and $N_{C_2}^{C_8}(-)$ is the Hill–Hopkins–Ravenel norm functor. To analyze the equivariant homotopy groups of $\Omega_{\mathbb{O}}$, Hill, Hopkins, and Ravenel generalized the C_2 -equivariant filtration of Hu–Kriz ([20]) and Dugger ([8]) to a G -equivariant Postnikov filtration for all finite groups G . They called this the slice filtration. Given any G -equivariant spectrum X , the slice filtration produces the slice tower $\{P^*X\}$, whose associated slice spectral sequence is strongly convergent and converges to the $RO(G)$ -graded homotopy groups $\pi_{\star}^G X$. Using the slice spectral sequence, Hill, Hopkins, and Ravenel proved that

$$\pi_{2j+1-2}\Omega = \pi_{2j+1-2}^{C_8}\Omega_{\mathbb{O}} = 0$$

for all $j \geq 7$, hence deducing the nonexistence of the corresponding Kervaire invariant elements.

We are interested in proving more detection theorems for the fixed points of the equivariant theories $MU^{((C_{2^n}))} := N_{C_2}^{C_{2^n}}MU_{\mathbb{R}}$ and their localizations. Our motivation is as follows: classically, π_*MU is a polynomial ring, hence torsion free, and the map $\pi_*\mathbb{S} \rightarrow \pi_*MU$ detects no nontrivial elements in the stable homotopy groups of spheres. Equivariantly, however, computations of Hu–Kriz [20], Dugger [8], Kitchloo–Wilson [22], and Hill–Hopkins–Ravenel [18,19] show that there are many torsion classes in the equivariant homotopy groups of the theories above. Since the Kervaire invariant elements are detected by the fixed point of a localization of $MU^{((C_8))}$, there should be other classes

in the stable homotopy groups of spheres that are also detected by such theories. We prove this is indeed the case.

Theorem 1.2 (*Theorem 6.11, detection theorems for $MU_{\mathbb{R}}$ and $BP_{\mathbb{R}}$*). *The Hopf elements, the Kervaire classes, and the $\bar{\kappa}$ -family (see Definition 1.4) are detected by the Hurewicz maps $\pi_*\mathbb{S} \rightarrow \pi_*MU_{\mathbb{R}}^{C_2}$ and $\pi_*\mathbb{S} \rightarrow \pi_*BP_{\mathbb{R}}^{C_2}$.*

Once we obtain the detection theorem for $\pi_*MU_{\mathbb{R}}^{C_2}$, we use the Hill–Hopkins–Ravenel norm functor to show that these elements are also detected by the C_{2^n} -fixed point of $MU^{(C_{2^n})}$:

Corollary 1.3 (*Corollary 6.13, detection theorem for $MU^{(G)}$*). *For any finite group G containing C_2 , the G -fixed point of $MU^{(G)}$ detects the Hopf elements, the Kervaire classes, and the $\bar{\kappa}$ -family.*

We pause here to discuss some implications of Theorem 1.2, as well as what we mean by the “ $\bar{\kappa}$ -family”. It is well known that the Hopf elements are represented by the elements

$$h_i \in \text{Ext}_{\mathcal{A}_*}^{1,2^i}(\mathbb{F}_2, \mathbb{F}_2)$$

on the E_2 -page of the classical Adams spectral sequence at the prime 2. By Adams’s solution of the Hopf invariant one problem [1], only h_0, h_1, h_2 , and h_3 survive to the E_{∞} -page. By Browder’s work [6], the Kervaire classes $\theta_j \in \pi_{2^{j+1}-2}\mathbb{S}$, if they exist, are represented by the elements

$$h_j^2 \in \text{Ext}_{\mathcal{A}_*}^{2,2^{j+1}}(\mathbb{F}_2, \mathbb{F}_2)$$

on the E_2 -page. For $j \leq 5$, h_j^2 survives. The case $\theta_4 \in \pi_{30}\mathbb{S}$ is due to Barratt, Mahowald, and Tangora [31,4], and the case $\theta_5 \in \pi_{62}\mathbb{S}$ is due to Barratt, Jones, and Mahowald [5]. The fate of h_6^2 is unknown. Hill, Hopkins, and Ravenel [18] showed that the h_j^2 , for $j \geq 7$, cannot survive to the E_{∞} -page. Given this information, Theorem 1.2 and Corollary 1.3 assert that the elements η, ν, σ , and θ_j , for $1 \leq j \leq 5$, are detected by $\pi_*^G MU^{(G)}$. The last unknown Kervaire class, θ_6 , will also be detected, should it survive the Adams spectral sequence.

To introduce the $\bar{\kappa}$ -family, we appeal to Lin’s complete classification of the groups $\text{Ext}_{\mathcal{A}_*}^{\leq 4,t}(\mathbb{F}_2, \mathbb{F}_2)$ [29]. In his classification, Lin showed that there is a family $\{g_k \mid k \geq 1\}$ of indecomposable elements with

$$g_k \in \text{Ext}_{\mathcal{A}_*}^{4,2^{k+2}+2^{k+3}}(\mathbb{F}_2, \mathbb{F}_2).$$

The first element of this family, g_1 , is in bidegree $(4, 24)$. It survives the Adams spectral sequence to become $\bar{\kappa} \in \pi_{20}\mathbb{S}$. It is for this reason that we name this family the $\bar{\kappa}$ -family. The element g_2 also survives to become the element $\bar{\kappa}_2 \in \pi_{44}\mathbb{S}$. Theorem 1.2

and Corollary 1.3 assert that they are both detected by $\pi_*^G MU^{((G))}$. Recent computations of Isaksen–Wang–Xu [21] show that g_3 supports a nontrivial d_5 -differential and therefore $\bar{\kappa}_3$ does not exist in $\pi_{92}\mathbb{S}$. For $k \geq 4$, the fate of g_k is unknown (g_4 is in stem 188). Nevertheless, they will be detected by $\pi_*^G MU^{((G))}$, should they survive the Adams spectral sequence.

Definition 1.4. The $\bar{\kappa}$ -family consists of the homotopy classes detected by the surviving g_k -family.

To prove Theorem 1.2, first observe that 2-locally, $MU_{\mathbb{R}}$ splits as a wedge of suspensions of $BP_{\mathbb{R}}$. Therefore we only need to prove the claim for $BP_{\mathbb{R}}$. To establish the link between the families $\{h_i\}$, $\{h_j^2\}$, and $\{g_k\}$ and the equivariant homotopy groups of $BP_{\mathbb{R}}$, we use the C_2 -equivariant Adams spectral sequence developed by Greenlees [11–13] and Hu–Kriz [20]. More precisely, we analyze the following maps of Adams spectral sequences

$$\begin{array}{ccc}
 \text{classical Adams spectral sequence of } \mathbb{S} & \Longrightarrow & (\pi_*\mathbb{S})_2^\wedge \\
 \downarrow & & \downarrow \\
 C_2\text{-equivariant Adams spectral sequence of } \mathbb{S} & \Longrightarrow & (\pi_{\star}^{C_2}\mathbb{S})_2^\wedge \\
 \downarrow & & \downarrow \\
 C_2\text{-equivariant Adams spectral sequence of } BP_{\mathbb{R}} & \Longrightarrow & (\pi_{\star}^{C_2}BP_{\mathbb{R}})_2^\wedge
 \end{array}$$

and prove the following.

Theorem 1.5 (Algebraic detection theorem). *The images of the elements $\{h_i \mid i \geq 1\}$, $\{h_j^2 \mid j \geq 1\}$, and $\{g_k \mid k \geq 1\}$ on the E_2 -page of the classical Adams spectral sequence of \mathbb{S} are nonzero on the E_2 -page of the C_2 -equivariant Adams spectral sequence of $BP_{\mathbb{R}}$.*

It turns out that for degree reasons, the C_2 -equivariant Adams spectral sequence of $BP_{\mathbb{R}}$ degenerates after the E_2 -page. From this, Theorem 1.2 easily follows from Theorem 1.5 because if any of h_i , h_j^2 , or g_k survives to the E_∞ -page of the classical Adams spectral sequence to represent an element in the stable homotopy groups of spheres, it must be detected by $\pi_{\star}^{C_2}BP_{\mathbb{R}}$.

The proof of Theorem 1.5 requires us to analyze the algebraic maps

$$\text{Ext}_{\mathcal{A}_*}(\mathbb{F}_2, \mathbb{F}_2) \rightarrow \text{Ext}_{\mathcal{A}_*^m}(H\underline{\mathbb{F}}_{2,\star}, H\underline{\mathbb{F}}_{2,\star}) \rightarrow \text{Ext}_{\Lambda_\star^m}(H\underline{\mathbb{F}}_{2,\star}, H\underline{\mathbb{F}}_{2,\star}).$$

They are maps on the E_2 -pages of the Adams spectral sequences above. Here, $\mathcal{A}_* := (H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2)_*$ is the classical dual Steenrod algebra; $\mathcal{A}_\star^m := (H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2)_\star$ is the genuine C_2 -equivariant dual Steenrod algebra; Λ_\star^m is a quotient of \mathcal{A}_\star^m . Hu and Kriz [20] studied \mathcal{A}_\star^m and completely computed the Hopf algebroid structure of $(H\underline{\mathbb{F}}_{2,\star}, \mathcal{A}_\star^m)$. We borrow extensively their formulas. More precisely, we use their formulas to describe the maps

$$(H\mathbb{F}_2, \mathcal{A}_*) \rightarrow (H\underline{\mathbb{F}}_{2\star}, \mathcal{A}_\star^m) \rightarrow (H\underline{\mathbb{F}}_{2\star}, \Lambda_\star^m)$$

of Hopf-algebroids. Then, by filtering these Hopf algebroids compatibly, we produce maps of May spectral sequences:

$$\begin{array}{ccc} \text{Modified May spectral sequence of } \mathbb{S} & \xlongequal{\quad} & \text{Ext}_{\mathcal{A}_*}(\mathbb{F}_2, \mathbb{F}_2) \\ \downarrow & & \downarrow \\ C_2\text{-equivariant May spectral sequence of } \mathbb{S} & \xrightarrow{\quad} & \text{Ext}_{\mathcal{A}_\star^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star}) \\ \downarrow & & \downarrow \\ C_2\text{-equivariant May spectral sequence of } BP_{\mathbb{R}} & \xrightarrow{\quad} & \text{Ext}_{\Lambda_\star^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star}). \end{array}$$

To analyze these maps, we appeal again to Hu and Kriz’s formulas. We compute the maps on the E_2 -page of the May spectral sequences above, as well as all the differentials in the C_2 -equivariant May spectral sequence of $BP_{\mathbb{R}}$.

The readers should be warned that the May spectral sequence at the top of the diagram above is *not* the classical May spectral sequence. The classical May spectral sequence is constructed from an increasing filtration of the dual Steenrod algebra \mathcal{A}_* . However, in constructing the equivariant May spectral sequence, we filtered \mathcal{A}_\star^m and Λ_\star^m by decreasing filtrations. To rectify this mismatch of filtrations, we need to change the filtration of \mathcal{A}_* to a decreasing filtration as well. This is necessary to ensure the compatibility of filtrations with respect to the map $\mathcal{A}_* \rightarrow \mathcal{A}_\star^m \rightarrow \Lambda_\star^m$ — or we won’t have a map of spectral sequences. Nevertheless, despite this change of filtration, we are able to compute this modified May spectral sequence. This computation, together with our knowledge of the C_2 -equivariant May spectral sequence of $BP_{\mathbb{R}}$, finishes the proof of Theorem 1.5.

While proving Theorem 1.5, we also prove a connection between the equivariant May spectral sequence of $BP_{\mathbb{R}}$ and the slice spectral sequence of $BP_{\mathbb{R}}$.

Theorem 1.6 (Theorem 4.9). *The integer-graded C_2 -equivariant May spectral sequence of $BP_{\mathbb{R}}$ is isomorphic to the associated-graded slice spectral sequence of $BP_{\mathbb{R}}$.*

By the “associated-graded slice spectral sequence”, we mean that whenever we see a \mathbb{Z} -class on the E_2 -page, we replace it by a tower of $\mathbb{Z}/2$ -classes. Theorem 1.6 can be intuitively explained as follows: since the Adams spectral sequence for $BP_{\mathbb{R}}$ collapses for degree reasons, the equivariant May spectral sequence of $BP_{\mathbb{R}}$ converges to an associated-graded of $(\pi_\star BP_{\mathbb{R}})_2^\wedge$. On the other hand, the slice spectral sequence

$$E_2^{s,V} = \pi_{V-s}^{C_2} P_{\dim V}^{\dim V} BP_{\mathbb{R}} \implies \pi_\star^{C_2} BP_{\mathbb{R}}$$

also computes the equivariant homotopy groups of $BP_{\mathbb{R}}$. Moreover, works of [20] and [17] essentially show that the C_2 -slice differentials are produced from equivariant cohomology

in $\text{Ext}_{A_*}(\mathbb{F}_2, \mathbb{F}_2)$, only finitely many elements survive to the E_∞ -page of the classical Adams spectral sequence.

Here, $Sq^0(-)$ is the Steenrod action defined on the Adams E_2 -page (see [7]). In particular, the families $\{h_i \mid i \geq 1\}$, $\{h_j^2 \mid j \geq 1\}$, and $\{g_k \mid k \geq 1\}$ are all Sq^0 -families on the 1-line, 2-line, and 4-line of the classical Adams spectral sequence, respectively. We are interested in the fate of the $\bar{\kappa}$ -family $\{g_k \mid k \geq 1\}$ in $\pi_*^G MU^{((G))}$ as we increase the order of G . As G grows bigger, it's possible that g_k will all support differentials in the slice spectral sequence of $\pi_*^G MU^{((G))}$ for k large enough, hence not surviving the classical Adams spectral sequence.

In [18], Hill, Hopkins, and Ravenel also used an algebraic detection theorem to prove that the Kervaire classes are detected by $\pi_*^{C_8} \Omega_{\mathbb{D}}$. They remarked that their algebraic detection theorem can be modified to prove that the C_{2^n} -fixed points of $MU^{((C_{2^n}))}$, for $n \geq 3$, detect the Kervaire classes. It's worth pointing out the differences between our algebraic detection theorem and their algebraic detection theorem. To prove their detection theorem, Hill, Hopkins, and Ravenel used the map of spectral sequences

$$\begin{array}{ccc}
 \text{Adams–Novikov spectral sequence} & \xrightarrow{\quad\quad\quad} & \pi_* \mathbb{S} \\
 \downarrow & & \downarrow \\
 C_{2^n}\text{-homotopy fixed point spectral sequence} & \xrightarrow{\quad\quad\quad} & \pi_*(MU^{((C_{2^n}))})^{C_{2^n}}.
 \end{array}$$

Their algebraic detection theorem [18, Theorem 11.2] shows that if $x \in \text{Ext}_{MU_* MU}^{2, 2^{j+1}} \times (MU_*, MU_*)$ is any element mapping to h_j^2 on the E_2 -page of the classical Adams spectral sequence, then the image of x in $H^2(C_{2^n}, \pi_{2^{j+1}}^u MU^{((C_{2^n}))})$ is not zero. Once this is proved, their detection theorem follows easily.

They further remarked that their algebraic detection theorem does not hold when G is C_2 or C_4 (see [18, Remark 11.14]). For these groups, there is a jump of filtration. In particular, for $n = 1$, the element $x \in \text{Ext}_{MU_* MU}^{2, 2^{j+1}}(MU_*, MU_*)$ maps to 0 on the E_2 -page of the C_2 -homotopy fixed point spectral sequence of $MU_{\mathbb{R}}$. However, because of Theorem 1.6, we deduce that there must be a nontrivial extension so that x actually corresponds to an element of filtration $2^{j+1} - 2$ in the C_2 -homotopy fixed point spectral sequence. For our algebraic detection theorem, this jump of filtration does not occur because we used maps of Adams spectral sequences.

As an application of Theorem 1.2, we study Hurewicz images of Real Johnson–Wilson theories. The Real Johnson–Wilson theories $E\mathbb{R}(n)$ were first constructed and studied by Hu and Kriz [20]. They constructed $E\mathbb{R}(n)$ from $BP_{\mathbb{R}}$ by mimicking the classical construction of $E(n)$. More precisely, there is an isomorphism

$$\mathbb{Z}[v_1, v_2, \dots] = \pi_{2*} BP \cong \pi_{*\rho_2}^{C_2} BP_{\mathbb{R}} = \mathbb{Z}[\bar{v}_1, \bar{v}_2, \dots],$$

where $\bar{v}_i \in \pi_{i\rho_2}^{C_2} BP_{\mathbb{R}}$ are lifts of the classical generators $v_i \in \pi_{2i} BP$. Quotienting out the \bar{v}_i generators for all $i \geq n + 1$ and inverting the \bar{v}_n produces the Real Johnson–Wilson

theory $E\mathbb{R}(n)$. It is a C_2 -equivariant spectrum whose underlying spectrum is $E(n)$, with an C_2 -action induced from the complex conjugation action of $BP_{\mathbb{R}}$.

Many people have also studied $E\mathbb{R}(n)$ after Hu and Kriz. Kitchloo and Wilson [22] proved that the fixed points $ER(n) := E\mathbb{R}(n)^{C_2}$ fits into the fiber sequence

$$\Sigma^{\lambda(n)}ER(n) \rightarrow ER(n) \rightarrow E(n),$$

where $\lambda(n) = 2^{2n+1} - 2^{n+2} + 1$. When $n = 1$, $E\mathbb{R}(1)$ is Atiyah’s Real K -theory $K\mathbb{R}$, with $ER(1) = K\mathbb{R}^{C_2} = KO$. In this case, Kitchloo and Wilson’s fibration recovers the classical fibration

$$\Sigma KO \xrightarrow{\eta} KO \rightarrow KU.$$

When $n = 2$, using the Bockstein spectral sequence associated to the fibration, Kitchloo and Wilson subsequently computed the cohomology groups $ER(2)^*(\mathbb{R}P^n)$ and $ER(2)^*(\mathbb{R}P^n \wedge \mathbb{R}P^m)$. From their computation, they deduced new nonimmersion results for even dimensional real projective spaces [23,24]. Most recently, Kitchloo, Lorman, and Wilson have used this Bockstein spectral sequence to further compute the $ER(n)$ cohomology of other spaces as well [30,25,26].

In [15], Hill and Meier studied the spectra $TMF_1(3)$ and $Tmf_1(3)$ of topological modular forms at level three. They proved that the spectrum $tmf_1(3)$, considered as an C_2 -equivariant spectrum, is a form of $BP_{\mathbb{R}}(2)$, and $tmf_1(3)[\bar{a}_3^{-1}]$ is a form of $E\mathbb{R}(2)$. Using this identification, they computed the C_2 -equivariant Picard groups and the C_2 -equivariant Anderson dual of $Tmf_1(3)$.

We are interested in the Hurewicz images of $\pi_*E\mathbb{R}(n)^{C_2}$. To do so, we study the map of slice spectral sequences

$$\text{SliceSS}(BP_{\mathbb{R}}) \rightarrow \text{SliceSS}(E\mathbb{R}(n)).$$

Theorem 1.5 and Theorem 1.6 identify the classes in the slice spectral sequence of $BP_{\mathbb{R}}$ that detect the families $\{h_i\}$, $\{h_j^2\}$, and $\{g_k\}$. Analyzing the images of these classes in the slice spectral sequence of $E\mathbb{R}(n)$ produces the detection theorem for $E\mathbb{R}(n)$.

Theorem 1.8 (*Detection theorem for $E\mathbb{R}(n)$*).

- (1) For $1 \leq i, j \leq n$, if the element $h_i \in \text{Ext}_{\mathcal{A}_*}^{1,2^i}(\mathbb{F}_2, \mathbb{F}_2)$ or $h_j^2 \in \text{Ext}_{\mathcal{A}_*}^{2,2^{j+1}}(\mathbb{F}_2, \mathbb{F}_2)$ survives to the E_{∞} -page of the Adams spectral sequence, then its image under the Hurewicz map $\pi_*\mathbb{S} \rightarrow \pi_*E\mathbb{R}(n)^{C_2}$ is nonzero.
- (2) For $1 \leq k \leq n-1$, if the element $g_k \in \text{Ext}_{\mathcal{A}_*}^{4,2^{k+2}+2^{k+3}}(\mathbb{F}_2, \mathbb{F}_2)$ survives to the E_{∞} -page of the Adams spectral sequence, then its image under the Hurewicz map $\pi_*\mathbb{S} \rightarrow \pi_*E\mathbb{R}(n)^{C_2}$ is nonzero.

Theorem 1.8 is extremely useful for computing $ER(n)^*(\mathbb{R}P^m)$. In [28], we use the fact that the Hopf elements are detected by $\pi_*^{C_2}E\mathbb{R}(n)$ to deduce the compatibility of

the slice differentials of $E\mathbb{R}(n)$ and the attaching maps of $\mathbb{R}P^m$. As a result, we are able to compute $ER(2)^*(\mathbb{R}P^m)$ by a double filtration spectral sequence, solving all the 2-extensions and some η and ν -extensions.

Hahn and the second author have shown that the Lubin–Tate theories E_n , equipped with the Goerss–Hopkins–Miller C_2 -action ([35,10]), is Real oriented. In other words, there is a C_2 -equivariant map $MU_{\mathbb{R}} \rightarrow E_n$. The proof for Theorem 1.8 can be modified to prove Hurewicz images for the homotopy fixed point spectra E_n^{hG} . In [14], the authors show that the Hurewicz images of $E\mathbb{R}(n)^{C_2}$ and $E_n^{hC_2}$ are the same. It follows that Theorem 1.8 holds for $\pi_*E_n^{hC_2}$ as well.

1.2. Summary of the contents

In Section 2, we provide the necessary background for the C_2 -equivariant dual Steenrod algebras — $(H\underline{\mathbb{F}}_{2\star}, \mathcal{A}_{\star}^m)$ and $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$ — and their C_2 -equivariant Adams spectral sequences. In Section 3, we compute the slice spectral sequence and the homotopy fixed point spectral sequence of $BP_{\mathbb{R}}$. In Section 4, we construct the equivariant May spectral sequence of $BP_{\mathbb{R}}$ and prove Theorem 1.6. In Section 5, we modify the filtration of the classical dual Steenrod algebra \mathcal{A}_{\star} to obtain a compatible filtration with respect to the map $\mathcal{A}_{\star} \rightarrow \mathcal{A}_{\star}^m \rightarrow \mathcal{A}_{\star}^{cc}$ of Steenrod algebras. We then analyze the resulting maps of May spectral sequences. Lastly, in Section 6, we combine results from the previous sections and prove Theorem 1.2, Corollary 1.3, Theorem 1.5, and Theorem 1.8.

1.3. Acknowledgments

The authors would like to thank the organizers of the 2016 Talbot workshop, Eva Belmont, Inbar Klang, and Dylan Wilson, for inviting them to the workshop. This project would not have come into being without the mentorship of Mike Hill and Doug Ravenel during the workshop. We would like to thank Vitaly Lorman for helpful conversations and a fruitful exchange of ideas. We are also grateful to Hood Chatham for his spectral sequence package, which produced all of our diagrams. Thanks are also due to Mark Behrens, Jeremy Hahn, Achim Krause, Peter May, Haynes Miller, Eric Peterson, Doug Ravenel, David B Rush, and Mingcong Zeng for helpful conversations. Finally, we would like to heartily thank Mike Hill and Mike Hopkins for sharing numerous insights with us during various stages of the project and many helpful conversations. The fourth author was partially supported by the National Science Foundation under Grant No. DMS-1810638.

2. The equivariant dual Steenrod algebra and Adams spectral sequence

In this section, we provide the necessary background for the C_2 -equivariant dual Steenrod algebra and the C_2 -equivariant Adams spectral sequence. These have been extensively studied by Hu–Kriz [20] and Greenlees [11–13]. Of the many ways to define

the C_2 -equivariant dual Steenrod algebra, two of them are of interest to us. The first one is the **Borel equivariant dual Steenrod algebra**

$$\mathcal{A}_\star^{cc} := F(EC_{2+}, H\mathbb{F}_2 \wedge H\mathbb{F}_2)_\star.$$

This has been studied by Greenlees. The second one is the **genuine equivariant dual Steenrod algebra**. It is defined by using the genuine Eilenberg–Mac Lane spectrum $H\underline{\mathbb{F}}_2$:

$$\mathcal{A}_\star^m := (H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2)_\star.$$

2.1. \mathcal{A}_\star^{cc} and \mathcal{A}_\star^m

To compute \mathcal{A}_\star^m , Hu and Kriz first computed the $RO(C_2)$ -graded homotopy groups \mathcal{A}_\star^{cc} . This computation can be further used to deduce the $RO(C_2)$ -graded homotopy groups \mathcal{A}_\star^m . We give a brief summary of Hu and Kriz’s computation of \mathcal{A}_\star^{cc} and \mathcal{A}_\star^m , focusing on the parts that we will need again for the later sections. For more details of their computation, see Section 6 of [20].

To start, we need the coefficient rings of the C_2 -equivariant Eilenberg–Mac Lane spectra $H^c := F(EC_{2+}, H\mathbb{F}_2)$ and $H^m := H\underline{\mathbb{F}}_2$. The following are some distinguished elements in their $RO(C_2)$ -graded homotopy groups.

Definition 2.1. The element

$$a_\sigma \in \pi_{-\sigma}^{C_2} S^0$$

is the element corresponding to the inclusion $S^0 \hookrightarrow S^\sigma$ (the one point compactification of the inclusion $\{0\} \subset \sigma$) under the suspension isomorphism $\pi_{-\sigma}^{C_2} S^0 \cong \pi_0^{C_2} S^\sigma$. Under the Hurewicz maps $\pi_\star^{C_2} \mathbb{S} \rightarrow H_\star^c$ and $\pi_\star^{C_2} \mathbb{S} \rightarrow H_\star^m$, the images of a_σ are nonzero. By an abuse of notation, we will denote the images by a_σ as well.

Definition 2.2. The element

$$u_\sigma \in \pi_{1-\sigma}^{C_2} H\underline{\mathbb{F}}_2$$

is the element corresponding to the generator of $H_1^{C_2}(S^\sigma; \mathbb{F}_2) = \pi_1^{C_2}(S^\sigma \wedge H\underline{\mathbb{F}}_2)$. It can also be regarded as an element in $\pi_{1-\sigma}^{C_2} F(EC_{2+}, H\mathbb{F}_2)$ via the map

$$H\underline{\mathbb{F}}_2 \rightarrow F(EC_{2+}, H\underline{\mathbb{F}}_2) \simeq F(EC_{2+}, H\mathbb{F}_2).$$

Hu and Kriz first computed H_\star^c . They then used it to analyze the cofiber of the map

$$H\underline{\mathbb{F}}_2 \rightarrow F(EC_{2+}, H\underline{\mathbb{F}}_2) \simeq F(EC_{2+}, H\mathbb{F}_2)$$

and subsequently computed the coefficient ring $H\underline{\mathbb{F}}_{2\star}$.




Fig. 1. The coefficient rings of $H\mathbb{F}_2$ (left) and $F(EC_{2+}, H\mathbb{F}_2)$ (right). The map $H\mathbb{F}_2 \rightarrow F(EC_{2+}, H\mathbb{F}_2)$ induces an isomorphism in the range $p \geq 0$.

Proposition 2.3 (Hu–Kriz).

(1) The coefficient ring $H_\star = F(EC_{2+}, H\mathbb{F}_2)_\star$ is the polynomial algebra

$$F(EC_{2+}, H\mathbb{F}_2)_{p+q\sigma} = \mathbb{F}_2[u_\sigma^\pm, a_\sigma].$$

(2) The coefficient ring $H_\star^m := H\mathbb{F}_2_\star^m$ is

$$H\mathbb{F}_2_{p+q\sigma} = \mathbb{F}_2[u_\sigma, a_\sigma] \oplus \mathbb{F}_2 \left\{ \frac{\theta}{u_\sigma^i a_\sigma^j} \right\}, \quad i, j \geq 0,$$

where θ is an element in $\pi_{2\sigma-2}^{C_2} H\mathbb{F}_2$. The element θ is infinitely u_σ and a_σ -divisible. It is also u_σ and a_σ -torsion. The product of any two elements $x, y \in \mathbb{F}_2 \left\{ \frac{\theta}{u_\sigma^i a_\sigma^j} \right\}$ is 0.

In particular, the map $H_{p+q\sigma}^m \rightarrow H_{p+q\sigma}^c$ is an isomorphism in the range $a \geq 0$. Fig. 1 shows $H\mathbb{F}_2$ and $F(EC_{2+}, H\mathbb{F}_2)$.

Remark 2.4. In [20], Hu and Kriz denoted u_σ by σ^{-1} and a_σ by a .

Remark 2.5. The element $\theta \in H_\star^m$ can be defined as follows: consider the Tate diagram

$$\begin{array}{ccccc} EC_{2+} \wedge H\mathbb{F}_2 & \longrightarrow & H\mathbb{F}_2 & \longrightarrow & \widetilde{EC}_2 \wedge H\mathbb{F}_2 \\ \downarrow \simeq & & \downarrow & & \downarrow \\ EC_{2+} \wedge F(EC_{2+}, H\mathbb{F}_2) & \longrightarrow & F(EC_{2+}, H\mathbb{F}_2) & \longrightarrow & \widetilde{EC}_2 \wedge F(EC_{2+}, H\mathbb{F}_2). \end{array}$$

Taking $\pi_{\star}^{C_2}(-)$ produces the following diagram on homotopy groups

$$\begin{CD} H_{\star}^f @>>> H_{\star}^m @>>> H_{\star}^g \\ @V \cong VV @VVV @VVV \\ H_{\star}^f @>>> H_{\star}^c @>>> H_{\star}^t \end{CD}$$

The coefficient rings of H_{\star}^t and H_{\star}^f can be computed to be

$$\begin{aligned} H_{\star}^t &= \mathbb{F}_2[u_{\sigma}^{\pm}, a_{\sigma}^{\pm}] \\ H_{\star}^f &= \mathbb{F}_2[u_{\sigma}^{\pm}, a_{\sigma}^{-1}]. \end{aligned}$$

Now, consider the boundary map ∂ defined by using the long exact sequences of homotopy groups for the top and bottom rows of the Tate diagram:

$$\partial : H_{\star}^t \rightarrow H_{\star-1}^f \rightarrow H_{\star-1}^m.$$

The element θ is the image of $u_{\sigma}^{-1}a_{\sigma}^{-1} \in H_{\star}^t$ under the boundary map ∂ .

With these coefficient groups in hand, we are now ready to compute the equivariant dual Steenrod algebras. When computing \mathcal{A}_{\star}^{cc} , we need to work in the category \mathcal{M} of bigraded $\mathbb{Z}[a_{\sigma}]$ -modules that are complete with respect to the topology associated with the principal ideal (a_{σ}) . The morphisms in this category are continuous homomorphisms. It turns out that even though \mathcal{A}_{\star}^{cc} is not flat over H_{\star}^c as \mathbb{F}_2 -modules, completion by (a_{σ}) ensures that \mathcal{A}_{\star}^{cc} is flat over H_{\star}^c in the category \mathcal{M} . Thus, we can regard $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$ as a \mathcal{M} -Hopf algebroid.

Theorem 2.6 (Hu–Kriz). *The \mathcal{M} -Hopf algebroid $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$ can be described by the following structure formulas:*

- (1) $\mathcal{A}_{\star}^{cc} = H_{\star}^c[\zeta_i \mid i \geq 1]_{a_{\sigma}}^{\wedge}$, $\dim \zeta_i = 2^i - 1$;
- (2) $\psi(\zeta_i) = \sum_{0 \leq j \leq i} \zeta_{i-j}^{2^j} \otimes \zeta_j$, with $\zeta_0 = 1$;
- (3) $\eta_R(u_{\sigma}^{-1}) = \sum_{i \geq 0} (u_{\sigma}^{-1})^{2^i} \zeta_i a_{\sigma}^{2^i - 1}$;
- (4) $\eta_R(a_{\sigma}) = a_{\sigma}$.

The formula for \mathcal{A}_{\star}^{cc} in Theorem 2.6 is obtained from the $RO(C_2)$ -graded homotopy fixed point spectral sequence

$$H^s(C_2; \pi_t(H\mathbb{F}_2 \wedge H\mathbb{F}_2) \otimes \text{sgn}^{\otimes r}) \implies \pi_{t-s+(\sigma-1)r} F(EC_{2+}, H\mathbb{F}_2 \wedge H\mathbb{F}_2),$$

where sgn is the sign representation. The ζ_i generators in \mathcal{A}_\star^{cc} are images of the ζ_i generators in the classical dual Steenrod algebra $\mathcal{A}_\star = \mathbb{F}_2[\zeta_i \mid i \geq 1]$ under the map

$$H\mathbb{F}_2 \wedge H\mathbb{F}_2 \rightarrow F(EC_{2+}, H\mathbb{F}_2 \wedge H\mathbb{F}_2).$$

Although Theorem 2.6 provides formulas describing $(H_\star^c, \mathcal{A}_\star^{cc})$ as a \mathcal{M} -Hopf algebroid, it is not very helpful for computing the Hopf algebroid structure of $(H_\star^m, \mathcal{A}_\star^m)$. To further compute \mathcal{A}_\star^m , Hu and Kriz constructed explicit equivariant generators ξ_i and τ_i in both \mathcal{A}_\star^m and \mathcal{A}_\star^{cc} . These generators are compatible in the sense that under the map $\mathcal{A}_\star^m \rightarrow \mathcal{A}_\star^{cc}$, $\tau_i \mapsto \tau_i$ and $\xi_i \mapsto \xi_i$. By computing the relations between the ξ_i 's and τ_i 's, Hu and Kriz obtained an alternative description of the \mathcal{M} -Hopf algebroid structure of $(H_\star^c, \mathcal{A}_\star^{cc})$. Afterwards, they observed that the exact same relations hold in \mathcal{A}_\star^m as well. This observation ultimately led them to conclude the Hopf algebroid structure of $(H\mathbb{F}_2, \mathcal{A}_\star^m)$.

We now introduce the ξ_i and τ_i generators. We structure our exposition to focus on describing the map

$$\mathcal{A}_\star \rightarrow \mathcal{A}_\star^m.$$

Understanding this map will be of great importance to us later on.

Definition 2.7. For X an C_2 -equivariant spectrum, let

$$\begin{aligned} H_\star^{cc} X &:= F(EC_{2+}, H\mathbb{F}_2 \wedge X)_\star, \\ H_\star^m X &:= (H\mathbb{F}_2 \wedge X)_\star. \end{aligned}$$

Classically, if a spectrum E is complex oriented, then one can easily compute E_*BP as follows: choose a complex orientation $b \in E^2(\mathbb{C}P^\infty)$. Associated to b is a coproduct formula

$$\psi(b) = \sum_{i \geq 0}^F b^{2^i} \otimes \xi_i,$$

where F is formal sum induced by the complex orientation of E . From this coproduct formula, one is led to conclude that

$$E_*BP = E_*[\xi_i \mid i \geq 1], \quad |\xi_i| = 2(2^i - 1).$$

This argument works C_2 -equivariantly as well. The genuine Eilenberg–Mac Lane spectrum $H\mathbb{F}_2$ is Real oriented via the Thom map $BP_{\mathbb{R}} \rightarrow H\mathbb{F}_2$. Applying the argument above produces equivariant polynomial generators for $H_\star^{cc}BP_{\mathbb{R}}$ and $H_\star^mBP_{\mathbb{R}}$.

Proposition 2.8 (Hu–Kriz). *There exist generators ξ_i of dimensions $|\xi_i| = (2^i - 1)\rho_2$ in both $H_\star^mBP_{\mathbb{R}}$ and $H_\star^{cc}BP_{\mathbb{R}}$, such that*

$$H_{\star}^m BP_{\mathbb{R}} = H_{\star}^m[\xi_i \mid i \geq 1],$$

$$H_{\star}^{cc} BP_{\mathbb{R}} = H_{\star}^c[\xi_i \mid i \geq 1].$$

Furthermore, the two sets of ξ_i generators are compatible in the sense that the map

$$H\underline{\mathbb{F}}_2 \wedge BP_{\mathbb{R}} \rightarrow F(EC_{2+}, H\underline{\mathbb{F}}_2 \wedge BP_{\mathbb{R}}) = F(EC_{2+}, H\underline{\mathbb{F}}_2 \wedge BP_{\mathbb{R}})$$

induces the map

$$H_{\star}^m BP_{\mathbb{R}} \rightarrow H_{\star}^{cc} BP_{\mathbb{R}}$$

sending $\xi_i \mapsto \xi_i$.

Definition 2.9. The orientation map $f : BP_{\mathbb{R}} \rightarrow H\underline{\mathbb{F}}_2$ induces the commutative diagram

$$\begin{array}{ccc} H\underline{\mathbb{F}}_2 \wedge BP_{\mathbb{R}} & \xrightarrow{\text{id} \wedge f} & H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2 \\ \downarrow & & \downarrow \\ F(EC_{2+}, H\underline{\mathbb{F}}_2 \wedge BP_{\mathbb{R}}) & \longrightarrow & F(EC_{2+}, H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2), \end{array}$$

which, after taking equivariant homotopy groups $\pi_{\star}^{C_2}(-)$, becomes

$$\begin{array}{ccc} H_{\star}^m BP_{\mathbb{R}} & \longrightarrow & \mathcal{A}_{\star}^m \\ \downarrow & & \downarrow \\ H_{\star}^{cc} BP_{\mathbb{R}} & \longrightarrow & \mathcal{A}_{\star}^{cc}. \end{array}$$

The image of the ξ_i generators in Proposition 2.8 produces generators $\xi_i \in \mathcal{A}_{\star}^m$ and \mathcal{A}_{\star}^{cc} .

Consider the commutative diagram

$$\begin{array}{ccc} H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2 = H\underline{\mathbb{F}}_2^{C_2} \wedge H\underline{\mathbb{F}}_2^{C_2} & \longrightarrow & (H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2)^{C_2} \\ & \searrow & \downarrow \\ & & F(EC_{2+}, H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2)^{C_2} \\ & & \parallel \\ & & F(EC_{2+}, H\underline{\mathbb{F}}_2 \wedge H\underline{\mathbb{F}}_2)^{C_2}. \end{array}$$

Taking $\pi_{\star}(-)$ produces the diagram

$$\begin{array}{ccc} \mathcal{A}_{\star} & \longrightarrow & \mathcal{A}_{\star}^m \\ & \searrow & \downarrow \\ & & \mathcal{A}_{\star}^{cc}. \end{array}$$

Here, \mathcal{A}_*^m and \mathcal{A}_*^{cc} are the integer graded parts of \mathcal{A}_*^m and \mathcal{A}_*^{cc} , respectively. The following theorem provides formulas relating the ξ_i generators and images of the ζ_i generators under the maps $\mathcal{A}_* \rightarrow \mathcal{A}_*^m \rightarrow \mathcal{A}_*^{cc}$.

Theorem 2.10 (Relations between ξ_i and ζ_i).

- (1) $\psi(\xi_i) = \sum_{0 \leq j \leq i} \xi_{i-j}^{2^j} \otimes \xi_j$;
- (2) The ξ_i generators are related to the images of the ζ_i generators (which, by an abuse of notation, will also be denoted by ζ_i) by the recursion formulas

$$\begin{aligned} \xi_0 &= 1 \\ a_\sigma^{2^i} \xi_i &= \zeta_{i-1}^2 \eta_R(u_\sigma) + \zeta_i a_\sigma + \xi_{i-1} u_\sigma^{2^{i-1}}, \quad i \geq 1. \end{aligned} \tag{2.1}$$

Proof. We prove the relations in \mathcal{A}_*^m . Once we have proven that they hold in \mathcal{A}_*^m , they will automatically hold in \mathcal{A}_*^{cc} as well. The proof is essentially the same as the proof of Theorem 6.18 in [20]. Let $b \in H\mathbb{F}_2^{\rho^2}(BS_+^1)$ be the Real orientation and $r \in H\mathbb{F}_2^1(B\mathbb{Z}/2_+)$ be the generator of $H\mathbb{F}_2^\star(B\mathbb{Z}/2_+)$. The coproduct formulas for b and r are, by definition,

$$\begin{aligned} \psi(b) &= \sum_{i \geq 0} b^{2^i} \otimes \xi_i \\ \psi(r) &= \sum_{i \geq 0} r^{2^i} \otimes \zeta_i. \end{aligned}$$

Part (1) is obtained by computing $\psi(\psi(b))$ in two ways through the commutative diagram

$$\begin{CD} H\mathbb{F}_2^\star(BS_+^1) @>\psi>> H\mathbb{F}_2^\star(BS_+^1) \wedge \mathcal{A}_*^m \\ @V\psi VV @VV\text{id} \wedge \psi V \\ H\mathbb{F}_2^\star(BS_+^1) \wedge \mathcal{A}_*^m @>\psi \wedge \text{id}>> H\mathbb{F}_2^\star(BS_+^1) \wedge \mathcal{A}_*^m \wedge \mathcal{A}_*^m \end{CD}$$

and comparing the coefficients of b^{2^i} :

$$\sum_{i \geq 0} \psi(b)^{2^i} \otimes \xi_i = \psi(\psi(b)) = \sum_{i \geq 0} b^{2^i} \otimes \psi(\xi_i).$$

For part (2), the map $B\mathbb{Z}/2_+ \rightarrow BS_+^1$ induces the map

$$H\mathbb{F}_2^\star(BS_+^1) \rightarrow H\mathbb{F}_2^\star(B\mathbb{Z}/2_+)$$

on equivariant cohomologies. This is a map

$$H\mathbb{F}_2^\star[[b]] \rightarrow H\mathbb{F}_2^\star[[r]],$$

where $|b| = -\rho_2$, $|r| = -1$. We would like to express the image of b in terms of r . The only terms on the right hand side that are of degree $-\rho_2$ are r^2u_σ and ra_σ . Hu and Kriz show that b maps to the sum of these two terms:

$$b \mapsto r^2u_\sigma + ra_\sigma.$$

The commutative diagram

$$\begin{CD} H\underline{\mathbb{F}}_2^\star(BS^1_+) @>f>> H\underline{\mathbb{F}}_2^\star(B\mathbb{Z}/2_+) \\ @VV\psi V @VV\psi V \\ H\underline{\mathbb{F}}_2^\star(BS^1_+) \wedge \mathcal{A}_\star^m @>f \wedge \text{id}>> H\underline{\mathbb{F}}_2^\star(B\mathbb{Z}/2_+) \wedge \mathcal{A}_\star^m, \end{CD}$$

obtained by the naturality of the coproduct, implies that

$$\sum_{i \geq 0} (r^2u_\sigma + ra_\sigma)^{2^i} \otimes \xi_i = \psi(b) = \psi(r^2u_\sigma + ra_\sigma) = \eta_R(u_\sigma) \sum_{i \geq 0} r^{2^{i+1}} \otimes \zeta_i^2 + a_\sigma \sum_{i \geq 0} r^{2^i} \otimes \zeta_i.$$

Comparing coefficients of r^{2^i} on both sides produces the recursion formulas, as desired. \square

We will now define the τ_i generators and compute their relations to the images of the classical ζ_i generators. Consider the C_2 -equivariant map $BS^1 \rightarrow BS^1$ classifying the squaring of Real line bundles. This produces the fiber sequence

$$B_{\mathbb{R}}\mathbb{Z}/2 \rightarrow BS^1 \xrightarrow{L^2} BS^1, \tag{2.2}$$

where the fiber $B_{\mathbb{R}}\mathbb{Z}/2$ is $\mathbb{R}P^\infty$, but with a nontrivial C_2 -action (the fixed point of $B_{\mathbb{R}}\mathbb{Z}/2$ under the C_2 action is $\mathbb{R}P^\infty \amalg \mathbb{R}P^\infty$). The Real orientation $b \in H\underline{\mathbb{Z}}^{\rho_2}(BS^1)$ restricts to a class $b' \in H\underline{\mathbb{Z}}^{\rho_2}(B_{\mathbb{R}}\mathbb{Z}/2)$. Under the map $H\underline{\mathbb{Z}} \rightarrow H\underline{\mathbb{F}}_2$, this gives a class $b'' \in H\underline{\mathbb{F}}_2^{\rho_2}(B_{\mathbb{R}}\mathbb{Z}/2)$:

$$\begin{CD} b \in H\underline{\mathbb{Z}}^{\rho_2}(\mathbb{C}P^\infty) @>>> b' \in H\underline{\mathbb{Z}}^{\rho_2}(B_{\mathbb{R}}\mathbb{Z}/2) \\ @VVV @VVV \\ b \in H\underline{\mathbb{F}}_2^{\rho_2}(\mathbb{C}P^\infty) @>>> b'' \in H\underline{\mathbb{F}}_2^{\rho_2}(B_{\mathbb{R}}\mathbb{Z}/2). \end{CD}$$

The composition

$$H\underline{\mathbb{Z}}^\star(BS^1) \rightarrow H\underline{\mathbb{Z}}^\star(BS^1) \rightarrow H\underline{\mathbb{Z}}^\star(B_{\mathbb{R}}\mathbb{Z}/2)$$

sends $b \mapsto 2b' = 0$. This implies that b' is in the image of the Bockstein β , induced by $H\underline{\mathbb{Z}} \xrightarrow{2} H\underline{\mathbb{Z}} \rightarrow H\underline{\mathbb{F}}_2$:

$$\cdots \rightarrow H\underline{\mathbb{F}}_2^\sigma(B_{\mathbb{R}}\mathbb{Z}/2) \xrightarrow{\beta} H\underline{\mathbb{Z}}^{\rho_2}(B_{\mathbb{R}}\mathbb{Z}/2) \xrightarrow{\cdot 2} H\underline{\mathbb{Z}}^{\rho_2}(B_{\mathbb{R}}\mathbb{Z}/2) \rightarrow \cdots$$

Let $c \in H\underline{\mathbb{F}}_2^\sigma(B_{\mathbb{R}}\mathbb{Z}/2)$ be a class such that $\beta c = b'$.

Proposition 2.11 (Hu–Kriz). $H\underline{\mathbb{F}}_2^\star(B_{\mathbb{R}}\mathbb{Z}/2)$ is a free $H\underline{\mathbb{F}}_2^\star$ -module with basis $\{1, b, b^2, \dots, c, cb, cb^2, \dots\}$.

Proof. Consider the cofiber sequence

$$B_{\mathbb{R}}\mathbb{Z}/2_+ \rightarrow BS_+^1 \rightarrow \text{Thom}(BS^1, L^2).$$

Taking $H\underline{\mathbb{F}}_2^\star(-)$ produces the Gysin sequence

$$H\underline{\mathbb{F}}_2^\star(\text{Thom}(BS^1, L^2)) \xrightarrow{0} H\underline{\mathbb{F}}_2^\star(BS_+^1) \rightarrow H\underline{\mathbb{F}}_2^\star(B_{\mathbb{R}}\mathbb{Z}/2_+).$$

By the Thom isomorphism theorem, $H\underline{\mathbb{F}}_2^\star(\text{Thom}(BS^1, L^2)) \cong H\underline{\mathbb{F}}_2^\star(BS_+^1)[x]$ as a free $H\underline{\mathbb{F}}_2^\star(BS_+^1)$ -module. The generator $x \in H\underline{\mathbb{F}}_2^{\rho_2}(\text{Thom}(BS^1, L^2))$ maps to $0 \in H\underline{\mathbb{F}}_2^\star(BS_+^1)$, and it is the image of $c \in H\underline{\mathbb{F}}_2^\sigma(B_{\mathbb{R}}\mathbb{Z}/2_+)$. It follows that as a $H\underline{\mathbb{F}}_2^\star$ -module,

$$H\underline{\mathbb{F}}_2^\star(B_{\mathbb{R}}\mathbb{Z}/2_+) \cong H\underline{\mathbb{F}}_2^\star(BS_+^1) \oplus H\underline{\mathbb{F}}_2^\star(BS_+^1)[c] = H\underline{\mathbb{F}}_2^\star\{1, b, b^2, \dots, c, cb, cb^2, \dots\}.$$

□

Since $\psi(x) = x \otimes 1$ and $c \mapsto x$, the coproduct formula for c must be of the form

$$\psi(c) = c \otimes 1 + \sum_{i \geq 0} (b'')^{2^i} \otimes \tau_i,$$

where τ_i are elements in \mathcal{A}_\star^m with dimensions $|\tau_i| = (2^i - 1)\rho_{C_2} + 1$.

Theorem 2.12 (Hu–Kriz).

- (1) $\psi(\tau_i) = \tau_i \otimes 1 + \sum_{0 \leq j \leq i} \xi_{i-j}^{2^j} \otimes \tau_j$.
- (2) The τ_i generators are related to the images of the ζ_i generators by the recursion formulas

$$\begin{aligned} a_\sigma \tau_0 &= \eta_R(u_\sigma) + u_\sigma, \\ a_\sigma^{2^i} \tau_i &= \tau_{i-1} u_\sigma^{2^{i-1}} + \zeta_i \eta_R(u_\sigma), \quad i \geq 1. \end{aligned}$$

Proof. The coproduct formula for b'' is the same as the one for b :

$$\psi(b'') = \sum_{i \geq 0} b''^{2^i} \otimes \xi_i.$$

Similar to Theorem 2.10, part (1) can be proved by computing $\psi(\psi(c))$ in two ways:

$$\psi(c) \otimes 1 + \sum_{i \geq 0} \psi(b'')^{2^i} \otimes \xi_i = \psi(\psi(c)) = c \otimes 1 \otimes 1 + \sum_{i \geq 0} b''^{2^i} \otimes \psi(\tau_i).$$

For part (2), similar to the proof of Theorem 2.10, we consider the map on cohomology

$$H\underline{\mathbb{F}}_2^\star(B_{\mathbb{R}}\mathbb{Z}/2) \rightarrow H\underline{\mathbb{F}}_2^\star(\mathbb{R}\mathbb{P}^\infty) = H\underline{\mathbb{F}}_{2\star}[[r]].$$

This map is induced by the map $\mathbb{R}\mathbb{P}^\infty \hookrightarrow B_{\mathbb{R}}\mathbb{Z}/2^{C_2} \rightarrow B_{\mathbb{R}}\mathbb{Z}/2$. The image of $b'' \in H\underline{\mathbb{F}}_2^\sigma(B_{\mathbb{R}}\mathbb{Z}/2)$ is the same as the one we found in Theorem 2.10, $r^2u_\sigma + ra_\sigma$. To find the image of $c \in H\underline{\mathbb{F}}_2^\sigma(B_{\mathbb{R}}\mathbb{Z}/2)$, note that the only terms in $H\underline{\mathbb{F}}_{2\star}[[r]]$ of degree $-\sigma$ are $\{ru_\sigma, a_\sigma\}$. Hu and Kriz showed that depending on our choices, c can either map to ru_σ or $ru_\sigma + a_\sigma$. Assume that we have chosen c so that $c \mapsto ru_\sigma$ (the relationship between τ_i and ζ_i is going to be the same regardless of this choice). There are two ways to compute $\psi(c)$. On one hand,

$$\psi(c) = c \otimes 1 + \sum_{i \geq 0} b''^{2^i} \otimes \tau_i = (ru_\sigma) \otimes 1 + \sum_{i \geq 0} (r^2u_\sigma + ra_\sigma)^{2^i} \otimes \tau_i.$$

On the other hand,

$$\psi(c) = \psi(ru_\sigma) = \eta_R(u_\sigma)\psi(r) = \eta_R(u_\sigma) \sum_{i \geq 0} r^{2^i} \otimes \zeta_i.$$

Comparing the coefficients of r^{2^i} for both expressions produces the recursion formulas, as desired. \square

Remark 2.13. The proof above also shows that in the ring $H\underline{\mathbb{F}}_2^\star(B_{\mathbb{R}}\mathbb{Z}/2)$, there is the relation $c^2 = b''u_\sigma + ca_\sigma$, regardless of the choice of c .

Using the formulas in Theorem 2.10 and Theorem 2.12, one can show that in both \mathcal{A}_{\star}^{cc} and \mathcal{A}_{\star}^m , the ξ_i and τ_i generators are related by the formula

$$\tau_i^2 = \tau_{i+1}a_\sigma + \xi_{i+1}\eta_R(u_\sigma) \quad ([20, \text{Proposition 6.37}]).$$

This is the last ingredient needed to compute the Hopf algebroids $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$ and $(H_{\star}^m, \mathcal{A}_{\star}^m)$.

Theorem 2.14 (Corollary 6.40 and Theorem 6.41 in [20]).

(1) The \mathcal{M} -Hopf algebroid $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$ can be described by

$$\mathcal{A}_{\star}^{cc} = H_{\star}^c[\xi_i, \tau_i]/(\tau_0a_\sigma = u_\sigma + \eta_R(u_\sigma), \tau_i^2 = \tau_{i+1}a_\sigma + \xi_{i+1}\eta_R(u_\sigma)),$$

with comultiplications

- (a) $\psi(\xi_i) = \sum_{0 \leq j \leq i} \xi_{i-j}^{2^j} \otimes \xi_j;$
- (b) $\psi(\tau_i) = \tau_i \otimes 1 + \sum_{0 \leq j \leq i} \xi_{i-j}^{2^j} \otimes \tau_j.$

(2) The Hopf algebroid $(H_{\star}^m, \mathcal{A}_{\star}^m)$ can be described by

$$\mathcal{A}_{\star}^m = H\underline{\mathbb{F}}_{2\star}[\xi_i, \tau_i]/(\tau_0 a_{\sigma} = u_{\sigma} + \eta_R(u_{\sigma}), \tau_i^2 = \tau_{i+1} a_{\sigma} + \xi_{i+1} \eta_R(u_{\sigma})).$$

The comultiplications are the same as the ones in $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$. The right unit on the elements $\frac{\theta}{u_{\sigma}^i a_{\sigma}^j} \in H\underline{\mathbb{F}}_{2\star}$ is given by the formula

$$\eta_R\left(\frac{\theta}{u_{\sigma}^i a_{\sigma}^j}\right) = \partial\left(\frac{1}{(u_{\sigma} + \tau_0 a_{\sigma})^{i+1} a_{\sigma}^{j+1}}\right), \quad i \geq 0, j \geq 0,$$

where ∂ is the boundary map in Remark 2.5.

There are certain extensions involving the Hopf algebroids $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$ and $(H_{\star}^m, \mathcal{A}_{\star}^m)$ that will produce change of rings theorems. As we will see later, these change of rings theorems will greatly simplify the computation of the C_2 -equivariant May and Adams spectral sequences of $BP_{\mathbb{R}}$.

Let $P_{\star} = \mathbb{F}_2[\xi_i \mid i \geq 1]$. Then with the coproduct formula

$$\psi(\xi_i) = \sum_{0 \leq j \leq i} \xi_{i-j}^{2^j} \otimes \xi_j,$$

$(\mathbb{F}_2, P_{\star})$ is a \mathbb{F}_2 -Hopf algebra. Similarly, $(\mathbb{F}_2[a_{\sigma}], P_{\star}[a_{\sigma}])$ is a $\mathbb{F}_2[a_{\sigma}]$ -Hopf algebra.

Proposition 2.15 (Proposition 6.29 and Theorem 6.41(b) in [20]).

(1) There is an extension of \mathcal{M} -Hopf algebroids

$$(\mathbb{F}_2[a_{\sigma}], P_{\star}[a_{\sigma}]) \rightarrow (H_{\star}^c, \mathcal{A}_{\star}^{cc}) \rightarrow (H_{\star}^c, \Lambda_{\star}^{cc}),$$

where

$$\Lambda_{\star}^{cc} = H_{\star}^c[\tau_i]/(\tau_i^2 = \tau_{i+1} a_{\sigma}),$$

with structure formulas

- (a) τ_i are primitive;
- (b) $\eta_R(a_{\sigma}) = a_{\sigma};$
- (c) $\eta_R(u_{\sigma}) = u_{\sigma} + \tau_0 a_{\sigma}.$

(2) There is an extension of Hopf algebroids

$$(\mathbb{F}_2[a_{\sigma}], P_{\star}[a_{\sigma}]) \rightarrow (H_{\star}^m, \mathcal{A}_{\star}^m) \rightarrow (H_{\star}^m, \Lambda_{\star}^m),$$

where

$$\Lambda_{\star}^m = H\underline{\mathbb{F}}_{2\star}[\tau_i]/(\tau_i^2 = \tau_{i+1}a_{\sigma}).$$

The structure formulas for τ_i , a_{σ} , and u_{σ} are the same as the ones in $(H_{\star}^c, \Lambda_{\star}^{cc})$.

2.2. The equivariant Adams spectral sequence

We now introduce the equivariant Adams spectral sequences that are associated to the Hopf algebroids $(H\underline{\mathbb{F}}_{2\star}, \mathcal{A}_{\star}^m)$ and $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$, respectively.

Given a C_2 -equivariant spectrum X , we can resolve X by $H\underline{\mathbb{F}}_2$. The resulting resolution is the equivariant Adams resolution of X . The spectral sequence associated to this resolution is the C_2 -equivariant Adams spectral sequence associated to $(H\underline{\mathbb{F}}_{2\star}, \mathcal{A}_{\star}^m)$. Hu and Kriz observed that the equivariant Steenrod algebra \mathcal{A}_{\star}^m is a free $H\underline{\mathbb{F}}_{2\star}$ -module, hence flat over $H\underline{\mathbb{F}}_{2\star}$. From this, they concluded that the E_2 -page of the $(H\underline{\mathbb{F}}_{2\star}, \mathcal{A}_{\star}^m)$ -Adams spectral sequence can be identified as

$$\text{Ext}_{\mathcal{A}_{\star}^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star} X) \implies (\pi_{\star}^{C_2} X)_2^{\wedge}$$

(cf. [20, Corollary 6.47]). Similar to the classical Adams spectral sequence, the equivariant Adams spectral sequence will converge in nice cases. In particular, it will converge for X a finite C_2 -spectrum, $MU_{\mathbb{R}}$, or $BP_{\mathbb{R}}$.

On the other hand, by work of Greenlees [11–13], we can also form the classical Adams resolution of the underlying spectrum of X , and then apply the functor $F(EC_{2+}, -)$ to the classical Adams tower. The resulting spectral sequence associated to this new tower has E_2 -page

$$\text{Ext}_{\mathcal{A}_{\star}^{cc}}(H_{\star}^c, H_{\star}^{cc} X) \implies (\pi_{\star}^{C_2} F(EC_{2+}, X))_2^{\wedge}.$$

Again, this equivariant Adams spectral sequence will converge in our cases of interest.

By applying the functor $F(EC_{2+}, -)$ to the equivariant Adams resolution of X by $H\underline{\mathbb{F}}_2$, we produce a map of towers, hence a map between the two Adams spectral sequences

$$\begin{array}{ccc} \text{Ext}_{\mathcal{A}_{\star}^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star} X) & \implies & (\pi_{\star}^{C_2} X)_2^{\wedge} \\ \downarrow & & \downarrow \\ \text{Ext}_{\mathcal{A}_{\star}^{cc}}(H_{\star}^c, H_{\star}^{cc} X) & \implies & (\pi_{\star}^{C_2} F(EC_{2+}, X))_2^{\wedge}. \end{array}$$

On the E_2 -page, the map

$$\text{Ext}_{\mathcal{A}_{\star}^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star} X) \rightarrow \text{Ext}_{\mathcal{A}_{\star}^{cc}}(H_{\star}^c, H_{\star}^{cc} X)$$

is induced from the map $(H\underline{\mathbb{F}}_{2\star}, \mathcal{A}_{\star}^m) \rightarrow (H_{\star}^c, \mathcal{A}_{\star}^{cc})$ of Hopf algebroids.

When $X = BP_{\mathbb{R}}$, we can simplify the E_2 -pages of both Adams spectral sequences using Proposition 2.8 and Proposition 2.15:

$$\begin{aligned} \text{Ext}_{A_{\star}^m}(H\underline{\mathbb{F}}_2_{\star}, H\underline{\mathbb{F}}_2_{\star}BP_{\mathbb{R}}) &= \text{Ext}_{\Lambda_{\star}^m}(H\underline{\mathbb{F}}_2_{\star}, H\underline{\mathbb{F}}_2_{\star}), \\ \text{Ext}_{A_{\star}^{cc}}(H_{\star}^c, H_{\star}^{cc}BP_{\mathbb{R}}) &= \text{Ext}_{\Lambda_{\star}^{cc}}(H_{\star}^c, H_{\star}^c). \end{aligned}$$

Here, $\Lambda^m = H\underline{\mathbb{F}}_2 \wedge_{BP_{\mathbb{R}}} H\underline{\mathbb{F}}_2$ and $\Lambda^{cc} = F(EC_{2+}, H\underline{\mathbb{F}}_2 \wedge_{BP_{\mathbb{R}}} H\underline{\mathbb{F}}_2)$.

3. The slice spectral sequence of $BP_{\mathbb{R}}$

We will now discuss the slice spectral sequence and the homotopy fixed point spectral sequence of $BP_{\mathbb{R}}$.

3.1. The slice spectral sequence of $BP_{\mathbb{R}}$

For definitions and properties of the slice filtration, we refer the readers to [18, Section 4]. We will be interested in both the integer-graded and the $RO(C_2)$ -graded slice spectral sequence of $BP_{\mathbb{R}}$:

$$\begin{aligned} E_2^{s,t} &= \pi_{t-s}^{C_2} P_t^t BP_{\mathbb{R}} \implies \pi_{t-s}^{C_2} BP_{\mathbb{R}} \\ E_2^{s,V} &= \pi_{V-s}^{C_2} P_{\dim V}^{\dim V} BP_{\mathbb{R}} \implies \pi_{V-s}^{C_2} BP_{\mathbb{R}} \end{aligned}$$

The gradings are the Adams grading, with r^{th} -differentials $d_r : E_2^{s,t} \rightarrow E_2^{s+r,t+(r-1)}$ and $d_r : E_2^{s,V} \rightarrow E_2^{s+r,V+(r-1)}$, respectively.

To produce the E_2 -page of the slice spectral sequence, we compute the slice sections $P_t^t BP_{\mathbb{R}}$. Let $\bar{v}_i \in \pi_{(2^i-1)\rho_2} BP_{\mathbb{R}}$ be the equivariant lifts of the usual generators $v_i \in \pi_{2(2^i-1)} BP$. Using the method of twisted monoid rings [18, Section 2.4], we construct the A_{∞} -map

$$S^0[\bar{v}_1, \bar{v}_2, \dots] \rightarrow BP_{\mathbb{R}}.$$

This map has the property that after taking $\pi_*^u(-)$, it becomes an isomorphism. Using terminologies developed in [18], this map is a *multiplicative refinement* of $\pi_*^u BP_{\mathbb{R}}$. Furthermore, this multiplicative refinement produces the slice sections of $BP_{\mathbb{R}}$. The following result is a special case of the Slice Theorem ([18, Theorem 6.1]) applied to $BP_{\mathbb{R}}$.

Proposition 3.1. *The only nonzero slice sections of $BP_{\mathbb{R}}$ are $P_{2^n}^{2^n} BP_{\mathbb{R}}$, where $n \geq 0$. They are*

$$P_{2^n}^{2^n} BP_{\mathbb{R}} = \left(\bigvee_I S^{n\rho_2} \right) \wedge H\underline{\mathbb{Z}}$$




Fig. 2. Coefficient ring of $H\mathbb{Z}_{p+q\sigma}^{C_2}$. Multiplication by a_σ are drawn with solid lines and multiplication by $u_{2\sigma}$ are drawn with dashed lines (not all multiplicative structures are drawn).

where I is the indexing set consisting of all monomials of the form $\bar{v}_1^{i_1}\bar{v}_2^{i_2}\bar{v}_3^{i_3}\cdots$ with

$$n\rho_2 = (\rho_2)i_1 + (3\rho_2)i_2 + (7\rho_2)i_3 + \cdots .$$

Proposition 3.1 shows that computing the E_2 -page of the slice spectral sequence of $BP_{\mathbb{R}}$ can be reduced to computing the coefficient group $H\mathbb{Z}_{\star}^{C_2}$.

Definition 3.2 (The classes a_V and u_V). Let V be a representation of G with $\dim V = d$.

- (1) $a_V \in \pi_{-V}^G S^0$ is the map corresponding to the inclusion $S^0 \hookrightarrow S^V$ induced by $\{0\} \subset V$.
- (2) If V is oriented, $u_V \in \pi_{d-V}^G H\mathbb{Z}$ is the class corresponding to the generator of $H_d^G(S^V; H\mathbb{Z})$.

A comprehensive computation for the coefficient ring of $H\mathbb{Z}_{\star}^{C_2}$ can be found in [8].

Theorem 3.3 (Theorem 2.8 in [8]). Fig. 2 shows the coefficient ring $H\mathbb{Z}_{p+q\sigma}^{C_2}$. Its product structures are as follows:




Fig. 3. Important d_3 -differentials and surviving torsion classes in $\text{SliceSS}(BP_{\mathbb{R}})$.

- (1) In the range $p \geq 0$, $H\mathbb{Z}_{p+q\sigma}^{C_2}$ is the polynomial algebra $\mathbb{Z}[u_{2\sigma}, a_{\sigma}]/(2a_{\sigma})$.
- (2) In the range $p < 0$, the class $\alpha = \frac{2}{u_{2\sigma}} \in H_{-2+2\sigma}^{C_2}$ is killed by a_{σ} and is infinitely $u_{2\sigma}$ divisible; the class $\theta \in H_{-3+3\sigma}^{C_2}$ is killed by $u_{2\sigma}$ and a_{σ} and it is infinitely $u_{2\sigma}$ divisible and a_{σ} divisible.

Proposition 3.1 and Theorem 3.3 enable us to compute the E_2 -page of the $RO(C_2)$ -graded slice spectral sequence of $BP_{\mathbb{R}}$. In particular, the positive part is the polynomial algebra $\mathbb{Z}[\bar{v}_i, u_{2\sigma}, a_{\sigma}]/(2a_{\sigma})$ with

$$\begin{aligned} |\bar{v}_i| &= (0, (2^i - 1) + (2^i - 1)\sigma), \\ |u_{2\sigma}| &= (0, 2 - 2\sigma), \\ |a_{\sigma}| &= (1, 1 - \sigma). \end{aligned}$$

The E_2 -page of the integer graded slice spectral sequence is the sub-algebra consisting of all the elements that have integer degrees in $t-s$. It is concentrated in the first quadrant with a vanishing line of slope 1.

Proposition 3.4. *In the $RO(C_2)$ -grade slice spectral sequence for $\pi_{\star}^{C_2} BP_{\mathbb{R}}$, a_{σ} and \bar{v}_i are permanent cycles. The differentials $d_i(u_{2\sigma}^{2^{k-1}})$ are zero for $i < 2^{k+1} - 1$, and*

$$d_{2^{k+1}-1}(u_{2\sigma}^{2^{k-1}}) = \bar{v}_k a_{\sigma}^{2^{k+1}-1}.$$

Proof. This is a special case of Hill–Hopkins–Ravenel’s Slice Differential Theorem ([18, Theorem 9.9]), applied to when $G = C_2$. \square

In Figure 3–5, we draw the first three sets of differentials of the integer-graded slice spectral sequence. To organize this information in a clean way, we have disassembled the spectral sequence into “stages”, corresponding to the differentials d_3, d_7, d_{15}, \dots . At each stage, the important surviving torsion elements are shown. Many classes with low filtrations (i.e., those on the 0-line) are not drawn because they are not torsion, and hence won’t be important for the purpose of this paper.




Fig. 4. Important d_7 -differentials and surviving torsion classes in $\text{SliceSS}(BP_{\mathbb{R}})$.

3.2. $\text{SliceSS}(BP_{\mathbb{R}}) \rightarrow \text{HFPS}(BP_{\mathbb{R}})$

The homotopy fixed point spectral sequence of $BP_{\mathbb{R}}$ is also going to be useful to us. It is a spectral sequence that computes the C_2 -equivariant homotopy groups of $F(EC_{2+}, BP_{\mathbb{R}})$. The integer-graded homotopy fixed point spectral sequence for $BP_{\mathbb{R}}$ is

$$E_2^{s,t} = H^s(C_2, \pi_t^u BP_{\mathbb{R}}) \implies \pi_{t-s}^{C_2} F(EC_{2+}, BP_{\mathbb{R}}).$$

Just like the slice spectral sequence, there is also an $RO(C_2)$ -graded version of this, with E_2 -page

$$E_2^{s,V} = H^s(C_2; \pi_0(S^{-V} \wedge BP_{\mathbb{R}})) \implies \pi_{V-s}^{C_2} F(EC_{2+}, BP_{\mathbb{R}}).$$

For a more general discussion of the $RO(G)$ -graded homotopy fixed point spectral sequence for any equivariant G -spectrum X , see [15, Section 2.3]. By [15, Corollary 4.7], the E_2 -page of the $RO(C_2)$ -graded homotopy fixed point spectral sequence of $BP_{\mathbb{R}}$ is isomorphic to the polynomial algebra

$$\mathbb{Z}[\bar{v}_i, u_{2\sigma}^{\pm}, a_{\sigma}] / (2a_{\sigma}).$$




Fig. 5. Important d_{15} -differentials and surviving torsion classes in $\text{SliceSS}(BP_{\mathbb{R}})$.

The differentials are given by

$$\begin{aligned}
 d_{2^{k+1}-1}(u_{2\sigma}^{2^{k-1}}) &= \bar{v}_k a_{\sigma}^{2^{k+1}-1}, \\
 d_{2^{k+1}-1}(u_{2\sigma}^{-2^{k-1}}) &= d_{2^{k+1}-1}(u_{2\sigma}^{-2^k} \cdot u_{2\sigma}^{2^{k-1}}) \\
 &= u_{2\sigma}^{-2^k} d_{2^{k+1}-1}(u_{2\sigma}^{2^{k-1}}) \\
 &= \bar{v}_k u_{2\sigma}^{-2^k} a_{\sigma}^{2^{k+1}-1}.
 \end{aligned}
 \tag{3.1}$$

They can be obtained by equivariant primary cohomology operations (see [20, Lemma 3.34]). The readers might have noticed at this point that the differentials on the positive powers of $u_{2\sigma}$ are the same as the differentials in the slice spectral sequence. Indeed, there is a map $\text{SliceSS}(BP_{\mathbb{R}}) \rightarrow \text{HFPS}(BP_{\mathbb{R}})$ that induces an isomorphism in a certain range.

To explain this map of spectral sequences, we will first construct a map of towers. Let X be a C_2 -spectrum. Let $\mathcal{S}_{>n}^S$ denote the localizing subcategory generated by all the slice cells of dimension $> n$, and $\mathcal{S}_{>n}^P$ the localizing subcategory generated by all the spheres of dimension $> n$. When $n \geq 0$, $\mathcal{S}_{>n}^P \subseteq \mathcal{S}_{>n}^S$, and this gives a natural map of towers

$$\text{Post}^{\bullet}(X) \rightarrow P^{\bullet}(X)$$

from the Postnikov tower of X to the slice tower of X . Non-equivariantly, this map is an isomorphism, because the slice tower is the Postnikov tower when we forget the C_2 -action. It follows that after taking $F(EC_{2+}, -)$ to both towers, the horizontal map in the following diagram is an isomorphism:

$$\begin{array}{ccc}
 F(EC_{2+}, \text{Post}^\bullet(X)) & \xrightarrow{\cong} & F(EC_{2+}, P^\bullet(X)) \\
 & & \uparrow \\
 & & P^\bullet(X).
 \end{array}$$

The top-left tower, $F(EC_{2+}, \text{Post}^\bullet(X))$, is the tower for constructing the homotopy fixed point spectral sequence. It follows that the vertical map induces a map of $RO(C_2)$ -graded spectral sequences:

$$\text{SliceSS}(X) \rightarrow \text{HFPSS}(X).$$

Proposition 3.5. *When $X = BP_{\mathbb{R}}$, the map $\text{SliceSS}(BP_{\mathbb{R}}) \rightarrow \text{HFPSS}(BP_{\mathbb{R}})$, considered as a map of integer graded spectral sequences, induces an isomorphism on the E_2 -page on or below the line of slope 1.*

Proof. The map of sections

$$P_{2t}^{2t}BP_{\mathbb{R}} \rightarrow F(EC_{2+}, P_{2t}^{2t}BP_{\mathbb{R}})$$

is the map $X \rightarrow F(EC_{2+}, X)$ induced by the collapse $EC_{2+} \rightarrow S^0$. To prove the desired isomorphism, it suffices to show that the map

$$\pi_{2t-s}^{C_2}(S^{t\rho_2} \wedge H\mathbb{Z}) \rightarrow \pi_{2t-s}^{C_2}F(EC_{2+}, S^{t\rho_2} \wedge H\mathbb{Z})$$

is an isomorphism for all $2t - s \geq s$, or $t \geq s$. This is equivalent to showing that the map

$$\pi_{(t-s)-t\sigma}^{C_2}H\mathbb{Z} \rightarrow \pi_{(t-s)-t\sigma}^{C_2}F(EC_{2+}, H\mathbb{Z}) \cong \pi_{(t-s)-t\sigma}^{C_2}F(EC_{2+}, H\mathbb{Z})$$

is an isomorphism for all $t \geq s \geq 0$, which is true by Lemma 3.6. \square

Lemma 3.6. *The coefficient ring $\pi_{\star}^{C_2}F(EC_{2+}, H\mathbb{Z})$ is the polynomial algebra*

$$\mathbb{Z}[u_{2\sigma}^\pm, a_\sigma]/(2a_\sigma).$$

The map

$$\pi_{p+q\sigma}^{C_2}H\mathbb{Z} \rightarrow \pi_{p+q\sigma}^{C_2}F(EC_{2+}, H\mathbb{Z}) \cong \pi_{p+q\sigma}^{C_2}F(EC_{2+}, H\mathbb{Z}),$$

is an isomorphism when $p \geq 0$, sending $u_{2\sigma} \mapsto u_{2\sigma}$, $a_\sigma \mapsto a_\sigma$, and zero when $p < 0$.

Proof. This is a standard computation. We refer the readers to [15, Corollary 4.7] and [8, Appendix B] for more details than what is written here. The key observation is that computing $\pi_{\star}^{C_2} F(EC_{2+}, H\mathbb{Z})$ is equivalent to computing $H^*(C_2; \oplus_{r \in \mathbb{Z}} \text{sgn}^{\otimes r})$, where sgn is the integral sign representation. The elements $u_{2\sigma} \in \pi_{2-2\sigma}^{C_2} H\mathbb{Z}$ and $a_{\sigma} \in \pi_{-\sigma}^{C_2} H\mathbb{Z}$, under the map $H\mathbb{Z} \rightarrow F(EC_{2+}, H\mathbb{Z})$, become the elements $u_{2\sigma} \in H^0(C_2; \text{sgn}^{\otimes 2})$ and $a_{\sigma} \in H^1(C_2; \text{sgn})$. \square

Remark 3.7. In [36, Section I.9], Ullman proves a general isomorphism result for any G -spectrum X . When $G = C_2$, his isomorphism range is the region *slightly below* the line of slope 1. In our case, however, $BP_{\mathbb{R}}$ has pure and isotropic slices and nonnegative slice sections. When this happens, we can extend his isomorphism range to be *on or below* the line of slope 1. Since this line is also the vanishing line of $\text{SliceSS}(BP_{\mathbb{R}})$, the map $\text{SliceSS}(BP_{\mathbb{R}}) \rightarrow \text{HFSS}(BP_{\mathbb{R}})$ is an inclusion on the E_2 -page.

Remark 3.8. Using Lemma 3.6, one can further show that the map of $RO(C_2)$ -graded spectral sequences

$$\begin{array}{ccc} \text{SliceSS}(BP_{\mathbb{R}}) & \xrightarrow{\cong} & \pi_{\star}^{C_2} BP_{\mathbb{R}} \\ \downarrow & & \downarrow \cong \\ \text{HFSS}(BP_{\mathbb{R}}) & \xrightarrow{\cong} & \pi_{\star}^{C_2} F(EC_{2+}, BP_{\mathbb{R}}) \end{array}$$

is an inclusion on the E_2 -page and an isomorphism in this range. It turns out that in $\text{HFSS}(BP_{\mathbb{R}})$, everything outside of this isomorphism range dies by the differentials in (3.1). As a result, we obtain an equivalence $BP_{\mathbb{R}} \simeq F(EC_{2+}, BP_{\mathbb{R}})$. This is called the strong completion theorem (or the homotopy fixed point theorem) of $BP_{\mathbb{R}}$. It is proved by Hu and Kriz in [20, Theorem 4.1]. As noted in their paper, the homotopy fixed point theorem holds for $MU_{\mathbb{R}}$ as well, but not for $BP_{\mathbb{R}}\langle n \rangle$. It fails for $BP_{\mathbb{R}}\langle n \rangle$ because not everything outside the isomorphism range dies in $\text{HFSS}(BP_{\mathbb{R}}\langle n \rangle)$. In particular, $\pi_{*} BP_{\mathbb{R}}\langle n \rangle^{hC_2}$ is not bounded below.

4. The equivariant May spectral sequence

We will now construct the equivariant May spectral sequence of $BP_{\mathbb{R}}$ by filtering the equivariant dual Steenrod algebra.

4.1. The equivariant May spectral sequence with respect to A_{\star}^m

Recall from Section 2.2 that for a C_2 -spectrum X with good properties, its equivariant Adams spectral sequence with respect to $(H\underline{\mathbb{F}}_{2\star}, A_{\star}^m)$ has E_2 -page

$$E_2 = \text{Ext}_{A_{\star}^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star} X) \implies (\pi_{\star}^{C_2} X)_{\wedge}^{\wedge}.$$

Table 1
Adams and May filtrations of elements.

| | stem | Adams filtration | May filtration |
|------------------|-----------------------|------------------|----------------|
| $w_i = [\tau_i]$ | $(2^i - 1)\rho_{C_2}$ | 1 | 0 |
| a_σ | $-\sigma$ | 0 | 1 |
| u_σ | $1 - \sigma$ | 0 | 0 |

To compute the Ext-groups on the E_2 -page, we filter the Hopf algebroid $(H\underline{\mathbb{F}}_{2\star}, A_\star^m)$ by powers of the ideal (a_σ) and set the filtration of the element $\xi_i^{2^j}$ to be $2^{j+1} - 1$. We also filter the $(H\underline{\mathbb{F}}_{2\star}, A_\star^m)$ -comodule $H\underline{\mathbb{F}}_{2\star}X$ to make it compatible with the filtration on $(H\underline{\mathbb{F}}_{2\star}, A_\star^m)$.

Definition 4.1. The spectral sequence

$$E_1 = \text{Ext}_{\text{gr}\bullet A_\star^m}(\text{gr}\bullet H\underline{\mathbb{F}}_{2\star}, \text{gr}\bullet H\underline{\mathbb{F}}_{2\star}X) \implies \text{Ext}_{A_\star^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star}X)$$

is the C_2 -equivariant May spectral sequence of X with respect to $(H\underline{\mathbb{F}}_{2\star}, A_\star^m)$. It is abbreviated by $C_2\text{-MaySS}_{A_\star^m}(X)$.

We are interested in the case when $X = BP_{\mathbb{R}}$. In this case, the E_2 -page of the equivariant Adams spectral sequence simplifies:

$$E_2 = \text{Ext}_{A_\star^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star}BP_{\mathbb{R}}) = \text{Ext}_{\Lambda_\star^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star}) \implies (\pi_\star^{C_2}BP_{\mathbb{R}})_2^\wedge.$$

Here, $\Lambda_\star^m = \pi_\star^{C_2}(H\underline{\mathbb{F}}_{2\star} \wedge_{BP_{\mathbb{R}}} H\underline{\mathbb{F}}_{2\star})$, and there is a quotient map

$$A_\star^m \rightarrow \Lambda_\star^m,$$

which quotients out the ξ_i -generators. The filtration on A_\star^m induces a filtration on Λ_\star^m , which is also by powers of (a_σ) . It follows that the equivariant May spectral sequence for $BP_{\mathbb{R}}$ has E_1 -page

$$E_1 = \text{Ext}_{\text{gr}\bullet \Lambda_\star^m}(\text{gr}\bullet H\underline{\mathbb{F}}_{2\star}, \text{gr}\bullet H\underline{\mathbb{F}}_{2\star}) \implies \text{Ext}_{\Lambda_\star^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star}).$$

To compute this E_1 -page and its differentials, we use Proposition 2.15. Denote the corresponding class of τ_i in the cobar complex $C_{\Lambda_\star^m}(H\underline{\mathbb{F}}_{2\star})$ by $w_i := [\tau_i]$.

Proposition 4.2. *The E_1 -page of $C_2\text{-MaySS}_{A_\star^m}(BP_{\mathbb{R}})$ is the polynomial ring $H\underline{\mathbb{F}}_{2\star}[w_i]$. The positive part of the E_1 -page (the elements in stems $p + q\sigma$ with $p \geq 0$) is the polynomial ring*

$$\mathbb{Z}/2[u_\sigma, a_\sigma][w_i].$$

The filtration of each element is given in Table 1.

Proof. Immediate from Proposition 2.15. \square

Proposition 4.3. *In C_2 -MaySS $_{\mathcal{A}_\star^m}(BP_{\mathbb{R}})$, the classes w_i and a_σ are permanent cycles. The class $u_\sigma^{2^n}$ ($n \geq 0$) supports a differential of length $(2^{n+1} - 1)$:*

$$d_{2^{n+1}-1}(u_\sigma^{2^n}) = w_n a_\sigma^{2^{n+1}-1}.$$

Proof. Since the τ_i generators are primitive and $\eta_R(a_\sigma) = a_\sigma$, the classes w_i and a_σ are permanent cycles. To obtain the differentials on the classes $u_\sigma^{2^n}$, we use the right unit formula for u_σ . By Proposition 2.15, the right unit formula for u_σ is

$$\eta_R(u_\sigma) = u_\sigma + \tau_0 a_\sigma.$$

This translates to the d_1 -differential

$$d_1(u_\sigma) = [\tau_0]a_\sigma = w_0 a_\sigma.$$

In general, taking the right unit formula to the 2^n -th power yields the formula

$$\begin{aligned} \eta_R(u_\sigma^{2^n}) &= \eta_R(u_\sigma)^{2^n} \\ &= u_\sigma^{2^n} + \tau_0^{2^n} a_\sigma^{2^n} \\ &= u_\sigma^{2^n} + \tau_n a_\sigma^{2^{n+1}-1}, \end{aligned}$$

where for the last equality we have repeatedly used the relations $\tau_i^2 = \tau_{i+1} a_\sigma$ in Λ_\star^m . This produces the $d_{2^{n+1}-1}$ -differential on $u_\sigma^{2^n}$, as desired. \square

4.2. *The equivariant May spectral sequence with respect to \mathcal{A}_\star^{cc}*

Everything we did in the previous section can be done with respect to $(H_\star^c, \mathcal{A}_\star^{cc})$ as well. Consider the equivariant Adams spectral sequence

$$E_2 = \text{Ext}_{\mathcal{A}_\star^{cc}}(H_\star^c, H_\star^c X) \implies (\pi_\star^{C_2} X)_2^\wedge.$$

We can filter the Hopf algebraoid $(H_\star^c, \mathcal{A}_\star^{cc})$ by powers of the ideal (a_σ) to obtain a similar equivariant May spectral sequence.

Definition 4.4. The spectral sequence

$$E_1 = \text{Ext}_{\text{gr}_\bullet \mathcal{A}_\star^{cc}}(\text{gr}_\bullet H_\star^c, \text{gr}_\bullet H_\star^c X) \implies \text{Ext}_{\mathcal{A}_\star^{cc}}(H_\star^c, H_\star^c X)$$

is the C_2 -equivariant May spectral sequence for X with respect to $(H_\star^c, \mathcal{A}_\star^{cc})$. It is abbreviated by C_2 -MaySS $_{\mathcal{A}_\star^{cc}}(X)$.

When $X = BP_{\mathbb{R}}$, we can make the same simplifications as we did in Section 4.1. Let $\Lambda_{\star}^{cc} := \pi_{\star}^{C_2} F(EC_{2+}, H\mathbb{F}_2 \wedge_{BP_{\mathbb{R}}} H\mathbb{F}_2)$. The E_2 -page of the equivariant Adams spectral sequence and E_1 -page of the equivariant May spectral sequence for $BP_{\mathbb{R}}$ are equal to

$$E_2 = \text{Ext}_{\Lambda_{\star}^{cc}}(H_{\star}^c, H_{\star}^c) \implies \pi_{\star}^{C_2}(BP_{\mathbb{R}})_2^{\wedge}$$

and

$$E_1 = \text{Ext}_{\text{gr}_{\bullet}\Lambda_{\star}^{cc}}(\text{gr}_{\bullet}H_{\star}^c, \text{gr}_{\bullet}H_{\star}^c) \implies \text{Ext}_{\Lambda_{\star}^{cc}}(H_{\star}^c, H_{\star}^c)$$

respectively. As before, denote the corresponding class of τ_i in the cobar complex $C_{\Lambda_{\star}^{cc}}(H_{\star}^c)$ by $w_i = [\tau_i]$.

Proposition 4.5. *The E_1 -page of C_2 -MaySS $_{\Lambda_{\star}^{cc}}(BP_{\mathbb{R}})$ is the polynomial ring*

$$H_{\star}^c[w_i] = \mathbb{Z}/2[u_{\sigma}^{\pm}, a_{\sigma}][w_i],$$

where the filtration of each element is the same as before (see Proposition 4.2).

Proof. The claim follows directly from [20, Proposition 6.29]. \square

Proposition 4.6. *In C_2 -MaySS $_{\Lambda_{\star}^{cc}}(BP_{\mathbb{R}})$, the classes w_i and a_i are permanent cycles. The classes $u_{\sigma}^{2^n}$ and $u_{\sigma}^{-2^n}$ ($n \geq 0$) support differentials of length $(2^{n+1} - 1)$:*

$$\begin{aligned} d_{2^{n+1}-1}(u_{\sigma}^{2^n}) &= w_n a_{\sigma}^{2^{n+1}-1}, \\ d_{2^{n+1}-1}(u_{\sigma}^{-2^n}) &= w_n u_{\sigma}^{-2^{n+1}} a_{\sigma}^{2^{n+1}-1}. \end{aligned}$$

Proof. The proof is exactly the same as the proof of Proposition 4.3. The differentials follow from the right unit formulas

$$\begin{aligned} \eta_R(u_{\sigma}) &= u_{\sigma} + \tau_0 a_{\sigma}, \\ \eta_R(u_{\sigma}^{-1}) &= (u_{\sigma} + \tau_0 a_{\sigma})^{-1} = u_{\sigma}^{-1} + \tau_0 u_{\sigma}^{-2} a_{\sigma} \quad (\text{modulo higher powers of } a_{\sigma}), \end{aligned}$$

and the relation $\tau_i^2 = \tau_{i+1} a_{\sigma}$ in Λ_{\star}^{cc} . \square

4.3. C_2 -MaySS $_{\Lambda_{\star}^{cc}}(BP_{\mathbb{R}})$ and HFPSS($BP_{\mathbb{R}}$)

The equivariant May spectral sequence C_2 -MaySS $_{\Lambda_{\star}^{cc}}(BP_{\mathbb{R}})$ and the homotopy fixed point spectral sequence HFPSS($BP_{\mathbb{R}}$) have the same E_2 -page and differentials under the correspondence $w_i \leftrightarrow \bar{v}_i$. This is first observed in (7.1) and (7.2) of [20]. The only slight difference is that in C_2 -MaySS $_{\Lambda_{\star}^{cc}}(BP_{\mathbb{R}})$, instead of a single \mathbb{Z} -class, we have a w_0 -tower of $\mathbb{Z}/2$ -classes. Our goal in this section is to prove this correspondence.

Theorem 4.7. *The C_2 -equivariant May spectral sequence for $BP_{\mathbb{R}}$ with respect to $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$ is isomorphic to the associated-graded homotopy fixed point spectral sequence for $BP_{\mathbb{R}}$.*

Remark 4.8. By the associated-graded homotopy fixed point spectral sequence, we mean that whenever we see a \mathbb{Z} -class on the E_2 -page, we replace it by a tower of $\mathbb{Z}/2$ -classes.

Proof. Consider the following diagram of spectral sequences:

$$\begin{array}{ccc}
 \text{algebraic } a_{\sigma}\text{-Bockstein} & \xleftarrow[\text{collapse}]{\cong} & \text{homotopy } a_{\sigma}\text{-Bockstein} \\
 \parallel & & \parallel \\
 C_2\text{-MaySS}_{\mathcal{A}_{\star}^{cc}}(BP_{\mathbb{R}}) & \xleftarrow{\cong} & \text{HFPSS}(BP_{\mathbb{R}}).
 \end{array}$$

We will explain each arrow in the diagram one by one:

(1) $C_2\text{-MaySS}_{\mathcal{A}_{\star}^{cc}}(BP_{\mathbb{R}}) = \text{algebraic } a_{\sigma}\text{-Bockstein}$. This is by definition: the equivariant May spectral sequence for $(H_{\star}^c, \mathcal{A}_{\star}^{cc})$ is defined by filtering Λ_{\star}^{cc} by powers of (a_{σ}) , which is the algebraic a_{σ} -Bockstein.

(2) $\text{Homotopy } a_{\sigma}\text{-Bockstein} = \text{HFPSS}(BP_{\mathbb{R}})$. This is proven in [15, Lemma 4.8]. We include their proof here because it is nice and short. Start with the cofiber sequence

$$S^0 \vee S^0 \longrightarrow S^0 \xrightarrow{a_{\sigma}} S^{\sigma}.$$

Taking $F(-, BP_{\mathbb{R}})$ yields the new sequence

$$\Sigma^{-\sigma} BP_{\mathbb{R}} \xrightarrow{a_{\sigma}} BP_{\mathbb{R}} \longrightarrow BP_{\mathbb{R}} \vee BP_{\mathbb{R}}.$$

The homotopy a_{σ} -Bockstein is the spectral sequence associated to this cofiber sequence. The key observation is that in the cofiber sequence

$$S(n\sigma)_+ \longrightarrow S^0 \xrightarrow{a_{\sigma}^n} S^{n\sigma},$$

$S(n\sigma)_+$ is the $(n-1)$ -skeleton for the standard equivariant decomposition of EC_{2+} , which is used to construct the homotopy fixed point spectral sequence. Taking $F(-, BP_{\mathbb{R}})$ again, we obtain the following commutative diagram:

$$\begin{array}{ccccc}
 \Sigma^{-(n+1)\sigma} BP_{\mathbb{R}} & \xrightarrow{a_{\sigma}^{n+1}} & BP_{\mathbb{R}} & \longrightarrow & F(S((n+1)\sigma)_+, BP_{\mathbb{R}}) \\
 \downarrow a_{\sigma} & & \downarrow \text{id} & & \downarrow \\
 \Sigma^{-n\sigma} BP_{\mathbb{R}} & \xrightarrow{a_{\sigma}^n} & BP_{\mathbb{R}} & \longrightarrow & F(S(n\sigma)_+, BP_{\mathbb{R}}).
 \end{array}$$

It follows that the towers for constructing the homotopy a_{σ} -Bockstein spectral sequence and the $\text{HFPSS}(BP_{\mathbb{R}})$ are the same.

(3) Algebraic a_σ -Bockstein \cong Homotopy a_σ -Bockstein. Consider the following diagram:

$$\begin{array}{ccccc}
 \vdots & & \vdots & & \vdots \\
 \uparrow & & \uparrow & & \uparrow \\
 (H^c)^{\wedge 2} \wedge \Sigma^{-\sigma} BP_{\mathbb{R}} & \xrightarrow{a_\sigma} & (H^c)^{\wedge 2} \wedge BP_{\mathbb{R}} & \longrightarrow & (H^c)^{\wedge 2} \wedge (BP_{\mathbb{R}} \vee BP_{\mathbb{R}}) \\
 \uparrow & & \uparrow & & \uparrow \\
 H^c \wedge \Sigma^{-\sigma} BP_{\mathbb{R}} & \xrightarrow{a_\sigma} & H^c \wedge BP_{\mathbb{R}} & \longrightarrow & H^c \wedge (BP_{\mathbb{R}} \vee BP_{\mathbb{R}}) \\
 \uparrow & & \uparrow & & \uparrow \\
 \Sigma^{-\sigma} BP_{\mathbb{R}} & \xrightarrow{a_\sigma} & BP_{\mathbb{R}} & \longrightarrow & BP_{\mathbb{R}} \vee BP_{\mathbb{R}}.
 \end{array}$$

The vertical direction is the Adams resolution by $H^c = F(EC_{2+}, H\mathbb{F}_2)$, and the horizontal direction is filtering by powers of a_σ (the a_σ -Bockstein). There are two ways to compute $\pi_*^{C_2}(BP_{\mathbb{R}})_2^\wedge$ from $BP_{\mathbb{R}} \vee BP_{\mathbb{R}}$: we can either first use the horizontal filtration, then the vertical filtration, or first use the vertical filtration, and then the horizontal filtration. This produces the following commutative diagram of spectral sequences

$$\begin{array}{ccc}
 \text{Ext}_{\mathcal{A}_\star^{cc}}(H_\star^c, H_\star^c(BP_{\mathbb{R}} \vee BP_{\mathbb{R}})) & \xrightarrow{\text{algebraic } a_\sigma\text{-Bockstein}} & \text{Ext}_{\mathcal{A}_\star^{cc}}(H_\star^c, H_\star^c BP_{\mathbb{R}}) \\
 \Downarrow \text{Adams (collapse)} & & \Downarrow \text{Adams (collapse)} \\
 (\pi_\star^{C_2} BP_{\mathbb{R}} \vee BP_{\mathbb{R}})_2^\wedge & \xrightarrow{\text{homotopy } a_\sigma\text{-Bockstein}} & (\pi_\star^{C_2} F(EC_{2+}, BP_{\mathbb{R}}))_2^\wedge = (\pi_\star^{C_2} BP_{\mathbb{R}})_2^\wedge.
 \end{array}$$

The vertical spectral sequences are from the Adams (vertical) filtrations, and the horizontal spectral sequences are from the a_σ -Bockstein (horizontal). The left vertical spectral sequence collapses by degree reasons. In particular, the integer-graded part is the non-equivariant Adams spectral sequence computing $\pi_* BP$, which collapses. The right vertical spectral sequence collapses by degree reasons as well. For both Adams spectral sequences, [20, Theorem 4.11] and our computation show there are no hidden a_σ -extensions. The top spectral sequence is the a_σ -Bockstein associated with the cofiber sequences

$$(H^c)^{\wedge n} \wedge \Sigma^{-\sigma} BP_{\mathbb{R}} \xrightarrow{a_\sigma} (H^c)^{\wedge n} \wedge BP_{\mathbb{R}} \rightarrow (H^c)^{\wedge n} \wedge (BP_{\mathbb{R}} \vee BP_{\mathbb{R}})$$

for $n \geq 1$. When we are computing the Ext groups, it is exactly the same as filtering $(H_\star^c, A_\star^{cc})$ by powers of a_σ . Therefore this spectral sequence is the algebraic a_σ -Bockstein, or in other words, the C_2 -equivariant May spectral sequence with respect to \mathcal{A}_\star^{cc} . Finally, the bottom arrow is the homotopy a_σ -Bockstein, which is the homotopy fixed point spectral sequence by the previous discussion. The collapse of the two Adams spec-

tral sequences (and no hidden a_σ -extensions) implies that the algebraic a_σ -Bockstein is isomorphic to the associated-graded homotopy a_σ -Bockstein, as desired. \square

4.4. C_2 -MaySS $_{A_*^m}(BP_{\mathbb{R}})$ and SliceSS($BP_{\mathbb{R}}$)

The map $(H\underline{\mathbb{F}}_{2,*}, A_*^m) \rightarrow (H_*^c, A_*^{cc})$ induces maps of the corresponding May and Adams spectral sequences:

$$\begin{CD} \text{Ext}_{\text{gr}, \Lambda_*^m}(\text{gr}_\bullet H\underline{\mathbb{F}}_{2,*}, \text{gr}_\bullet H\underline{\mathbb{F}}_{2,*}) @>\text{May}>> \text{Ext}_{\Lambda_*^m}(H\underline{\mathbb{F}}_{2,*}, H\underline{\mathbb{F}}_{2,*}) @>\text{Adams}>> \pi_*^{C_2}(BP_{\mathbb{R}})_2^\wedge \\ @VVV @VVV @VV\text{id}V \\ \text{Ext}_{\text{gr}, \Lambda_*^{cc}}(\text{gr}_\bullet H_*^c, \text{gr}_\bullet H_*^c) @>\text{May}>> \text{Ext}_{\Lambda_*^{cc}}(H_*^c, H_*^c) @>\text{Adams}>> \pi_*^{C_2}(BP_{\mathbb{R}})_2^\wedge \end{CD}$$

For the purpose of finding Hurewicz images, we restrict our attention to the maps between integer-graded spectral sequences. They are induced from the map of integer-graded Hopf algebroids $(H\underline{\mathbb{F}}_{2,*}, A_*^m) \rightarrow (H_*^c, A_*^{cc})$:

$$\begin{CD} \text{Ext}_{\text{gr}, \Lambda_*^m}(\text{gr}_\bullet H\underline{\mathbb{F}}_{2,*}, \text{gr}_\bullet H\underline{\mathbb{F}}_{2,*}) @>\text{May}>> \text{Ext}_{\Lambda_*^m}(H\underline{\mathbb{F}}_{2,*}, H\underline{\mathbb{F}}_{2,*}) @>\text{Adams}>> \pi_*^{C_2}(BP_{\mathbb{R}})_2^\wedge \\ @VVV @VVV @VV\text{id}V \\ \text{Ext}_{\text{gr}, \Lambda_*^{cc}}(\text{gr}_\bullet H_*^c, \text{gr}_\bullet H_*^c) @>\text{May}>> \text{Ext}_{\Lambda_*^{cc}}(H_*^c, H_*^c) @>\text{Adams}>> \pi_*^{C_2}(BP_{\mathbb{R}})_2^\wedge \end{CD}$$

The E_2 -page of C_2 -MaySS $_{A_*^m}(BP_{\mathbb{R}})$ is the subring of the polynomial ring $H\underline{\mathbb{F}}_{2,*}[w_i]$ that contains only integer-graded elements. For degree reasons, monomials of the form $u_\sigma^{-k} a_\sigma^{-l} w_{i_1}^{e_1} w_{i_2}^{e_2} \cdots w_{i_j}^{e_j}$ with $k > 0, l \geq 0$ do not have integer grading. It follows that the integer graded elements are all contained in the subring

$$\mathbb{Z}/2[u_\sigma, a_\sigma][w_i] \subset H\underline{\mathbb{F}}_{2,*}[w_i].$$

Theorem 4.9. *The integer-graded C_2 -equivariant May spectral sequence of $BP_{\mathbb{R}}$ with respect to $(H\underline{\mathbb{F}}_{2,*}, A_*^m)$ is isomorphic to the associated-graded slice spectral sequence of $BP_{\mathbb{R}}$.*

Proof. Consider the following diagram:

$$\begin{CD} C_2\text{-MaySS}_{A_*^m}(BP_{\mathbb{R}}) @<\cong<< \text{SliceSS}(BP_{\mathbb{R}}) \\ @VVV @VVV \\ C_2\text{-MaySS}_{A_*^{cc}}(BP_{\mathbb{R}}) @<\cong<< \text{HFPSS}(BP_{\mathbb{R}}). \end{CD}$$

The above discussion, together with Proposition 4.3 and Proposition 4.6, show that the left vertical map is an inclusion on the E_2 -page and an isomorphism on or below the line of slope 1. Proposition 3.5 shows that the right vertical map is also an inclusion on

the E_2 -page and an isomorphism on or below the line of slope 1. Given the isomorphism already established in Theorem 4.7, it follows that C_2 -MaySS $_{A_*^m}(BP_{\mathbb{R}})$ is isomorphic to the associated-graded SliceSS($BP_{\mathbb{R}}$), as desired. \square

Remark 4.10. The isomorphism in Theorem 4.7 is of $RO(C_2)$ -graded spectral sequences. We have only proven the isomorphism in Theorem 4.9 as integer-graded spectral sequences. This is enough for the purpose of proving Hurewicz images.

5. Map of May spectral sequences

5.1. Map of dual Steenrod algebras

The maps $\mathcal{A}_* \rightarrow \mathcal{A}_*^m \rightarrow \Lambda_*^m$ induce maps of Adams E_2 -pages:

$$E_2(\text{ASS}(\mathbb{S})) \rightarrow E_2(C_2\text{-ASS}(\mathbb{S})) \rightarrow E_2(C_2\text{-ASS}(BP_{\mathbb{R}})).$$

Filtering \mathcal{A}_* , \mathcal{A}_*^m , and Λ_*^m compatibly with respect to the map above produces maps of May spectral sequences

$$\text{MMaySS}(\mathbb{S}) \rightarrow C_2\text{-MaySS}(\mathbb{S}) \rightarrow C_2\text{-MaySS}(BP_{\mathbb{R}}).$$

Here, $\text{MMaySS}(\mathbb{S})$ is the modified May spectral sequence, which will be defined in section 5.2. These maps of May spectral sequences will later help us prove our detection theorem for $BP_{\mathbb{R}}^{C_2}$.

Recall that

$$\begin{aligned} \mathcal{A}_* &= \mathbb{F}_2[\zeta_1, \zeta_2, \dots], \\ \mathcal{A}_*^m &= H\underline{\mathbb{F}}_{2,*}[\xi_i, \tau_i]/(\tau_i^2 = \tau_{i+1}a_\sigma + \xi_{i+1}\eta_R(u_\sigma)), \\ \Lambda_*^m &= H\underline{\mathbb{F}}_{2,*}[\tau_i]/(\tau_i^2 = \tau_{i+1}a_\sigma). \end{aligned}$$

The following theorem will be used later for

- (1) Constructing and computing the map $\text{MMaySS}(\mathbb{S}) \rightarrow C_2\text{-MaySS}(BP_{\mathbb{R}})$ of May spectral sequences.
- (2) Computing the images of elements in the classical Adams spectral sequence $\text{ASS}(\mathbb{S})$ under the map

$$\text{ASS}(\mathbb{S}) \rightarrow C_2\text{-ASS}(BP_{\mathbb{R}}).$$

Theorem 5.1. *The composite map $\mathcal{A}_* \rightarrow \mathcal{A}_*^m \rightarrow \Lambda_*^m$ sends the element*

$$\zeta_i^{2^j} \mapsto \tau_{i+j-1}u_\sigma^{2^{i+j-1}-2^j}a_\sigma^{2^j-1} \text{ (modulo higher powers of } a_\sigma\text{)}.$$

Proof. By an abuse of notation, we will denote ζ_i to be the image of $\zeta_i \in \mathcal{A}_*$ in \mathcal{A}_*^m and Λ_*^m . The following relations hold in \mathcal{A}_*^m (cf. [20, Theorem 6.18, Theorem 6.41] and Theorem 2.10):

$$a_\sigma^{2^i} \zeta_i = \zeta_{i-1}^2 \eta_R(u_\sigma) + \zeta_i a_\sigma + \zeta_{i-1} u_\sigma^{2^{i-1}}, \tag{5.1}$$

$$a_\sigma \tau_0 = u_\sigma + \eta_R(u_\sigma). \tag{5.2}$$

To prove our claim, we will show using induction on i that

$$\zeta_i \mapsto \tau_{i-1} \eta_R(u_\sigma)^{2^{i-1}-1} \pmod{\xi_1, \xi_2, \dots}.$$

For the base case when $i = 1$, equations (5.1) and (5.2) imply

$$\begin{aligned} a_\sigma^2 \zeta_1 &= \eta_R(u_\sigma) + \zeta_1 a_\sigma + u_\sigma \\ \implies \zeta_1 a_\sigma &= u_\sigma + \eta_R(u_\sigma) = a_\sigma \tau_0 \pmod{\xi_1, \xi_2, \dots} \\ \implies \zeta_1 &= \tau_0 \pmod{\xi_1, \xi_2, \dots}. \end{aligned}$$

Therefore the base case holds. Now, suppose we have shown that

$$\zeta_{i-1} = \tau_{i-2} \eta_R(u_\sigma)^{2^{i-2}-1} \pmod{\xi_1, \xi_2, \dots}.$$

To prove the relation for ζ_i , we use relation (5.1) again:

$$\begin{aligned} \zeta_i a_\sigma &= \zeta_{i-1}^2 \eta_R(u_\sigma) \pmod{\xi_1, \xi_2, \dots} \\ &= (\tau_{i-2} \eta_R(u_\sigma)^{2^{i-2}-1})^2 \eta_R(u_\sigma) \pmod{\xi_1, \xi_2, \dots} \text{ (induction hypothesis)} \\ &= \tau_{i-2}^2 \eta_R(u_\sigma)^{2^{i-1}-1} \pmod{\xi_1, \xi_2, \dots} \\ &= (\tau_{i-1} a_\sigma) \eta_R(u_\sigma)^{2^{i-1}-1} \pmod{\xi_1, \xi_2, \dots} \\ \implies \zeta_i &= \tau_{i-1} \eta_R(u_\sigma)^{2^{i-1}-1} \pmod{\xi_1, \xi_2, \dots}, \end{aligned}$$

as desired.

To finish the proof of the theorem, we need to simplify the expression

$$\zeta_i^{2^j} = \left(\tau_{i-1} \eta_R(u_\sigma)^{2^{i-1}-1} \right)^{2^j}$$

modulo higher powers of a_σ . Since $u_\sigma + \eta_R(u_\sigma) = \tau_0 a_\sigma$,

$$u_\sigma \equiv \eta_R(u_\sigma) \pmod{\text{modulo higher powers of } a_\sigma}.$$

After applying the relation $\tau_n^2 = \tau_{n+1} a_\sigma$ j -times, we obtain the equality

$$\zeta_i^{2^j} \equiv \tau_{i+j-1} u_\sigma^{2^{i+j-1}-2^j} a_\sigma^{2^j-1} \pmod{\text{modulo higher powers of } a_\sigma},$$

as desired. \square

5.2. Change of filtration

The maps $\mathcal{A}_* \rightarrow A_*^m \rightarrow \Lambda_*^m$ induce maps of Ext groups

$$\begin{array}{ccc} \text{Ext}_{\mathcal{A}_*}(\mathbb{F}_2, \mathbb{F}_2) & \longrightarrow & \text{Ext}_{A_*^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star}) \\ & \searrow & \downarrow \\ & & \text{Ext}_{\Lambda_*^m}(H\underline{\mathbb{F}}_{2\star}, H\underline{\mathbb{F}}_{2\star}). \end{array}$$

To analyze these maps, we will construct maps of May spectral sequences:

$$\text{MMaySS}(\mathbb{S}) \rightarrow C_2\text{-MaySS}(\mathbb{S}) \rightarrow C_2\text{-MaySS}(BP_{\mathbb{R}}).$$

We do so by filtering the classical dual Steenrod algebra \mathcal{A}_* and the equivariant dual Steenrod algebras A_*^m and Λ_*^m compatibly with respect to the maps $\mathcal{A}_* \rightarrow A_*^m \rightarrow \Lambda_*^m$.

Recall that in constructing the classical May spectral sequence, the dual Steenrod algebra $\mathcal{A}_* = \mathbb{F}_2[\zeta_1, \zeta_2, \dots]$ is filtered by powers of its unit coideal, hence producing an increasing filtration. More specifically, we can define a grading on \mathcal{A}_* by setting the degree of $h_{i,j} := \zeta_i^{2^j}$ to be

$$|h_{i,j}| = 2i - 1$$

and extend additively to all unique representatives. The increasing filtration associated to this grading is

$$F_0\mathcal{A}_* \subset F_1\mathcal{A}_* \subset \dots \subset F_p\mathcal{A}_* \subset F_{p+1}\mathcal{A}_* \subset \dots \subset \mathcal{A}_*,$$

where at stage p , $F_p\mathcal{A}_*$ consists of all elements of total degrees $\leq p$.

However, in constructing the equivariant May spectral sequence $C_2\text{-MaySS}(\mathbb{S})$ and $C_2\text{-MaySS}(BP_{\mathbb{R}})$, we filtered \mathcal{A}_*^m and Λ_*^m by powers of (a_σ) and produced decreasing filtrations instead. To rectify this mismatch of filtrations, we need to change the filtration of the classical dual Steenrod algebra \mathcal{A}_* to make it compatible with the decreasing filtrations on \mathcal{A}_*^m and Λ_*^m . In particular, it must be a decreasing filtration. To do this, notice that by Theorem 5.1, the element $h_{i,j}$ is sent to

$$h_{i,j} \mapsto \tau_{i+j-1} u_\sigma^{2^{i+j-1}-2^j} a_\sigma^{2^j-1} \text{ (Modulo higher powers of } a_\sigma). \tag{5.3}$$

We can define a new grading on \mathcal{A}_* on by setting the degree of the generators $h_{i,j}$ to be

$$|h_{i,j}| = 2^j - 1$$

and extend linearly to all unique representatives. The decreasing filtration associated to this is

$$\mathcal{A}_* = F_0\mathcal{A}_* \supset F_1\mathcal{A}_* \supset F_2\mathcal{A}_* \supset \dots,$$

where $F_p\mathcal{A}_*$ contains elements whose total degrees are $\geq p$. From (5.3), it is immediate that this filtration is compatible with the decreasing filtration of A_\star^m and Λ_\star^m with respect to the maps $\mathcal{A}_* \rightarrow A_\star^m \rightarrow \Lambda_\star^m$. Therefore, we obtain maps of May spectral sequences

$$\begin{array}{ccc} \text{MMaySS}(\mathbb{S}) & \longrightarrow & C_2\text{-MaySS}(\mathbb{S}) \\ & \searrow & \downarrow \\ & & C_2\text{-MaySS}(BP_{\mathbb{R}}). \end{array}$$

To compute the E_1 -page of our modified May spectral sequence $\text{MMaySS}(\mathbb{S})$, consider the coproduct formula for $h_{i,j}$:

$$\begin{aligned} \Psi(h_{i,j}) &= 1 \otimes h_{i,j} + h_{i,j} \otimes 1 + \sum_{k=1}^{i-1} h_{i-k,j+k} \otimes h_{k,j} \\ &= \underbrace{1 \otimes h_{i,j}}_{\text{deg}=2^j-1} + \underbrace{h_{i,j} \otimes 1}_{\text{deg}=2^j-1} + \underbrace{h_{i-1,j+1} \otimes h_{1,j}}_{\text{deg}=(2^{j+1}-1)+(2^j-1)} + \underbrace{h_{i-2,j+2} \otimes h_{2,j}}_{\text{deg}=(2^{j+2}-1)+(2^j-1)} + \\ &\quad \dots + \underbrace{h_{1,j+i-1} \otimes h_{i-1,j}}_{\text{deg}=(2^{j+i-1}-1)+(2^j-1)}. \end{aligned}$$

With the old filtration, $|h_{i,j}| = 2i - 1$, and everything in the summation sign on the right is of degree exactly $2i - 2$. After changing to our new filtration, everything in this sum is of degree at least

$$\begin{aligned} |h_{i-1,j+1} \otimes h_{1,j}| &= (2^{j+1} - 1) + (2^j - 1) \\ &> 2^j - 1 \\ &= |h_{i,j}|. \end{aligned}$$

Therefore after projecting to the associated-graded $\mathcal{A}_* \rightarrow \text{gr}_\bullet \mathcal{A}_*$, the elements $h_{i,j}$ are primitive. It follows that the E_1 -page of our modified May spectral sequence is still the polynomial algebra generated by the $h_{i,j}$:

$$E_1 = \text{Ext}_{\text{gr}_\bullet \mathcal{A}_*}(\mathbb{F}_2, \mathbb{F}_2) = \mathbb{F}_2[\{h_{i,j}\}_{i \geq 1, j \geq 0}] \implies \text{Ext}_{\mathcal{A}_*}(\mathbb{F}_2, \mathbb{F}_2).$$

The differentials obtained from the coproduct formula for $h_{i,j}$ is now of length $2^{j+1} - 1$:

$$d_{2^{j+1}-1}(h_{i,j}) = h_{i-1,j+1} \otimes h_{1,j}$$

(before changing the filtration, it was $d_1(h_{i,j}) = \sum_{k=1}^{i-1} h_{i-k,j+k} \otimes h_{k,j}$). Intuitively, with our new filtration, the differentials are being “stretched out”.

6. Detection theorem

6.1. Detection theorems for $BP_{\mathbb{R}}$ and $MU_{\mathbb{R}}$

We will now prove our detection theorems for the Hopf, Kervaire, and $\bar{\kappa}$ -family by analyzing the map of spectral sequences

$$MMaySS(\mathbb{S}) \rightarrow C_2\text{-MaySS}_{\mathcal{A}_{\star}^m}(BP_{\mathbb{R}}) \cong \text{SliceSS}(BP_{\mathbb{R}}).$$

Using Theorem 5.1, the following proposition is immediate.

Proposition 6.1. *On the E_2 -page of the map $MMaySS(\mathbb{S}) \rightarrow C_2\text{-MaySS}(BP_{\mathbb{R}}) \cong \text{SliceSS}(BP_{\mathbb{R}})$,*

$$\begin{aligned} h_{1n} &\mapsto \bar{v}_n a_{\sigma}^{2^n-1}, \\ h_{1n}^2 &\mapsto \bar{v}_n^2 a_{\sigma}^{2(2^n-1)}, \\ h_{2n}^4 &\mapsto \bar{v}_{n+1}^4 u_{\sigma}^{2^{n+2}} a_{\sigma}^{4(2^n-1)}. \end{aligned}$$

Proposition 6.2. *In the slice spectral sequence of $BP_{\mathbb{R}}$, the classes $\bar{v}_n a_{\sigma}^{2^n-1}$, $\bar{v}_n^2 a_{\sigma}^{2(2^n-1)}$, and $\bar{v}_{n+1}^4 u_{\sigma}^{2^{n+2}} a_{\sigma}^{4(2^n-1)}$ survive to the E_{∞} -page.*

Proof. As discussed in Proposition 3.4, all of the differentials in the slice spectral sequence for $BP_{\mathbb{R}}$ are completely classified by the slice differential theorem ([18, Theorem 9.9]). They are

$$d_{2^{k+1}-1}(u_{\sigma}^{2^k}) = a_{\sigma}^{2^{k+1}-1} \bar{v}_k, \quad k \geq 1.$$

The longest possible differentials that could possibly kill the classes mentioned are differentials of length $2^{n+1} - 1$. The survival of these classes is a straightforward computation (see Fig. 6). \square

Theorem 6.3 (*Detection of Hopf elements and Kervaire classes*). *If the element h_n or h_n^2 in $\text{Ext}_{\mathcal{A}_{\star}}(\mathbb{F}_2, \mathbb{F}_2)$ survives to the E_{∞} -page of the Adams spectral sequence, then its image under the Hurewicz map $\pi_{\star}\mathbb{S} \rightarrow \pi_{\star}BP_{\mathbb{R}}^{C_2}$ is nonzero.*

Proof. In the modified May spectral sequence, the classes h_{1n} and h_{1n}^2 are permanent cycles. Furthermore, since h_{1n} is of Adams filtration 1 and h_{1n}^2 is of Adams filtration 2, if they are targets of differentials in the modified May spectral sequence, then the source of these differentials must be of Adams filtrations 0 and 1, respectively. This is clearly impossible. Therefore they are not targets of differentials and hence survive to the E_{∞} -page of the modified May spectral sequence.




Fig. 6. A part of the slice spectral sequence for $BP_{\mathbb{R}}$. The classes $\bar{v}_n a_{\sigma}^{2^n-1}$, $\bar{v}_n^2 a_{\sigma}^{2(2^n-1)}$, and $\bar{v}_{n+1}^4 u_{\sigma}^{2^{n+2}} a_{\sigma}^{4(2^n-1)}$ survive to the E_{2n+1} -page, hence to the E_{∞} -page.

On the E_{∞} -page of the modified May spectral sequence, these classes represent the unique elements $h_n \in \text{Ext}_{\mathcal{A}_*}^{1,2^n}(\mathbb{F}_2, \mathbb{F}_2)$ and $h_n^2 \in \text{Ext}_{\mathcal{A}_*}^{2,2^{n+1}}(\mathbb{F}_2, \mathbb{F}_2)$ in their respective bidegrees. The theorem now follows from Proposition 6.1 and Proposition 6.2. \square

Theorem 6.4 (Detection of $\bar{\kappa}$ -family). *If the element $g_n \in \text{Ext}_{\mathcal{A}_*}^{4,2^{n+2}+2^{n+3}}(\mathbb{F}_2, \mathbb{F}_2)$ survives to the E_{∞} -page of the Adams spectral sequence, then its image under the Hurewicz map $\pi_* \mathbb{S} \rightarrow \pi_* BP_{\mathbb{R}}^{C_2}$ is nonzero.*

Remark 6.5. As mentioned previously, the element g_1 survives to the element $\bar{\kappa} \in \pi_{20} \mathbb{S}$. The element g_2 also survives to an element in $\pi_{44} \mathbb{S}$. They are both detected in $\pi_* BP_{\mathbb{R}}^{C_2}$. The fate of the g_n elements for $n \geq 3$ is unknown.

The proof of Theorem 6.4 requires the following facts:

Lemma 6.6. *The element g_n is the only nonzero element in $\text{Ext}_{\mathcal{A}_*}^{4,2^{n+2}+2^{n+3}}(\mathbb{F}_2, \mathbb{F}_2)$.*

Proof. We appeal to the classification theorem of Lin ([29, Theorem 1.3]). For indecomposable elements in $\text{Ext}_{\mathcal{A}_*}^{4,t}(\mathbb{F}_2, \mathbb{F}_2)$, binary expansion of t shows that g_n is the only indecomposable element in $\text{Ext}_{\mathcal{A}_*}^{4,2^{n+2}+2^{n+3}}(\mathbb{F}_2, \mathbb{F}_2)$. The only possible decomposable ele-

ments in the same bidegree as g_n are $h_n c_n$ and $h_{n+1}^2 h_{n+2}^2$. However, they are both zero due to relations in $\text{Ext}_{\mathcal{A}_*}^{s,t}(\mathbb{F}_2, \mathbb{F}_2)$. \square

Lemma 6.7. *Let $1 \leq m \leq n$, if*

$$\text{deg}(h_{i_1, j_1} h_{i_2, j_2} \cdots h_{i_m, j_m}) < \text{deg}(h_{i'_1, j'_1} h_{i'_2, j'_2} \cdots h_{i'_n, j'_n}),$$

then for any $k \geq 1$,

$$\text{deg}(h_{i_1, j_1+k} h_{i_2, j_2+k} \cdots h_{i_m, j_m+k}) < \text{deg}(h_{i'_1, j'_1+k} h_{i'_2, j'_2+k} \cdots h_{i'_n, j'_n+k}).$$

Proof. The given condition translates to the inequality

$$\sum (2^{j_m} - 1) < \sum (2^{j'_n} - 1),$$

which, after rearranging, is

$$\sum 2^{j_m} < \left(\sum 2^{j'_n} \right) - (n - m).$$

Multiplying both sides of the inequality by 2^k produces the new inequality

$$\begin{aligned} \sum 2^{j_m+k} &< \left(\sum 2^{j'_n+k} \right) - 2^k (n - m) \\ &< \left(\sum 2^{j'_n+k} \right) - (n - m). \end{aligned}$$

Rearranging this inequality gives

$$\sum (2^{j_m+k} - 1) < \sum (2^{j'_n+k} - 1),$$

as desired. \square

Lemma 6.8. *The element h_{21}^4 is a permanent cycle in the modified May spectral sequence of the sphere.*

Proof. It is clear from the modified May filtration that all the differentials are of odd length. First, we have

$$d_3(h_{21}) = h_{11} h_{12},$$

so $d_3(h_{21}^2) = 0$. In fact, $d_5(h_{21}^2) = 0$ as well. To show this, notice that h_{21}^2 is in tridegree $(s, t, m) = (2, 12, 2)$, where (s, t) is the degree associated to $\text{Ext}_{\mathcal{A}_*}^{s,t}(\mathbb{F}_2, \mathbb{F}_2)$ and m is the modified May filtration. If h_{21}^2 supports a d_5 , then the target must be of tridegree $(3, 12, 7)$. We will characterize all $h_{i_1 j_1} h_{i_2 j_2} h_{i_3 j_3}$ of this tridegree. The equations that need to be satisfied are

$$(2^{i_1} - 1)2^{j_1} + (2^{i_2} - 1)2^{j_2} + (2^{i_3} - 1)2^{j_3} = 12,$$

$$(2^{j_1} - 1) + (2^{j_2} - 1) + (2^{j_3} - 1) = 7.$$

Since $7 = 7 + 0 + 0 = 3 + 3 + 1$, it's not hard to check that the only possibility for $h_{i_1j_1}h_{i_2j_2}h_{i_3j_3}$ is $h_{10}h_{20}h_{13}$. However, this element cannot be the target of a differential because it supports a nontrivial d_1 -differential

$$d_1(h_{10}h_{20}h_{13}) = h_{10}^2h_{11}h_{13}.$$

Therefore $d_5(h_{21}^2) = 0$.

By computations in the cobar complex $C(\mathcal{A}_*)$ (see [34, Lemma 3.2.10(b)]), we deduce that

$$d_7(h_{21}^2) = h_{12}^3 + h_{11}^2h_{13}.$$

(Note that the computation in the cobar complex also shows that h_{21}^2 is a d_5 -cycle.) By the Leibneiz rule, this differential implies that $d_7(h_{21}^4) = 0$.

The element h_{21}^4 is in tridegree $(s, t, m) = (4, 24, 4)$. The target of a differential d_r with source h_{21}^4 must be of tridegree $(5, 24, 4 + r)$. In particular, it must be a linear combination of elements of the form $h_{i_1j_1} \cdots h_{i_5j_5}$ satisfying the equations

$$\sum_{k=1}^5 (2^{i_k} - 1)2^{j_k} = 24, \tag{6.1}$$

$$\sum_{k=1}^5 (2^{j_k} - 1) = 4 + r. \tag{6.2}$$

Since $d_7(h_{21}^4) = 0$, r must be at least 9. Moreover, subtracting Equation (6.2) from Equation (6.1) yields the equation

$$\sum_{k=1}^5 (2^{i_k+j_k} - 2^{j_k+1} + 1) = 20 - r.$$

The left hand side is at least 5 because $i_k \geq 1$. It follows that $r \leq 15$. We will now rule out each possibility in the range $9 \leq r \leq 15$ case-by-case:

Case 1: $r = 9$. Equation (6.2) becomes

$$\sum_{k=1}^5 (2^{j_k} - 1) = 13.$$

The possibilities are

$$\begin{aligned} 13 &= 7 + 3 + 3 + 0 + 0 \\ &= 7 + 3 + 1 + 1 + 1 \\ &= 3 + 3 + 3 + 3 + 1. \end{aligned}$$

The second and third possibilities give nothing. The first possibility gives $h_{10}h_{30}h_{12}^2h_{13}$, which supports a nontrivial d_1 differential

$$d_1(h_{10}h_{30}h_{12}^2h_{13}) = h_{10}^2h_{21}h_{12}^2h_{13}.$$

Therefore $d_9(h_{21}^4) = 0$.

Case 2: $r = 11$. Equation (6.1) becomes

$$\sum_{k=1}^5 (2^{j_k} - 1) = 15.$$

The possibilities are

$$\begin{aligned} 15 &= 15 + 0 + 0 + 0 + 0 \\ &= 7 + 7 + 1 + 0 + 0 \\ &= 7 + 3 + 3 + 1 + 1 \\ &= 3 + 3 + 3 + 3 + 3. \end{aligned}$$

The first possibility gives $h_{10}^2h_{20}^2h_{14}$. The second possibility gives $h_{10}^2h_{21}h_{13}^2$ and $h_{11}h_{20}^2h_{13}^2$. The third possibility gives $h_{11}h_{21}h_{12}^2h_{13}$. The fourth possibility gives nothing. Now, we will rule them out one by one.

- For $h_{10}^2h_{20}^2h_{14}$, we can first argue using the cobar complex that

$$d_3(h_{20}^2) = h_{10}^2h_{12} + h_{11}^3.$$

So there is a nontrivial d_3 -differential

$$\begin{aligned} d_3(h_{10}^2h_{20}^2h_{14}) &= h_{10}^2h_{14}(h_{10}^2h_{12} + h_{11}^3) \\ &= h_{10}^4h_{12}h_{14} + h_{10}^2h_{11}^3h_{14} \\ &= h_{10}^4h_{12}h_{14} \end{aligned}$$

(the element $h_{10}^2h_{11}^3h_{14} = 0$ on the E_3 -page because $d_1(h_{20}) = h_{10}h_{11}$).

- The element $h_{10}^2 h_{21} h_{13}^2$ is the target of the d_1 -differential

$$d_1(h_{10} h_{30} h_{13}^2) = h_{10}^2 h_{21} h_{13}^2,$$

and hence 0 on the E_3 -page.

- The element $h_{11} h_{20}^2 h_{13}^2$ supports a nontrivial d_3 -differential

$$d_3(h_{11} h_{20}^2 h_{13}^2) = h_{11} h_{13}^2 (h_{10}^2 h_{12} + h_{11}^3) = h_{11}^4 h_{13}^2$$

(the first term in the sum is 0 because $h_{10} h_{11} = 0$ on the E_3 -page).

- The element $h_{11} h_{21} h_{12}^2 h_{13}$ supports a nontrivial d_3 differential

$$d_3(h_{11} h_{21} h_{12}^2 h_{13}) = h_{11} h_{12}^2 h_{13} d_3(h_{21}) = h_{11}^2 h_{12}^3 h_{13},$$

hence does not survive past the E_3 -page.

Therefore $d_{11}(h_{21}^4) = 0$.

Case 3: $r = 13$. Equation (6.2) becomes

$$\sum_{k=1}^5 (2^{j_k} - 1) = 17.$$

The possibilities are

$$\begin{aligned} 17 &= 15 + 1 + 1 + 0 + 0 \\ &= 7 + 7 + 3 + 0 + 0 \\ &= 7 + 7 + 1 + 1 + 1 \\ &= 7 + 3 + 3 + 3 + 1. \end{aligned}$$

The first possibility gives $h_{10} h_{20} h_{11}^2 h_{14}$. The second possibility gives $h_{10} h_{20} h_{12} h_{13}^2$. The third and fourth possibilities give nothing. Both elements support nontrivial d_1 differentials:

$$\begin{aligned} d_1(h_{10} h_{20} h_{11}^2 h_{14}) &= h_{10}^2 h_{11}^3 h_{14}, \\ d_1(h_{10} h_{20} h_{12} h_{13}^2) &= h_{10}^2 h_{11} h_{12} h_{13}^2. \end{aligned}$$

Therefore $d_{13}(h_{21}^4) = 0$.

Case 4: $r = 15$. Equation (6.2) becomes

$$\sum_{k=1}^5 (2^{j_k} - 1) = 19.$$

The possibilities are

$$\begin{aligned}
 19 &= 15 + 3 + 1 + 0 + 0 \\
 &= 15 + 1 + 1 + 1 + 1 \\
 &= 7 + 7 + 3 + 1 + 1 \\
 &= 7 + 3 + 3 + 3 + 3.
 \end{aligned}$$

The first possibility gives $h_{10}^2 h_{11} h_{12} h_{14}$. The second possibility gives $h_{11}^4 h_{14}$. The third possibility gives $h_{11}^2 h_{12} h_{13}^2$. The fourth possibility gives $h_{12}^4 h_{13}$. They do not survive because of the following differentials:

$$\begin{aligned}
 d_1(h_{10} h_{20} h_{12} h_{14}) &= h_{10}^2 h_{11} h_{12} h_{14}, \\
 d_3(h_{11} h_{20}^2 h_{14}) &= h_{11} h_{14} d_3(h_{20}^2) \\
 &= h_{11} h_{14} (h_{10}^2 h_{12} + h_{11}^3) \\
 &= h_{11}^4 h_{14}, \\
 d_3(h_{11} h_{21} h_{13}^2) &= h_{11}^2 h_{12} h_{13}^2, \\
 d_7(h_{12}^3 h_{22}) &= h_{12}^4 h_{13}.
 \end{aligned}$$

Therefore $d_{15}(h_{21}^4) = 0$. This concludes the proof of the Lemma. \square

Proposition 6.9. *For $n \geq 1$, the elements h_{2n}^4 survive to the E_∞ -page of the modified May spectral sequence of the sphere.*

Proof. We will first show that for all $n \geq 1$, h_{2n}^4 is not the target of a differential. By Proposition 6.1 and Proposition 6.2, the image of h_{2n}^4 on the E_2 -page under the map $\text{MMaySS}(\mathbb{S}) \rightarrow \text{SliceSS}(BP_{\mathbb{R}})$ is $\bar{v}_{n+1}^4 u_\sigma^{2^{n+2}} a_\sigma^{4(2^n-1)}$, which survives to the E_∞ -page of the slice spectral sequence. However, if h_{2n}^4 is the target of a differential d_r , then its image must also be the target of a differential $d_{r'}$, with $r' \leq r$. This is a contradiction. Therefore h_{2n}^4 is not the target of a differential.

We now show that h_{2n}^4 is also a permanent cycle. By Lemma 6.8, h_{21}^4 is a permanent cycle. This means that we can find an element x in the cobar complex $C(\mathcal{A}_*)$ of the form

$$x = \underbrace{\xi_2^{2^1} | \xi_2^{2^1} | \xi_2^{2^1} | \xi_2^{2^1}}_{\text{deg}=4 \cdot (2^1-1)=4} + \underbrace{S}_{\text{deg}>4}$$

such that $d(x) = 0$. In the expression for x , S is a sum containing elements of the form

$$s = h_{i_1, j_1} \cdots | \cdots | \cdots | \cdots h_{i_k, j_k},$$

with $k \geq 4$ and

$$\begin{aligned} \deg(\xi_2^{2^1} | \xi_2^{2^1} | \xi_2^{2^1} | \xi_2^{2^1}) &= (2^1 - 1) + (2^1 - 1) + (2^1 - 1) + (2^1 - 1) \\ &< \sum_k (2^{j_k} - 1) \\ &= \deg(s). \end{aligned} \tag{6.3}$$

To show that h_{2n}^4 is a permanent cycle, we apply the Sq^0 -operation introduced by Nakamura [33] $(n - 1)$ -times to the element x , obtaining a new element

$$(Sq^0)^{n-1}(x) = \underbrace{\xi_2^{2^n} | \xi_2^{2^n} | \xi_2^{2^n} | \xi_2^{2^n}}_{\deg=4 \cdot (2^n - 1)} + \underbrace{(Sq^0)^{n-1}(S)}_{\deg>4 \cdot (2^n - 1)}$$

in the cobar complex. Everything in $(Sq^0)^{n-1}(S)$ is of the form

$$(Sq^0)^{n-1}(s) = h_{i_1, j_1+n-1} \cdots | \cdots | \cdots | \cdots | \cdots h_{i_k, j_k+n-1}.$$

Lemma 6.7, applied to inequality (6.3), shows that

$$\deg((Sq^0)^{n-1}(\xi_2^{2^1} | \xi_2^{2^1} | \xi_2^{2^1} | \xi_2^{2^1})) = 4 \cdot (2^n - 1) < \deg((Sq^0)^{n-1}(s)).$$

By [33, Lemma 3.1], the Sq^0 -operation preserves the coboundary operator of the cobar complex $C(\mathcal{A}_*)$. Therefore

$$d((Sq^0)^{n-1}(x)) = (Sq^0)^{n-1}d(x) = (Sq^0)^{n-1}(0) = 0.$$

It follows that h_{2n}^4 is a permanent cycle, as desired. \square

Proof of Theorem 6.4. By Proposition 6.9, the elements $\{h_{2n}^4 \mid n \geq 1\}$ survive to the E_∞ -page of the modified May spectral sequence, hence they detect some nonzero elements in $\text{Ext}_{\mathcal{A}_*}^{4, 2^{n+2}+2^{n+3}}(\mathbb{F}_2, \mathbb{F}_2)$. By Lemma 6.6, these elements must be $\{g_n \mid n \geq 1\}$. The theorem now follows from Proposition 6.1 and Propostion 6.2. \square

Remark 6.10. By [20, Theorem 4.1], the map $\pi_*BP_{\mathbb{R}}^{C_2} \rightarrow \pi_*BP_{\mathbb{R}}^{hC_2}$ is an isomorphism. Therefore Theorem 6.3 and Theorem 6.4 hold for the homotopy fixed point of $BP_{\mathbb{R}}$ as well.

Theorem 6.3 and Theorem 6.4 combine to produce our detection theorems for $MU_{\mathbb{R}}$ and $BP_{\mathbb{R}}$:

Theorem 6.11 (*Detection theorems for $MU_{\mathbb{R}}$ and $BP_{\mathbb{R}}$*). *The Hopf elements, the Kervaire classes, and the $\bar{\kappa}$ -family are detected by the Hurewicz maps $\pi_*\mathbb{S} \rightarrow \pi_*MU_{\mathbb{R}}^{C_2}$ and $\pi_*\mathbb{S} \rightarrow \pi_*^{C_2}BP_{\mathbb{R}}^{C_2}$.*

Theorem 6.12. *Let E be an E_∞ H -spectrum. If the H -fixed point spectrum of E detects a class $x \in \pi_*\mathbb{S}$, then the G -fixed point spectrum of $(N_H^G E)$ detects x as well.*

Proof. This follows from the following commutative diagram:

$$\begin{array}{ccccc}
 (N_H^G E)^G & \longrightarrow & (i_H^* N_H^G E)^H = (E \wedge \cdots \wedge E)^H & \longrightarrow & E^H \\
 & & \uparrow & & \nearrow \\
 & & \mathbb{S} & &
 \end{array}$$

The first horizontal map is the map from the G -fixed point to the H -fixed point. The second horizontal map is obtained by the multiplicative structure on E . Taking $\pi_*(-)$ to the entire diagram gives the maps

$$\pi_* \mathbb{S} \rightarrow \pi_*(N_H^G E)^G \rightarrow \pi_*(E^H).$$

Since x maps to a nonzero element in $\pi_*(E^H)$ under the composition map, x must map to a nonzero element in $\pi_*(N_H^G E)^G$ as well. \square

Letting $E = MU_{\mathbb{R}}$ in Theorem 6.12 gives the following:

Corollary 6.13. For any finite group G containing C_2 , the G -fixed point of $MU^{((G))}$ detects the Hopf elements, the Kervaire classes, and the $\bar{\kappa}$ -family.

Remark 6.14. Theorem 6.12 produces the detection tower

$$\begin{array}{c}
 \vdots \\
 \downarrow \\
 \pi_*(MU^{((C_{2^n}))})_{C_{2^n}} \\
 \downarrow \\
 \vdots \\
 \downarrow \\
 \pi_* \mathbb{S} \longrightarrow \pi_*(MU^{((C_8))})_{C_8} \\
 \searrow \quad \downarrow \\
 \pi_*(MU^{((C_4))})_{C_4} \\
 \downarrow \\
 \pi_*(MU_{\mathbb{R}})_{C_2}.
 \end{array}$$

As we go up the tower, the size of the cyclic group increases, and $\pi_*(MU^{((C_{2^n}))})_{C_{2^n}}$ will detect more classes in the homotopy groups of spheres.

6.2. Detection theorem for $E\mathbb{R}(n)$

Recall that the Real Johnson–Wilson theory $E\mathbb{R}(n)$ is constructed from $BP_{\mathbb{R}}$ by killing \bar{v}_i for $i \geq n + 1$ and inverting \bar{v}_n . Its refinement is

$$\begin{array}{ccc} S^0[\bar{v}_1, \bar{v}_2, \dots] & \longrightarrow & BP_{\mathbb{R}} \\ \downarrow & & \downarrow \\ S^0[\bar{v}_1, \dots, \bar{v}_{n-1}, \bar{v}_n^{\pm}] & \longrightarrow & E\mathbb{R}(n). \end{array}$$

To prove the detection theorem for $E\mathbb{R}(n)$, we analyze the composite map

$$\text{MMaySS}(\mathbb{S}) \rightarrow \text{SliceSS}(BP_{\mathbb{R}}) \rightarrow \text{SliceSS}(E\mathbb{R}(n)).$$

Lemma 6.15. *In the slice spectral sequence for $\pi_{\star}^{C_2} E\mathbb{R}(n)$, the classes $a_{\sigma}, \bar{v}_1, \dots, \bar{v}_{n-1}, \bar{v}_n^{\pm}$ are permanent cycles. For $1 \leq k \leq n$, the differentials $d_i(u_{2\sigma}^{2^{k-1}})$ are zero for $i < 2^{k+1} - 1$, and*

$$d_{2^{k+1}-1}(u_{2\sigma}^{2^{k-1}}) = \bar{v}_k a_{\sigma}^{2^{k+1}-1}.$$

The class $u_{2\sigma}^{2^n}$ is a permanent cycle.

Proof. This is immediate by comparing to the slice spectral sequence of $BP_{\mathbb{R}}$ (Proposition 3.4). For the class $u_{2\sigma}^{2^n}$, it is supposed to support a differential to $\bar{v}_{n+1} a_{\sigma}^{2^{n+2}-1}$. However, \bar{v}_{n+1} is 0 in the slice spectral sequence for $E\mathbb{R}(n)$. This implies that $u_{2\sigma}^{2^n}$ is a $d_{2^{n+2}-1}$ -cycle. Furthermore, for degree reasons, there are no classes in the appropriate degrees that can be hit by longer differentials from $u_{2\sigma}^{2^n}$. It follows that the class $u_{2\sigma}^{2^n}$ is a permanent cycle. \square

Theorem 6.16 (Detection theorem for $E\mathbb{R}(n)$).

- (1) For $1 \leq k \leq n$, if the element h_k or h_k^2 in $\text{Ext}_{\mathcal{A}_{\star}}(\mathbb{F}_2, \mathbb{F}_2)$ survives to the E_{∞} -page of the Adams spectral sequence, then its image under the Hurewicz map $\pi_{\star}\mathbb{S} \rightarrow \pi_{\star}E\mathbb{R}(n)^{C_2}$ is nonzero.
- (2) For $1 \leq k \leq n - 1$, if the element $g_k \in \text{Ext}_{\mathcal{A}_{\star}}^{4, 2^{n+2}+2^{n+3}}(\mathbb{F}_2, \mathbb{F}_2)$ survives to the E_{∞} -page of the Adams spectral sequence, then its image under the Hurewicz map $\pi_{\star}\mathbb{S} \rightarrow \pi_{\star}E\mathbb{R}(n)^{C_2}$ is nonzero.

Proof. By Proposition 6.1, Theorem 6.3, and Theorem 6.4, it suffices to show that the classes $\bar{v}_k a_{\sigma}^{2^k-1}$ ($1 \leq k \leq n$), $\bar{v}_k^2 a_{\sigma}^{2(2^k-1)}$ ($1 \leq k \leq n$), and $\bar{v}_{k+1}^4 u_{\sigma}^{2^{k+2}} a_{\sigma}^{4(2^k-1)}$ ($1 \leq k \leq n - 1$) survive to the E_{∞} -page of the slice spectral sequence of $E\mathbb{R}(n)$. This is immediate from Lemma 6.15. \square

Remark 6.17. By the homotopy fixed point theorem ([18, Theorem 10.8]), the spectrum $E\mathbb{R}(n)$ is cofree. This means that $\pi_*E\mathbb{R}(n)^{C_2} \rightarrow \pi_*E\mathbb{R}(n)^{hC_2}$ is an isomorphism. Therefore Theorem 6.16 holds for the homotopy fixed point of $E\mathbb{R}(n)$ as well.

Remark 6.18. The detection theorem for $E\mathbb{R}(n)$ also holds for $v_n^{-1}MU_{\mathbb{R}}$ as it splits as a wedge of suspensions of $E\mathbb{R}(n)$ (with $E\mathbb{R}(n)$ itself being one of the wedge summands).

References

- [1] J.F. Adams, On the non-existence of elements of Hopf invariant one, *Ann. of Math.* (2) 72 (1960) 20–104.
- [2] S. Araki, Orientations in τ -cohomology theories, *Jpn. J. Math. (N. S.)* 5 (2) (1979) 403–430.
- [3] M.F. Atiyah, K -theory and reality, *Quart. J. Math. Oxford Ser. (2)* 17 (1966) 367–386.
- [4] M.G. Barratt, M.E. Mahowald, M.C. Tangora, Some differentials in the Adams spectral sequence. II, *Topology* 9 (1970) 309–316.
- [5] M.G. Barratt, J.D.S. Jones, M.E. Mahowald, Relations amongst Toda brackets and the Kervaire invariant in dimension 62, *J. Lond. Math. Soc. (2)* 30 (3) (1984) 533–550.
- [6] W. Browder, The Kervaire invariant of framed manifolds and its generalization, *Ann. of Math.* (2) 90 (1969) 157–186.
- [7] R.R. Bruner, J.P. May, J.E. McClure, M. Steinberger, H_{∞} Ring Spectra and Their Applications, *Lecture Notes in Mathematics*, vol. 1176, Springer-Verlag, Berlin, 1986.
- [8] D. Dugger, An Atiyah–Hirzebruch spectral sequence for KR-theory, *K-Theory* 35 (3) (2005) 213–256.
- [9] M. Fujii, Cobordism theory with reality, *Math. J. Okayama Univ.* 18 (2) (1975–1976) 171–188.
- [10] P.G. Goerss, M.J. Hopkins, Moduli spaces of commutative ring spectra, in: *Structured Ring Spectra*, in: *London Math. Soc. Lecture Note Ser.*, vol. 315, Cambridge Univ. Press, Cambridge, 2004, pp. 151–200.
- [11] J.P.C. Greenlees, *Adams Spectral Sequences in Equivariant Topology*, Ph.D. Thesis, University of Cambridge, 1985.
- [12] J.P.C. Greenlees, Stable maps into free G -spaces, *Trans. Amer. Math. Soc.* 310 (1) (1988) 199–215.
- [13] J.P.C. Greenlees, The power of mod p Borel homology, in: *Homotopy Theory and Related Topics*, Kinoshita, 1988, in: *Lecture Notes in Math.*, vol. 1418, Springer, Berlin, 1990, pp. 140–151.
- [14] J. Hahn, X. Shi, Real orientations of Lubin–Tate spectra, *ArXiv e-prints*, 2017.
- [15] M.A. Hill, L. Meier, The C_2 -spectrum $\mathrm{Tmf}_1(3)$ and its invertible modules, *Algebr. Geom. Topol.* 17 (4) (2017) 1953–2011.
- [16] M.A. Hill, M.J. Hopkins, D.C. Ravenel, The arf-Kervaire invariant problem in algebraic topology: introduction, *Curr. Dev. Math.* 2009 (2010) 23–57.
- [17] M.A. Hill, M.J. Hopkins, D.C. Ravenel, The Arf–Kervaire problem in algebraic topology: sketch of the proof, *Curr. Dev. Math.* 2010 (2011) 1–44.
- [18] M.A. Hill, M.J. Hopkins, D.C. Ravenel, On the nonexistence of elements of Kervaire invariant one, *Ann. of Math.* (2) 184 (1) (2016) 1–262.
- [19] M.A. Hill, M.J. Hopkins, D.C. Ravenel, The Slice spectral sequence for the C_4 analogue of Real K -theory, *arXiv:1502.07611*, February 2016.
- [20] P. Hu, I. Kriz, Real-oriented homotopy theory and an analogue of the Adams–Novikov spectral sequence, *Topology* 40 (2) (2001) 317–399.
- [21] D.C. Isaksen, G. Wang, Z. Xu, More stable stems, In preparation.
- [22] N. Kitchloo, W.S. Wilson, On fibrations related to real spectra, in: *Proceedings of the Nishida Fest*, Kinoshita 2003, in: *Geom. Topol. Monogr.*, vol. 10, Geom. Topol. Publ., Coventry, 2007, pp. 237–244.
- [23] N. Kitchloo, W.S. Wilson, The second real Johnson–Wilson theory and nonimmersions of RP^n , *Homology, Homotopy Appl.* 10 (3) (2008) 223–268.
- [24] N. Kitchloo, W.S. Wilson, The second real Johnson–Wilson theory and nonimmersions of RP^n , II *Homology, Homotopy Appl.* 10 (3) (2008) 269–290.
- [25] N. Kitchloo, V. Lorman, W.S. Wilson, Landweber flat real pairs and $ER(n)$ -cohomology, *ArXiv e-prints*, March 2016.
- [26] N. Kitchloo, V. Lorman, W.S. Wilson, The $ER(2)$ -cohomology of $B\mathbb{Z}/(2^q)$ and CP^n , *ArXiv e-prints*, May 2016.

- [27] P.S. Landweber, Conjugations on complex manifolds and equivariant homotopy of MU , *Bull. Amer. Math. Soc.* 74 (1968) 271–274.
- [28] G. Li, X.D. Shi, G. Wang, Z. Xu, The $er(2)$ cohomology of real projective spaces, 2017, In preparation.
- [29] W.-H. Lin, $\text{Ext}_A^{4,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $\text{Ext}_A^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$, *Topology Appl.* 155 (5) (2008) 459–496.
- [30] V. Lorman, The real Johnson–Wilson cohomology of $\mathbb{C}P^\infty$, ArXiv e-prints, September 2015.
- [31] M. Mahowald, M. Tangora, Some differentials in the Adams spectral sequence, *Topology* 6 (1967) 349–369.
- [32] H. Miller, Kervaire invariant one [after M.A. Hill, M.J. Hopkins, and D.C. Ravenel], ArXiv e-prints, April 2011.
- [33] O. Nakamura, On the squaring operations in the May spectral sequence, *Mem. Fac. Sci., Kyushu Univ., Ser. A, Math.* 26 (1972) 293–308.
- [34] D.C. Ravenel, *Complex Cobordism and Stable Homotopy Groups of Spheres*, American Mathematical Soc., 2003.
- [35] C. Rezk, Notes on the Hopkins–Miller theorem, in: *Homotopy Theory via Algebraic Geometry and Group Representations*, Evanston, IL, 1997, in: *Contemp. Math.*, vol. 220, Amer. Math. Soc., Providence, RI, 1998, pp. 313–366.
- [36] J. Ullman, *On the Regular Slice Spectral Sequence*, Ph.D. Thesis, Massachusetts Institute of Technology, 2013.