

A SIMPLE UNIVERSAL PROPERTY OF THOM RING SPECTRA

OMAR ANTOLÍN-CAMARENA AND TOBIAS BARTHEL

ABSTRACT. We give a simple universal property of the multiplicative structure on the Thom spectrum of an n -fold loop map, obtained as a special case of a characterization of the algebra structure on the colimit of a lax \mathcal{O} -monoidal functor. This allows us to relate Thom spectra to \mathbb{E}_n -algebras of a given characteristic in the sense of Szymik. As applications, we recover the Hopkins–Mahowald theorem realizing $H\mathbb{F}_p$ and $H\mathbb{Z}$ as Thom spectra, and compute the topological Hochschild homology and the cotangent complex of various Thom spectra.

CONTENTS

1. Introduction	1
1.1. Motivation	1
1.2. Outline and main results	2
Acknowledgments	3
2. Colimits of lax functors	3
2.1. Ordinary lax functors	3
2.2. A few words about ∞ -operads	4
2.3. The \mathcal{O} -algebra structure on the colimit of a lax \mathcal{O} -monoidal functor	7
3. The universal multiplicative property of the Thom spectrum	8
3.1. The universal property and its consequences	9
3.2. \mathbb{E}_n -orientations	11
4. Thom spectra and versal \mathbb{E}_n -algebras	14
4.1. Characteristics of ring spectra	14
4.2. Thom spectra as versal characteristic χ algebras	16
5. Applications	18
5.1. The Hopkins–Mahowald theorem I: $H\mathbb{F}_p$	18
5.2. The Hopkins–Mahowald theorem II: $H\mathbb{Z}$	19
5.3. Topological Hochschild homology and the cotangent complex	21
References	22

1. INTRODUCTION

1.1. Motivation. Many spectra of interest fall in one of two classes: either they contain important geometric or arithmetic information and are difficult to compute with, e.g., the sphere spectrum or various algebraic K -theory spectra, or they belong

Date: June 14, 2017.

2010 Mathematics Subject Classification. 55N20, 55P42, 55P43, 55P48.

to a family of what Hopkins calls “designer homotopy types”, like Eilenberg–Mac Lane spectra or Brown–Gitler spectra.

Thom spectra are remarkable in that they often belong to the intersection: Classical examples include cobordism spectra, whose homotopy groups and other invariants often turn out to be completely computable. As another beautiful example, Mahowald [Mah79] proves that the Eilenberg–Mac Lane spectra HF_2 and $H\mathbb{Z}$ can be realized as Thom spectra, an observation which he applies in his construction of new elements in the homotopy groups of spheres [Mah77]. Similar ideas also play a role in the proof of the nilpotence theorem by Devinatz, Hopkins, and Smith [DHS88].

From a more conceptual point of view, work of Lewis and May [LMSM86], Sullivan [Sul05], and May and Sigurdsson [MS06] places the theory of Thom spectra in the context of parametrized spectra and thus reveals the underlying geometry of the construction. We are interested here in how this framework accounts for some of the good algebraic properties of Thom spectra.

1.2. Outline and main results. In [ABG⁺14a] and [ABG11], the authors describe a convenient model of Thom spectra as homotopy colimits of diagrams whose shape is given by the base space. Their approach is closely related and essentially equivalent to the earlier models cited above, but is expressed in the language of ∞ -categories. The study of multiplicative structures on Thom spectra can thus take place in the setting of \mathcal{O} -monoidal functors between \mathcal{O} -monoidal ∞ -categories.

Section 2 first explains the situation in ordinary category theory that we then extend to \mathcal{O} -monoidal ∞ -categories, proving our general characterization of the algebra structure of colimits of lax \mathcal{O} -monoidal functors in Theorem 2.13. This is used in Section 3 to deduce the universal property of the multiplicative structure of Thom spectra of \mathbb{E}_n -maps, see Theorem 3.5.

Theorem. *If X is an \mathbb{E}_n -space and $f: X \rightarrow \text{Pic}(S^0)$ an \mathbb{E}_n -map, then the space $\text{Map}_{\text{Alg}_{S^0}^{\mathbb{E}_n}}(Mf, A)$ is equivalent to the space of \mathbb{E}_n -lifts of f indicated below:*

$$\begin{array}{ccc} & & \text{Pic}(S^0)_{/A} \\ & \nearrow & \downarrow \\ X & \xrightarrow{f} & \text{Pic}(S^0). \end{array}$$

This generalizes earlier structural results about Thom spectra proven by Lewis [LMSM86] and in [ABG⁺14a]. We then show that any n -fold loop map with Thom spectrum Mf is canonically \mathbb{E}_{n-1} Mf -orientable, thereby establishing a structured version of the Thom isomorphism. Moreover, we deduce a theorem of Chadwick and Mandell [CM15], describing \mathbb{E}_n -orientations.

We start Section 4 by introducing a notion of characteristic for \mathbb{E}_n -algebras in spectra, a straightforward extension of the \mathbb{E}_∞ -case studied previously by Szymik [Szy14, Szy13]; see also [Bak] for related ideas. It is easy to construct a weakly initial example of an algebra of characteristic χ , denoted $S^0 //_{\mathbb{E}_n} \chi$. We then show that these algebras satisfy the same universal property as certain Thom spectra naturally corresponding to them, which gives our second main result, Theorem 4.10.

Theorem. *For any n and any $k \geq 1$ and $f: S^k \rightarrow BGL_1 S^0$ with corresponding n -fold loop map $\bar{f}: \Omega^n \Sigma^n S^k \rightarrow BGL_1 R$ and with associated characteristic $\chi = \chi(f)$, there is an equivalence $M\bar{f} = S^0 //_{\mathbb{E}_n} \chi$.*

This theorem establishes a connection between Thom spectra and versal algebras which allows to transfer results proven for one class to the other. We illustrate this idea in Section 5. For instance, by specializing to $n = 2$ and $(1 - p) = f_p: S^1 \rightarrow BGL_1 S_p^0$ and using Cohen's computation [CLM76] of the mod p cohomology of the free \mathbb{E}_2 -algebra on S^1 , we recover the Hopkins–Mahowald theorem

$$M\bar{f}_p = H\mathbb{F}_p.$$

Our argument simplifies the proofs given by Priddy [Pri78] and Cohen–May–Taylor [CMT81] in that it does not require Kochman's calculations of the homology operations of BO or computations of Steenrod operations, respectively. Furthermore, it allows us to directly deduce the identification of $H\mathbb{Z}$ as an \mathbb{E}_2 -Thom ring spectrum. Finally, we describe the topological Hochschild homology and the \mathbb{E}_n -cotangent complex of the \mathbb{E}_n -algebras considered above.

Acknowledgments. We are grateful to Jacob Lurie for suggesting to use the universal property of Thom ring spectra to reprove the Hopkins–Mahowald theorem. We would like to thank Jon Beardsley, Andrew Blumberg, David Gepner, Gijs Heuts, Mike Hopkins, Tyler Lawson, and Chris Schommer-Pries for helpful conversations. The second author also thanks the Max Planck Institute for Mathematics for its hospitality.

2. COLIMITS OF LAX FUNCTORS

2.1. Ordinary lax functors. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a lax monoidal functor between two (ordinary) monoidal categories. Recall that this means that we are given a family of morphisms $FX \otimes FY \rightarrow F(X \otimes Y)$ natural in X and Y which are compatible with the associativity constraints in \mathcal{C} and \mathcal{D} . If \mathcal{C} is small and \mathcal{D} is *monoidally cocomplete*, by which we mean that \mathcal{D} is cocomplete and $\otimes: \mathcal{D} \times \mathcal{D} \rightarrow \mathcal{D}$ preserves colimits in each variable separately, then $M := \text{colim} F$ acquires the structure of an algebra object in \mathcal{D} . The multiplication $\mu: M \otimes M \rightarrow M$ is obtained from the lax structure of F as follows: the composite morphisms

$$FX \otimes FY \rightarrow F(X \otimes Y) \rightarrow M,$$

induce a morphism

$$M \otimes M \cong \text{colim}_{(X,Y) \in \mathcal{C} \times \mathcal{C}} (FX \otimes FY) \rightarrow M,$$

where the first isomorphism comes from the assumption that tensor products in \mathcal{D} commute with colimits in each variable separately. This is an analogue in ordinary category theory of Lewis result that the Thom spectrum of a loop map is an \mathbb{E}_1 -ring spectrum.

If we are additionally given an algebra object A in \mathcal{D} , the slice category $\mathcal{D}_{/A}$ acquires a monoidal structure in which $(W \xrightarrow{f} A) \otimes (Z \xrightarrow{g} A)$ is given by the composite

$$W \otimes Z \xrightarrow{f \otimes g} A \otimes A \xrightarrow{\mu_A} A.$$

The projection functor $\pi: \mathcal{D}_{/A} \rightarrow \mathcal{D}$ is, of course, monoidal.

The universal property we will give for the multiplicative structure of Thom spectra is an analogue of the following fact: morphisms $M \rightarrow A$ of algebra objects in \mathcal{D} are in one to one correspondence with lax monoidal lifts of F through π :

$$\begin{array}{ccc} & & \mathcal{D}/A \\ & \nearrow & \downarrow \pi \\ \mathcal{C} & \xrightarrow{F} & \mathcal{D} \end{array}$$

In this setting, this fact is straightforward to verify: lax monoidal lifts are “lax monoidal cocones” over F with vertex A , and such cocones induce algebra morphisms $M = \operatorname{colim} F \rightarrow A$.

In Section 2.3 we will generalize this discussion in two ways: first, we will work with ∞ -categories instead of ordinary categories; second, instead of considering only monoidal ∞ -categories we will work with \mathcal{O} -monoidal ∞ -categories for an arbitrary ∞ -operad \mathcal{O} . (For applications we will take \mathcal{O} to be the \mathbb{E}_n operad for some $0 \leq n \leq \infty$.) Before doing that, however, let us recall some definitions from the theory of ∞ -operads.

2.2. A few words about ∞ -operads. We recall some definitions about ∞ -operads from chapters 2 and 3 of Lurie’s *Higher Algebra* [Lur17]. The theory of ∞ -operads developed there is the $(\infty, 1)$ -generalization of the theory of colored operads or symmetric multicategories. So an ∞ -operad \mathcal{O} has a collection of objects and for every finite (unordered) family $\{X_i\}_{i \in I}$ of objects and any object Y there is a space of morphisms $\operatorname{Mul}_{\mathcal{O}}(\{X_i\}, Y)$. The definitions in [Lur17] encode this data indirectly: an ∞ -operad \mathcal{O} is specified by an ∞ -category \mathcal{O}^{\otimes} and a functor $\mathcal{O}^{\otimes} \rightarrow N(\operatorname{Fin}_*)$ to the category of finite pointed sets satisfying certain conditions. Morphisms of ∞ -operads $\mathcal{O} \rightarrow \mathcal{O}'$ are functors over $N(\operatorname{Fin}_*)$ (that is, functors $\mathcal{O}^{\otimes} \rightarrow \mathcal{O}'^{\otimes}$ that form a commutative triangle with the structure maps to $N(\operatorname{Fin}_*)$) which satisfy one additional condition. See [Lur17, Definitions 2.1.1.10 and 2.1.2.7] for the precise definition.

Remark 2.1. To give some intuition for how the definition captures the notion of colored operad, think of the case of symmetric monoidal categories. These can be regarded as a special kind of operad in which $\operatorname{Mul}_{\mathcal{C}}(\{X_i\}_{i \in I}, Y) = \operatorname{Hom}_{\mathcal{C}}(\bigotimes_{i \in I} X_i, Y)$; but they can also be thought of as commutative monoids in the category of all categories, which using Segal’s idea of Γ -spaces, can be encoded as appropriately weak functors $\operatorname{Fin}_* \rightarrow \operatorname{Cat}$ satisfying certain conditions. Applying the (covariant) Grothendieck construction to such a functor one obtains a coCartesian fibration $\mathcal{C}^{\otimes} \rightarrow \operatorname{Fin}_*$. The definition of symmetric monoidal ∞ -category is exactly what this would suggest: a coCartesian fibration of ∞ -categories $\mathcal{C}^{\otimes} \rightarrow N(\operatorname{Fin}_*)$ satisfying an analogue of the Segal condition, and the definition of ∞ -operad is a generalization of this.

An \mathcal{O} -monoidal ∞ -category is defined to be a coCartesian fibration $\mathcal{C}^{\otimes} \rightarrow \mathcal{O}^{\otimes}$ such that the composite $\mathcal{C}^{\otimes} \rightarrow \mathcal{O}^{\otimes} \rightarrow N(\operatorname{Fin}_*)$ presents \mathcal{C} as an ∞ -operad. If \mathcal{O} is the \mathbb{E}_1 or \mathbb{E}_{∞} operad, this recovers the notion of monoidal ∞ -category or symmetric monoidal ∞ -category, respectively.

Remark 2.2. Again, the intuition for this definition comes from the Grothendieck construction: morally, one would want to say that \mathcal{O} -monoidal categories are \mathcal{O} -algebras in categories, or more precisely, morphisms of operads from \mathcal{O} to the $(\infty, 2)$ -category of ∞ -categories equipped with its symmetric monoidal structure. But instead of appealing to the theory of $(\infty, 2)$ -categories, it is much easier to describe what the result of applying a version of the Grothendieck construction would be and adopt the resulting coCartesian fibration as the definition.

Warning 2.3. In the case that \mathcal{O} is an operad with a single color X , an \mathcal{O} -monoidal ∞ -category $p: \mathcal{C}^\otimes \rightarrow \mathcal{O}^\otimes$ can be thought of as an ∞ -category $\mathcal{C} = p^{-1}(X)$ equipped with functors $\mathcal{C}^n \rightarrow \mathcal{C}$ for each n -ary operation in \mathcal{O} . Of course, something similar is true for multicolored \mathcal{O} , except that instead of a single underlying ∞ -category, there is an ∞ -category $\mathcal{C}_X = p^{-1}(X)$ for each object X of \mathcal{O} , and a functor $\prod \mathcal{C}_{X_i} \rightarrow \mathcal{C}_Y$ for each operation in $\text{Mul}_{\mathcal{O}}(\{X_i\}, Y)$. The reader can pretend that whenever we mention an \mathcal{O} -monoidal ∞ -category, \mathcal{O} has a single object without missing out on anything essential.

Between two \mathcal{O} -monoidal ∞ -categories \mathcal{C} and \mathcal{D} we can consider \mathcal{O} -monoidal functors and lax \mathcal{O} -monoidal functors. Lax \mathcal{O} -monoidal functors are simply morphisms of ∞ -operads $\mathcal{C} \rightarrow \mathcal{D}$ over \mathcal{O} ; they form an ∞ -category denoted $\text{Fun}_{\mathcal{O}}^{\text{lax}}(\mathcal{C}, \mathcal{D})$ or $\text{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D})$. The lax \mathcal{O} -monoidal functors that send coCartesian morphisms to coCartesian morphisms are the \mathcal{O} -monoidal functors; they form an ∞ -category $\text{Fun}_{\mathcal{O}}^\otimes(\mathcal{C}, \mathcal{D})$.

Definition 2.4. Let \mathcal{O}^\otimes be an ∞ -operad. An \mathcal{O} -monoidally cocomplete ∞ -category is an \mathcal{O} -monoidal category, that is, a coCartesian fibration of ∞ -operads $q: \mathcal{C}^\otimes \rightarrow \mathcal{O}^\otimes$, such that q is compatible with all small colimits in the sense of [Lur17, Definition 3.1.1.18]. This means that for every object X of the underlying ∞ -category \mathcal{O} of \mathcal{O}^\otimes , the ∞ -category $\mathcal{C}_X := q^{-1}(X)$ is cocomplete, and for every morphism $f \in \text{Mul}_{\mathcal{O}}(\{X_i\}_{1 \leq i \leq n}, Y)$, the functor $\otimes_f: \prod_{1 \leq i \leq n} \mathcal{C}_{X_i} \rightarrow \mathcal{C}_Y$ preserves colimits in each variable separately.

Recall that the *core* of an ∞ -category \mathcal{C} is the maximal ∞ -groupoid contained in \mathcal{C} . If \mathcal{C} is incarnated as a quasi-category, then the core \mathcal{C}^\simeq is easily described as a Kan complex: it is the subsimplicial set of \mathcal{C} consisting of simplices all of whose edges are invertible morphisms of \mathcal{C} . If \mathcal{C} has an \mathcal{O} -monoidal structure, then, as expected from the theory of ordinary monoidal categories, \mathcal{C}^\simeq inherits an \mathcal{O} -monoidal structure as well. More precisely, we have:

Proposition 2.5. *Let \mathcal{O} be an ∞ -operad and let $q: \mathcal{C}^\otimes \rightarrow \mathcal{O}^\otimes$ exhibit \mathcal{C} as an \mathcal{O} -monoidal ∞ -category. Define $\mathcal{C}_{\text{coCart}}^\otimes$ to be subcategory of \mathcal{C}^\otimes spanned by the q -coCartesian morphisms. The restriction of q to this subcategory, $\tilde{q}: \mathcal{C}_{\text{coCart}}^\otimes \rightarrow \mathcal{O}^\otimes$, then exhibits $\mathcal{C}_{\text{coCart}}^\otimes$ as an \mathcal{O} -monoidal category such that for each object X of \mathcal{O} , the underlying ∞ -category $(\mathcal{C}_{\text{coCart}}^\otimes)_X$ is the core of \mathcal{C}_X .*

Proof. We will show that:

- (1) \tilde{q} is a coCartesian fibration.
- (2) $(\mathcal{C}_{\text{coCart}}^\otimes)_X = \mathcal{C}_X^\simeq$ for any object X of \mathcal{O} or even of \mathcal{O}^\otimes .
- (3) The composite $\mathcal{C}_{\text{coCart}}^\otimes \xrightarrow{\tilde{q}} \mathcal{O}^\otimes \rightarrow N(\text{Fin}_*)$ exhibits $\mathcal{C}_{\text{coCart}}^\otimes$ as an ∞ -operad.

The items 1 and 3 guarantee $\mathcal{C}_{\text{coCart}}^\otimes$ is an \mathcal{O} -monoidal category. We have placed item 2 in the middle because it will be used in the proof of 3.

Proof of 1. First we need to show that \tilde{q} is an inner fibration of simplicial sets, i.e., we must show it has the right lifting property against inner horn inclusions. But q has this lifting property and the fillers only have 1-simplices not present in the horn in a single case: $\Lambda_1^2 \hookrightarrow \Delta^2$. Since coCartesian morphisms are closed under composition, the filler for q will also serve as a filler for \tilde{q} .

Now we will show that all morphisms in $\mathcal{C}_{\text{coCart}}^\otimes$ are \tilde{q} -coCartesian. This is very similar to the above proof that \tilde{q} is an inner fibration. A morphism f is coCartesian if and only if for every $n \geq 2$ and every commutative diagram

$$\begin{array}{ccc}
 \Delta^1 & & \\
 i_{01} \downarrow & \searrow f & \\
 \Lambda_n^0 & \longrightarrow & \mathcal{C}_{\text{coCart}}^\otimes \\
 \downarrow & \nearrow & \downarrow \\
 \Delta^n & \longrightarrow & \mathcal{O}^\otimes
 \end{array}$$

there is a dotted arrow that makes the diagram commute (this is dual to [Lur09, Remark 2.4.1.4]). As above, we can always take the same filler as for the corresponding diagram with \mathcal{C}^\otimes in place of $\mathcal{C}_{\text{coCart}}^\otimes$: indeed, the simplex Δ^n has no 1-simplices absent from Λ_n^0 unless $n = 2$; and if $n = 2$, we can still use the same filler because if g and $h \circ g$ are coCartesian then so is h (this is the dual of [Lur09, Proposition 2.4.1.7]).

Proof of 2. The morphisms in $(\mathcal{C}_{\text{coCart}}^\otimes)_X$ project to the identity morphism of $X \in \mathcal{O}^\otimes$ by definition. Morphisms projecting to the identity are coCartesian if and only if they are invertible [Lur09, Proposition 2.4.1.5].

Proof of 3. If $p: \mathcal{O}^\otimes \rightarrow N(\text{Fin}_*)$ is an ∞ -operad and $\langle n \rangle \in \text{Fin}_*$ denotes the pointed set $\{*, 1, 2, \dots, n\}$, there is an equivalence $\mathcal{O}_{\langle n \rangle}^\otimes \rightarrow \mathcal{O}^n$. What we need to check according to [Lur17, Proposition 2.1.2.12] is that for every $X \in \mathcal{O}_{\langle n \rangle}^\otimes$ with corresponding sequence $(X_1, \dots, X_n) \in \mathcal{O}^n$, we get an equivalence $(\mathcal{C}_{\text{coCart}}^\otimes)_X \cong \prod_{i=1}^n (\mathcal{C}_{\text{coCart}}^\otimes)_{X_i}$ induced by taking coCartesian lifts of the inert morphisms $X \rightarrow X_i$. For the purposes of this argument, the definition of inert does not matter: we simply note that by the same proposition applied to \mathcal{C}^\otimes , we *do* have such an equivalence for \mathcal{C}^\otimes instead of $\mathcal{C}_{\text{coCart}}^\otimes$. This equivalence restricts to the required equivalence because coCartesian morphisms are closed under composition. \square

Remark 2.6. This proposition encodes the \mathcal{O} -monoidal structure of the space \mathcal{C}^\simeq as an ∞ -category $\mathcal{C}_{\text{coCart}}^\otimes$ with a coCartesian fibration over \mathcal{O}^\otimes . A more direct way to encode an \mathcal{O} -space is as an \mathcal{O} -algebra in the ∞ -category of spaces. These two forms are equivalent: the coCartesian fibration \tilde{q} constructed in the proposition has the feature that all morphisms in the domain are coCartesian, and thus by [Lur17, Proposition 2.4.2.4], \tilde{q} is a left fibration, and therefore classifies a functor $\mathcal{O}^\otimes \rightarrow \text{Spaces}$ which exhibits \mathcal{C}^\simeq as an \mathcal{O} -monoid in spaces. This is the correspondence of [Lur17, Example 2.4.2.4] restricted to left fibrations.

Remark 2.7. That the core of an \mathcal{O} -monoidal ∞ -category inherits an \mathcal{O} -monoidal structure has been used implicitly before (see for example, [ABG11, Definition 8.5 and Remark 8.6] or [MS16, Proposition 2.2.3]), but as far the authors know, no

explicit description has appeared in the literature, which is why we feel justified in giving it in such detail.

2.3. The \mathcal{O} -algebra structure on the colimit of a lax \mathcal{O} -monoidal functor.

For \mathcal{C} an \mathcal{O} -monoidal category, the ∞ -category of \mathcal{O} -algebras in \mathcal{C} , denoted by $\text{Alg}_{/\mathcal{O}}(\mathcal{C})$, is defined to be $\text{Alg}_{\mathcal{O}/\mathcal{O}}(\mathcal{C})$, the ∞ -category of lax \mathcal{O} -monoidal functors $\mathcal{O} \rightarrow \mathcal{C}$. Unwinding the definitions we see that \mathcal{O} -algebras in \mathcal{C} are sections of the coCartesian fibration $\mathcal{C}^\otimes \rightarrow \mathcal{O}^\otimes$ presenting the \mathcal{O} -monoidal structure on \mathcal{C} ; more precisely \mathcal{O} -algebras are those sections which are also maps of ∞ -operads, i.e., maps sending inert morphisms to inert morphisms.

Theorem 2.8. *Let \mathcal{O} be an ∞ -operad, \mathcal{C} be a small \mathcal{O} -monoidal ∞ -category, and \mathcal{D} be an \mathcal{O} -monoidally cocomplete ∞ -category. If $F: \mathcal{C}^\otimes \rightarrow \mathcal{D}^\otimes$ is a lax \mathcal{O} -monoidal functor, then there is an \mathcal{O} -algebra in \mathcal{D} given by a functor $M: \mathcal{O} \rightarrow \mathcal{D}$ such that for every object X of \mathcal{O} , $M(X) = \text{colim}(F_X: \mathcal{C}_X \rightarrow \mathcal{D}_X)$.*

Remark 2.9. In the case that \mathcal{O} has a single object, the conclusion should be thought of as saying that the colimit of the functor $F: \mathcal{C} \rightarrow \mathcal{D}$ has a canonical structure of an \mathcal{O} -algebra in \mathcal{D} .

Remark 2.10. This theorem is due to Lewis [LMSM86, Section IX.7] in the case that \mathcal{C} is an ∞ -groupoid and \mathcal{D} is the category of spectra — and \mathcal{O} has a single color. In [ABG11], the authors prove a similar result under slightly stronger assumptions: that \mathcal{O} is coherent and \mathcal{D} is an \mathcal{O} -algebra object in the ∞ -category of presentable ∞ -categories (and functors which are left adjoints).

Proof. Let $p: \mathcal{C}^\otimes \rightarrow \mathcal{O}^\otimes$ and $q: \mathcal{D}^\otimes \rightarrow \mathcal{O}^\otimes$ be the coCartesian fibrations of ∞ -operads presenting \mathcal{C} and \mathcal{D} as \mathcal{O} -monoidal categories. Since \mathcal{D} is assumed to be \mathcal{O} -monoidally cocomplete, by [Lur17, Proposition 3.1.1.20], there exists an operadic left Kan extension M to F along p relative to q :

$$\begin{array}{ccc} \mathcal{C}^\otimes & \xrightarrow{F} & \mathcal{D}^\otimes \\ p \downarrow & \nearrow M & \downarrow q \\ \mathcal{O}^\otimes & \xrightarrow{\text{id}} & \mathcal{O}^\otimes. \end{array}$$

This M is the desired algebra structure on $\text{colim} F$. To show this we need only check that for each object X in \mathcal{O} , $M(X) \cong \text{colim}(F|_{\mathcal{C}_X}: \mathcal{C}_X \rightarrow \mathcal{D}_X)$. According to the $n = 1$ case of [Lur17, Theorem 3.1.2.3], the diagram $((\mathcal{C}_{\text{act}}^\otimes)_{/X})^\triangleright \rightarrow \mathcal{D}^\otimes$ induced from the commutative triangle in the top left half of the above square is an operadic colimit diagram relative to q . Now [Lur17, Proposition 3.1.1.16] states that an operadic colimit relative to a coCartesian fibration (here, q) becomes a colimit in each underlying ∞ -category \mathcal{D}_X . \square

Corollary 2.11. *There exists an algebra structure on $\text{colim} F$ for each F . In fact, there is a left adjoint to the functor $p^* = (- \circ p): \text{Alg}_{/\mathcal{O}}(\mathcal{D}) \rightarrow \text{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D})$ that gives this algebra structure functorially.*

Proof. This follows from the proof of Theorem 2.8 by [Lur17, Corollary 3.1.3.4]. \square

Recall that for an \mathcal{O} -algebra $A \in \text{Alg}_{/\mathcal{O}}(\mathcal{D})$, the slice category $\mathcal{D}_{/A}$ has the structure of an \mathcal{O} -monoidal ∞ -category; in other words, $\mathcal{D}_{/A}$ is the underlying ∞ -category of an \mathcal{O} -monoidal ∞ -category $\mathcal{D}_{/A}^\otimes \rightarrow \mathcal{O}^\otimes$ (see [Lur17, Theorem 2.2.2.4]). The slice category has the following universal property:

Lemma 2.12. *Let \mathcal{C} and \mathcal{D} be \mathcal{O} -monoidal ∞ -categories, A be an \mathcal{O} -algebra in \mathcal{D} , and $F: \mathcal{C}^\otimes \rightarrow \mathcal{D}^\otimes$ be a lax \mathcal{O} -monoidal functor. Lax \mathcal{O} -monoidal lifts of F through the projection $\mathcal{D}^\otimes_{/A\mathcal{O}} \rightarrow \mathcal{D}^\otimes$ then correspond to \mathcal{O} -monoidal natural transformations $F \rightarrow A \circ p$, where $p: \mathcal{C}^\otimes \rightarrow \mathcal{O}^\otimes$ exhibits \mathcal{C} as \mathcal{O} -monoidal. More precisely, there is a homotopy equivalence*

$$\mathrm{Map}_{\mathrm{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D})}(F, A \circ p) \cong \{F\} \times_{\mathrm{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D})} \mathrm{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D}/A).$$

Proof. Unwinding the definition of the slice category [Lur17, Section 2.2.2], we see that the quasi-category $\{F\} \times_{\mathrm{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D})} \mathrm{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D}/A)$ is the simplicial set such that morphisms into it from another simplicial set Z are in bijection with morphisms $\tilde{F}: \mathcal{C}^\otimes \times Z^\triangleright \rightarrow \mathcal{D}^\otimes$ such that

- (1) The following diagram commutes:

$$\begin{array}{ccccc} \mathcal{C}^\otimes & \xrightarrow{i} & \mathcal{C}^\otimes \times Z^\triangleright & \xrightarrow{\pi_1} & \mathcal{C}^\otimes \\ p \downarrow & & \tilde{F} \downarrow & & \downarrow p \\ \mathcal{O}^\otimes & \xrightarrow{A} & \mathcal{D}^\otimes & \xrightarrow{q} & \mathcal{O}^\otimes \end{array}$$

where

- $p: \mathcal{C}^\otimes \rightarrow \mathcal{O}^\otimes$ and $q: \mathcal{D}^\otimes \rightarrow \mathcal{O}^\otimes$ are the coCartesian fibrations of ∞ -operads that exhibit \mathcal{C} and \mathcal{D} as \mathcal{O} -monoidal ∞ -categories,
 - i includes \mathcal{C}^\otimes as $\mathcal{C}^\otimes \times \{c\}$ where c is the cocone point of Z^\triangleright ,
- (2) $\tilde{F}|_{\mathcal{C}^\otimes \times Z} = F \circ \pi_1$ (note that in particular, this restriction is independent of the second argument), and
- (3) \tilde{F} sends pairs (f, z) with f an inert morphism of \mathcal{C}^\otimes to inert morphisms of \mathcal{D}^\otimes .

This is readily seen to be isomorphic to the simplicial set that is denoted by

$$\mathrm{Hom}_{\mathrm{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D})}^R(F, A \circ p)$$

in [Lur09, Section 1.2.2], a model of the mapping space $\mathrm{Map}_{\mathrm{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D})}(F, A \circ p)$. \square

Theorem 2.13. *The \mathcal{O} -algebra M from Theorem 2.8 is characterized by the following universal property: For any \mathcal{O} -algebra A in \mathcal{D} the space of \mathcal{O} -algebra maps $\mathrm{Map}_{\mathrm{Alg}_{\mathcal{C}/\mathcal{O}}(\mathcal{D})}(M, A)$ is homotopy equivalent to the space of lax \mathcal{O} -monoidal lifts*

$$\begin{array}{ccc} & & \mathcal{D}^\otimes_{/A\mathcal{O}} \\ & \nearrow & \downarrow \\ \mathcal{C}^\otimes & \longrightarrow & \mathcal{D}^\otimes \end{array}$$

Proof. Just combine Corollary 2.11 and Lemma 2.12. \square

3. THE UNIVERSAL MULTIPLICATIVE PROPERTY OF THE THOM SPECTRUM

Let $n \geq 0$ and R be an \mathbb{E}_{n+1} -ring spectrum, then the ∞ -category of (left) R -module spectra, Mod_R , can be equipped with the structure of an \mathbb{E}_n -monoidal ∞ -category, see [Lur17, Corollary 5.1.2.6]. This means that we can talk about \mathbb{E}_n -algebra objects in Mod_R . We will call these \mathbb{E}_n R -algebras and denote the ∞ -category they form by $\mathrm{Alg}_R^{\mathbb{E}_n}$, that is, we set $\mathrm{Alg}_R^{\mathbb{E}_n} = \mathrm{Alg}_{/\mathbb{E}_n}(\mathrm{Mod}_R)$. Our multiplicative Thom spectra will be \mathbb{E}_n R -algebras.

As in [ABG⁺14a], we associate two ∞ -groupoids to R :

- Let $\text{Pic}(R)$ be the subcategory of invertible R -modules and all equivalences between them, i.e., the core of the subcategory of Mod_R on the invertible objects.
- Let BGL_1R denote the subcategory of modules equivalent to R and all equivalences between them. This is a full subcategory of $\text{Pic}(R)$, namely, the component of the R -module R .

By Proposition 2.5 $(\text{Mod}_R)^\simeq$ inherits an \mathbb{E}_n -monoidal structure from Mod_R . Since the proposed sets of objects for both $\text{Pic}(R)$ and BGL_1R are closed under tensor products, by [Lur17, Proposition 2.2.1.1], both categories inherit the structure of \mathbb{E}_n -monoidal ∞ -groupoids, or equivalently, \mathbb{E}_n -spaces. By construction both are grouplike.

Definition 3.1. A *local system* of R -modules on a space X is simply a map $f: X \rightarrow \text{Pic}(R)$. The *Thom spectrum* of f is given by

$$M_R f = Mf := \text{colim}(X \rightarrow \text{Pic}(R) \rightarrow \text{Mod}_R).$$

This construction of Thom spectrum appears in [ABG⁺14a] where it is shown to agree with the definitions in [MS06] and [LMSM86]. We will often apply the above definition in the special case that the map f factors through BGL_1R .

3.1. The universal property and its consequences. Applying Theorem 2.8 to Mod_R we directly obtain Lewis’s theorem, or rather the generalization from the sphere spectrum to arbitrary R given in [ABG11]:

Corollary 3.2. *If X is an \mathbb{E}_n -space and $f: X \rightarrow \text{Pic}(R)$ is an \mathbb{E}_n -map, then Mf becomes an \mathbb{E}_n R -algebra.*

Definition 3.3. Given an \mathbb{E}_n R -algebra A , we define \mathbb{E}_n -spaces $\text{Pic}(R)_{/A}$ and $BGL_1R_{/A}$ by requiring the following squares to be pullbacks of \mathbb{E}_n -monoidal categories:

$$\begin{array}{ccccc} BGL_1R_{/A} & \longrightarrow & \text{Pic}(R)_{/A} & \longrightarrow & (\text{Mod}_R)_{/A} \\ \downarrow & & \downarrow & & \downarrow \\ BGL_1R & \longrightarrow & \text{Pic}(R) & \longrightarrow & \text{Mod}_R, \end{array}$$

where $(\text{Mod}_R)_{/A}$ is the slice of Mod_R over the underlying R -module of A .

Alternatively, we could define $BGL_1R_{/A}$ and $\text{Pic}(R)_{/A}$ merely as ∞ -groupoids by requiring the above diagram to consist of pullback squares of ∞ -categories. They would then inherit \mathbb{E}_n -monoidal structures from $(\text{Mod}_R)_{/A}$ in the same way that BGL_1R and $\text{Pic}(R)$ inherit their structure from Mod_R .

Warning 3.4. The above definition is an abuse of notation: $\text{Pic}(R)_{/A}$ and $BGL_1R_{/A}$ are *not* slice categories. We can think of the objects of $\text{Pic}(R)$ as invertible R -modules M equipped with a map $M \rightarrow A$ of R -modules, and of the morphisms as commuting triangles where the arrow $M \rightarrow M'$ is an equivalence.

We can now state a characterization of the \mathbb{E}_n -structure on Mf :

Theorem 3.5. *Let X be an \mathbb{E}_n -space and $f: X \rightarrow \text{Pic}(R)$ be an \mathbb{E}_n -map. The \mathbb{E}_n -algebra structure of Mf is characterized by the following universal property: the*

space of \mathbb{E}_n R -algebra maps $\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(Mf, A)$ is equivalent to the space of \mathbb{E}_n -lifts of f indicated below:

$$\begin{array}{ccc} & & \text{Pic}(R)_{/A} \\ & \nearrow & \downarrow \\ X & \xrightarrow{f} & \text{Pic}(R). \end{array}$$

Proof. Theorem 2.13 tells us directly that $\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(Mf, A)$ is the space of lifts of $X \xrightarrow{f} \text{Pic}(R) \rightarrow \text{Mod}_R$ to a lax \mathbb{E}_n -functor $X \rightarrow (\text{Mod}_R)_{/A}$. Now, edges in $(\text{Mod}_R)_{/A}^{\otimes}$ are coCartesian for the projection $(\text{Mod}_R)_{/A}^{\otimes} \rightarrow \mathbb{E}_n^{\otimes}$ if and only if their image in Mod_R^{\otimes} is coCartesian. So a lax \mathbb{E}_n -functor lifting f is automatically monoidal and factors through $\text{Pic}(R)_{/A}$. \square

Notation 3.6. If A is an \mathbb{E}_{n+1} -ring spectrum under R , that is an \mathbb{E}_{n+1} -ring spectrum equipped with an \mathbb{E}_{n+1} -morphism $\eta: R \rightarrow A$, we use Ind_R^A to denote the continuous \mathbb{E}_n -monoidal functor $\text{Mod}_R \rightarrow \text{Mod}_A$ induced by η (see [Lur17, Proposition 7.1.2.6]). Since this functor is \mathbb{E}_n -monoidal, there is also an induced functor $\text{Alg}_R^{\mathbb{E}_n} \rightarrow \text{Alg}_A^{\mathbb{E}_n}$ for which we also use the same notation.

Remark 3.7. Let us explain the relation between the notions of \mathbb{E}_{n+1} -ring spectrum under R and \mathbb{E}_n R -algebra. First, if A is an \mathbb{E}_{n+1} -ring spectrum under R , we can canonically equip it with the structure of an \mathbb{E}_n R -algebra: by [Lur17, Corollary 7.3.2.7], the right adjoint of $\text{Ind}_R^A: \text{Mod}_R \rightarrow \text{Mod}_A$ is lax \mathbb{E}_n -monoidal, so it preserves \mathbb{E}_n -algebra objects. In particular, this right adjoint lets us view the unit object A as an \mathbb{E}_n R -algebra.

Now, for $n = \infty$, the concepts of \mathbb{E}_∞ R -algebra and \mathbb{E}_∞ -ring spectrum under R actually coincide by [Lur17, Variant 7.1.3.8]. They do not for $n < \infty$, but an \mathbb{E}_n R -algebra structure on A still makes A an \mathbb{E}_n -ring spectrum under R [Lur17, Warning 7.1.3.9], so for such an A there is still an \mathbb{E}_{n-1} -monoidal functor Ind_R^A .

Remark 3.8. In case the \mathbb{E}_n R -algebra A is actually an \mathbb{E}_{n+1} -ring spectrum under R , we can simplify the description in Theorem 3.5: Ind_R^A restricts to an \mathbb{E}_n -monoidal functor $\text{Pic}(R) \rightarrow \text{Pic}(A)$, and giving a lift to $\text{Pic}(R)_{/A}$ is equivalent to giving a null homotopy of the composite $X \rightarrow \text{Pic}(R) \rightarrow \text{Pic}(A)$, because $\text{Pic}(R)_{/A}$ is the fiber of the functor $\text{Pic}(R) \rightarrow \text{Pic}(A)$ over the object $A \in \text{Pic}(A)$. Similar remarks apply to the BGL_1 case.

As a corollary, we easily obtain another of Lewis' results [LMSM86, IX.7.1]. To state it we will use the following notation.

Notation 3.9. For $m \leq n$, let $\text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n}$ be the free \mathbb{E}_n R -algebra on an \mathbb{E}_m R -algebra, i.e., the left adjoint to the forgetful functor $\text{Alg}_R^{\mathbb{E}_n} \rightarrow \text{Alg}_R^{\mathbb{E}_m}$.

Corollary 3.10. For any unital map of spaces $f: X \rightarrow BGL_1 R$, there is a natural equivalence

$$\text{Ind}_{\mathbb{E}_0}^{\mathbb{E}_n}(Mf) \xrightarrow{\sim} M\bar{f},$$

where $\bar{f}: \Omega^n \Sigma^n X \rightarrow BGL_1 R$ is the n -fold loop map adjoint to the pointed map $\Sigma^n f$.

Proof. We will check that both sides represent the same functor on $\text{Alg}_R^{\mathbb{E}_n}$; this yields the claim. Let A be an arbitrary \mathbb{E}_n - R -algebra. By adjunction, we have that $\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(\text{Ind}_{\mathbb{E}_0}^{\mathbb{E}_n}(Mf), A) \cong \text{Map}_{\text{Alg}_R^{\mathbb{E}_0}}(Mf, A)$ and by Theorem 3.5, this is the space L_0 of lifts of f to an \mathbb{E}_0 -map (i.e., a pointed map) $X \rightarrow BGL_1R/A$. Moreover, and again by Theorem 3.5, $\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(M\bar{f}, A)$ is the space L_n of lifts of \bar{f} to an \mathbb{E}_n -map $\Omega^n \Sigma^n X \rightarrow BGL_1R/A$.

Since $\Omega^n \Sigma^n X$ is the free grouplike \mathbb{E}_n -space on the pointed space X , the solid vertical maps (induced by composition with the unit $X \rightarrow \Omega^n \Sigma^n X$) in the following diagram of fiber sequences are equivalences:

$$\begin{array}{ccccc} L_n & \longrightarrow & \text{Map}_{\mathbb{E}_n}(\Omega^n \Sigma^n X, BGL_1R/A) & \longrightarrow & \text{Map}_{\mathbb{E}_n}(\Omega^n \Sigma^n X, BGLR) \\ \downarrow & & \downarrow \sim & & \downarrow \sim \\ L_0 & \longrightarrow & \text{Map}_{\mathbb{E}_0}(X, BGL_1R/A) & \longrightarrow & \text{Map}_{\mathbb{E}_0}(X, BGL_1R). \end{array}$$

It follows that the induced map $L_0 \rightarrow L_n$ is an equivalence, too. \square

Remark 3.11. This result also effortlessly extends to more general operads in place of \mathbb{E}_n , which in fact is the version proved by Lewis. We can replace \mathbb{E}_0 as well; for example, the same argument as used above also proves that if f is an m -fold loop map and \bar{f} denotes the universal extension to an n -fold loop for $n \geq m$, then $\text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n}(Mf) \xrightarrow{\sim} M\bar{f}$.

3.2. \mathbb{E}_n -orientations. The Thom spectrum MG for a topological group $G \rightarrow O$ arises classically in the theory of orientations of vector bundles, representing the universal cohomology theory that orients manifolds with structure group G . This point of view admits a generalization to \mathbb{E}_n -ring spectra, as we briefly summarize now; see [ABG⁺14a] and [ABG⁺14b] for a comparison of the various notions of orientations in the \mathbb{E}_1 and \mathbb{E}_∞ cases, as well as [ABG11].

For the remainder of this section let $n > 0$, R be an \mathbb{E}_{n+1} -ring spectrum and A be an \mathbb{E}_{n+1} -ring spectrum under R (see Notation 3.6).

Definition 3.12. Let $B(R, A)$ be the full subgroupoid of $\text{Pic}(R)_{/A}$ consisting of morphisms of R -modules $h: M \rightarrow A$ such that the adjoint $h^\dagger: \text{Ind}_R^A(M) \rightarrow A$ is an equivalence.

To study the relation between $B(R, A)$ and orientations, we need a lemma about these R -module morphisms:

Lemma 3.13. *Let $h_i: M_i \rightarrow A$ for $i = 1, 2$ be two morphisms of R -modules. Then $(h_1 \otimes_A h_2)^\dagger$ is an equivalence of A -modules if and only if both h_1^\dagger and h_2^\dagger are (here \otimes_A denotes the tensor product in $\text{Pic}(R)_{/A}$).*

Proof. Recall that in the monoidal structure of $\text{Pic}(R)_{/A}$, $h_1 \otimes_A h_2$ is given by the composite $M \otimes_R N \xrightarrow{h_1 \otimes_R h_2} A \otimes_R A \xrightarrow{\mu_A} A$, where μ_A is the multiplication on A . Since Ind_R^A is \mathbb{E}_n -monoidal, we see that $(h_1 \otimes_A h_2)^\dagger = h_1^\dagger \otimes_A h_2^\dagger$.

This makes it clear that if both h_i^\dagger are equivalences, then so is $(h_1 \otimes_A h_2)^\dagger$.

Now assume $(h_1 \otimes_A h_2)^\dagger$ is an equivalence with inverse g and let us prove that both h_i^\dagger are equivalences. To lighten the notation, set $M_i^\dagger := \text{Ind}_R^A(M_i)$. Notice

that by definition of g , the morphism of A -modules given by the composite

$$A \xrightarrow{g} M_1^\dagger \otimes_A M_2^\dagger \xrightarrow{\text{id}_{M_1^\dagger} \otimes_A h_2^\dagger} M_1^\dagger \otimes_A A$$

is a section of h_1^\dagger , which shows that M_1^\dagger splits as $A \oplus F_1$ where $F_1 := \text{fib}(h_1^\dagger)$. Under this splitting, h_1^\dagger corresponds to the projection $A \oplus F_1 \rightarrow A$. Of course there is an analogous splitting of M_2^\dagger as $A \oplus F_2$.

Then we get that $h_1^\dagger \otimes_A h_2^\dagger$ is the projection of $(A \oplus F_1) \otimes_A (A \oplus F_2) = A \oplus F_1 \oplus F_2 \oplus F_1 \otimes F_2$ onto the first summand. Since this map is assumed to be an equivalence, we conclude $F_1 = F_2 = 0$ and thus both h_1^\dagger and h_2^\dagger are equivalences. \square

As a corollary of (one direction) of the lemma, the set of objects in $B(R, A)$ is closed under tensor products in $\text{Pic}(R)_{/A}$, and thus $B(R, A)$ inherits an \mathbb{E}_n -monoidal structure.

We can now define orientations analogously to the definitions in [ABG⁺14a] and [ABG⁺14b] for the \mathbb{E}_1 and \mathbb{E}_∞ cases.

Definition 3.14. The *space of \mathbb{E}_n A -orientations* of an \mathbb{E}_n -map $f: X \rightarrow \text{Pic}(R)$ is the space of \mathbb{E}_n -lifts of f indicated below:

$$\begin{array}{ccc} & & B(R, A) \\ & \nearrow & \downarrow \\ X & \longrightarrow & \text{Pic}(R) \end{array}$$

Notice that because $B(R, A)$ is an \mathbb{E}_n -subspace of $\text{Pic}(R)_{/A}$, the universal property (Theorem 3.5) implies that an \mathbb{E}_n A -orientation of f determines a map of \mathbb{E}_n R -algebras $Mf \rightarrow A$. Of course, not every such map of algebras is an orientation; one way to state the requirement of factoring through $B(R, A)$ is to say that for every point $x: * \rightarrow X$, the adjoint θ_x^\dagger of the R -module map $\theta_x: M(f \circ x) \rightarrow Mf \rightarrow A$ is an equivalence. But if the \mathbb{E}_n -space X is grouplike, then this condition is automatically satisfied, as we show next.

Lemma 3.15. *If $f: X \rightarrow \text{Pic}(R)$ is an n -fold loop map, then the space of \mathbb{E}_n R -algebra morphisms $Mf \rightarrow A$ is equivalent to the space of \mathbb{E}_n A -orientations of f .*

Proof. We just need to show that the lifts of $f: X \rightarrow \text{Pic}(R)$ to $\text{Pic}(R)_{/A}$ as considered in Theorem 3.5 factor through $B(R, A)$ (we defined $B(R, A)$ as the union of some of the connected components of $\text{Pic}(R)_{/A}$).

Such a lift gives for each point $x \in X$ an R -linear map $\alpha_x: f(x) \rightarrow A$. If we have at least one multiplication on X and inverses for it, the map $\alpha_{x^{-1}}: f(x^{-1}) \rightarrow A$ satisfies that $\alpha_x \otimes \alpha_{x^{-1}} \simeq \alpha_{1_X}: f(1_X) \rightarrow A$ is homotopic to the unit $R \rightarrow A$ and thus is an equivalence when induced up to A . By Lemma 3.13, this implies α_x is in $B(R, A)$. \square

As expected \mathbb{E}_n A -orientations give rise to Thom isomorphisms of \mathbb{E}_n algebras.

Proposition 3.16. *An \mathbb{E}_n A -orientation of an \mathbb{E}_n -map $f: X \rightarrow \text{Pic}(R)$ give rise to a Thom isomorphism $\text{Ind}_R^A(Mf) \cong \text{Ind}_S^A(\Sigma_+^\infty X)$ of \mathbb{E}_n A -algebras (where S is the sphere spectrum). The equivalence of R -modules underlying this Thom isomorphism is a map $A \otimes_R Mf \rightarrow A \otimes \Sigma_+^\infty X$.*

Proof. We can also describe $B(R, A)$ as a pullback:

$$\begin{array}{ccc} B(R, A) & \longrightarrow & B(A, A) \\ \downarrow & & \downarrow \\ \text{Pic}(R) & \xrightarrow{\text{Ind}_R^A} & \text{Pic}(A) \end{array}$$

Notice that from the definition, $B(A, A)$ is the space of trivialized A -modules denoted by $A\text{-triv}$ in [ABG⁺14a]. Since it is the (entire) slice category $\text{Pic}(A)_{/A}$, it is contractible. Thus a lift of f to $B(R, A)$ gives a null-homotopy of $\text{Ind}_R^A \circ f$, and consequently, $\text{Ind}_R^A \circ f$ has the same Thom spectrum as the null map $X \rightarrow \text{Pic}(A)$.

Since $\text{Ind}_R^A: \text{Mod}_R \rightarrow \text{Mod}_A$ is a left adjoint, it preserves the colimit computing the Thom spectrum of f , that is, $M(\text{Ind}_R^A \circ f) \cong \text{Ind}_R^A(Mf)$. On the other hand the null map can be expressed as $\text{Ind}_R^S \circ c$ where $c: X \rightarrow \text{Pic}(S)$ is null. The same argument shows that $M(\text{Ind}_R^S \circ c) \cong \text{Ind}_R^S(\text{colim}_X S) = \text{Ind}_R^S(\Sigma_+^\infty X)$.

Finally, the underlying R -module spectrum of $\text{Ind}_R^A(M)$ is given by $Mf \otimes_R A$. \square

Corollary 3.17. *Let $n \geq 2$ and $f: X \rightarrow \text{Pic}(R)$ be an n -fold loop map with Thom spectrum Mf . Then f is canonically \mathbb{E}_{n-1} Mf -orientable.*

Proof. Recall that Mf is an \mathbb{E}_n R -algebra, which we can regard as an \mathbb{E}_n -ring spectrum under R (see Remark 3.7). The identity morphism $Mf \rightarrow Mf$ gives the required orientation. \square

The corresponding Thom isomorphism is an equivalence of \mathbb{E}_{n-1} R -algebras, $\text{Ind}_R^{Mf}(Mf) \cong Mf \otimes_R \text{Ind}_S^{Mf}(\Sigma_+^\infty X)$ whose underlying R -module equivalence is a morphism $Mf \otimes_R Mf \cong Mf \otimes \Sigma_+^\infty X$. This Thom isomorphism is due to Mahowald, see [Mah79, Theorem 1.2].

Now we give an alternative proof and mild generalization of the description of \mathbb{E}_n -orientations of ring spectra due to Chadwick and Mandell, see [CM15, Theorem 3.2]. While their argument uses the general Thom isomorphism, we deduce this result directly from the universal property of Thom ring spectra.

Corollary 3.18. *Let $f: X \rightarrow BGL_1 R$ be an \mathbb{E}_n -map and let A be a \mathbb{E}_{n+1} -ring spectrum under R . The space of \mathbb{E}_n R -algebra maps from Mf to A (regarded as an \mathbb{E}_n R -algebra) is either empty or equivalent to the space of \mathbb{E}_n R -algebra maps from $R \otimes \Sigma_+^\infty X$ to A , that is*

$$\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(Mf, A) \simeq \text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(R \otimes \Sigma_+^\infty X, A).$$

Proof. As explained in Remark 3.8, in this setting the space $\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(Mf, A)$ can be described as the space of null homotopies of the composite \mathbb{E}_n -map $X \xrightarrow{f} BGL_1 R \xrightarrow{\text{Ind}_R^A} BGL_1 A$. So we have the following pullback square of spaces:

$$\begin{array}{ccc} \text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(Mf, A) & \longrightarrow & * \\ \downarrow & & \downarrow \text{Ind}_R^A \circ f \\ * & \xrightarrow{c_A} & \text{Map}_{\mathbb{E}_n}(X, BGL_1 A) \end{array}$$

Here we have labeled the arrows from $*$ by the maps they pick out and c_A denotes the constant map with value the base point of $BGL_1 A$ which, when thought of as a functor, is the constant functor with value $A \in \text{Mod}_A$.

Now, the \mathbb{E}_n R -algebra $R \otimes \Sigma_+^\infty X$ is the Thom spectrum of c_R , so by the same reasoning as above we have the following pullback square:

$$\begin{array}{ccc} \text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(R \otimes \Sigma_+^\infty X, A) & \longrightarrow & * \\ \downarrow & & \downarrow \text{Ind}_R^A \circ c_R \\ * & \xrightarrow{c_A} & \text{Map}_{\mathbb{E}_n}(X, BGL_1 A) \end{array}$$

Since $\text{Ind}_R^A \circ c_R = c_A$, we see that in case $\text{Ind}_R^A \circ f \sim c_A$, the two pullbacks are equivalent, while if $\text{Ind}_R^A \circ f \not\sim c_A$, the first pullback is empty. \square

4. THOM SPECTRA AND VERSAL \mathbb{E}_n -ALGEBRAS

4.1. Characteristics of ring spectra. Let $0 \leq n \leq \infty$ be an integer or ∞ and consider an \mathbb{E}_{n+1} -ring spectrum R with associated category Mod_R of left R -module. As explained above, this category admits a monoidal product $\otimes = \otimes_R$ equipping it with the structure of an \mathbb{E}_n -monoidal category, so that there is a category $\text{Alg}_R^{\mathbb{E}_n}$ of \mathbb{E}_n -algebras over R .

We now extend the definition of characteristic introduced in [Szy14, Szy13] in two ways: Firstly, we allow arbitrary homotopy classes $S^k \rightarrow R$ in non-negative degrees and secondly we consider \mathbb{E}_n -algebras. For background and further results in the \mathbb{E}_∞ case we refer to Szymik's papers.

Definition 4.1. Given $k \geq 0$ and $\chi \in \pi_k R$ thought of as an R -linear map $\Sigma^k R \rightarrow R$, an algebra $A \in \text{Alg}_R^{\mathbb{E}_n}$ with unit $\eta: R \rightarrow A$ is said to be of *characteristic* χ if $\eta \circ \chi: \Sigma^k R \rightarrow A$ is null-homotopic.

Remark 4.2. If $n \geq 1$ and $A \in \text{Alg}_R^{\mathbb{E}_n}$ is of characteristic p , then the existence of the multiplication map $\mu: A \otimes_R A \rightarrow A$ shows $p \cdot \text{id}_A = 0$. However, note that for $n = 0$ an algebra being of characteristic p does not imply $p \cdot \text{id}_A$ is null, as the mod 2 Moore spectrum demonstrates.

Definition 4.3. For a given $\chi: \Sigma^k R \rightarrow R$, we define the *versal R -algebra* $R //_{\mathbb{E}_n} \chi$ of characteristic χ as the following pushout in $\text{Alg}_R^{\mathbb{E}_n}$:

$$\begin{array}{ccc} F_{\mathbb{E}_n} \Sigma^k R & \xrightarrow{\bar{0}} & R \\ \bar{\chi} \downarrow & & \downarrow \\ R & \longrightarrow & R //_{\mathbb{E}_n} \chi, \end{array}$$

where

- $F_{\mathbb{E}_n}: \text{Mod}_R \rightarrow \text{Alg}_R^{\mathbb{E}_n}$ is the free \mathbb{E}_n R -algebra functor, the left adjoint to the forgetful functor $\text{Alg}_R^{\mathbb{E}_n} \rightarrow \text{Mod}_R$.
- $\bar{g}: F_{\mathbb{E}_n} M \rightarrow N$ denotes the algebra map which is adjoint to the R -linear map $M \rightarrow N$ with $N \in \text{Alg}_R^{\mathbb{E}_n}$.

We use the term *versal* rather than *universal* as these algebras are not initial but only weakly initial in general, see [Szy14, Prop. 3.11]. When there is no risk of confusion, we shall write $R // \chi$ instead of $R //_{\mathbb{E}_n} \chi$.

The versal \mathbb{E}_n R -algebra $R // \chi$ admits a different characterization as the free \mathbb{E}_n R -algebra on the *pointed spectrum* $R \rightarrow R/\chi = \text{cof}(\Sigma^k R \xrightarrow{\chi} R)$, that is, $R //_{\mathbb{E}_n} \chi = \text{Ind}_{\mathbb{E}_0}^{\mathbb{E}_n}(R //_{\mathbb{E}_0} \chi)$. More generally we have:

Lemma 4.4. *Let $0 \leq m \leq n$, then the following \mathbb{E}_n R -algebras are equivalent:*

- (1) *The versal characteristic χ algebra $R //_{\mathbb{E}_n} \chi$,*
- (2) *the free \mathbb{E}_n R -algebra $\text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n}(R //_{\mathbb{E}_m} \chi)$ on the \mathbb{E}_m R -algebra $R //_{\mathbb{E}_m} \chi$,*
- (3) *the coequalizer $C_R(\chi) = \text{coeq}\left(F_{\mathbb{E}_n} \Sigma^k R \begin{array}{c} \xrightarrow{\bar{\chi}} \\ \xrightarrow{\bar{0}} \end{array} R\right)$ taken in $\text{Alg}_R^{\mathbb{E}_n}$.*

Proof. We first prove the equivalence of the algebras in (1) and (2). On the one hand, applying the left adjoint $\text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n}$ to the defining pushout diagram for $R //_{\mathbb{E}_m} \chi$ gives a pushout square

$$\begin{array}{ccc} \text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n} F_{\mathbb{E}_m} \Sigma^k R & \xrightarrow{\bar{0}} & \text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n} R \\ \text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n} \bar{\chi} \downarrow & & \downarrow \\ \text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n} R & \longrightarrow & \text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n}(R //_{\mathbb{E}_m} \chi). \end{array}$$

On the other hand, since $\text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n}$ preserves the initial R -algebra R and $\text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n} F_{\mathbb{E}_m} \simeq F_{\mathbb{E}_n}$, this diagram is naturally equivalent to the pushout square

$$\begin{array}{ccc} F_{\mathbb{E}_n} \Sigma^k R & \xrightarrow{\bar{0}} & R \\ \bar{\chi} \downarrow & & \downarrow \\ R & \longrightarrow & R //_{\mathbb{E}_n} \chi, \end{array}$$

hence it follows immediately that $R //_{\mathbb{E}_n} \chi$ is equivalent to $\text{Ind}_{\mathbb{E}_m}^{\mathbb{E}_n}(R //_{\mathbb{E}_m} \chi)$.

To prove that $R //_{\mathbb{E}_n} \chi$ is equivalent to the coequalizer $C_R(\chi)$, we observe that both algebras corepresent the same functor on $\text{Alg}_R^{\mathbb{E}_n}$, which maps a given algebra A with unit map η to the space of null-homotopies of the composite $\Sigma^k R \xrightarrow{\chi} R \xrightarrow{\eta} A$. \square

There is a Thom-isomorphism type lemma for algebras of a given characteristic, which is a straightforward generalization of [Szy14, Prop. 3.2].

Lemma 4.5. *Let $n \geq 1$ and suppose $\eta: R \rightarrow A$ is a map of \mathbb{E}_{n+1} -ring spectra where A has characteristic χ , then there is a natural equivalence of \mathbb{E}_n A -algebras*

$$\text{Ind}_R^A(F_{\mathbb{E}_n}(\Sigma^{k+1} R)) \xrightarrow{\sim} \text{Ind}_R^A(R //_{\mathbb{E}_n} \chi).$$

Proof. The two algebras in question are constructed as pushouts in $\text{Alg}_R^{\mathbb{E}_n}$

$$\begin{array}{ccc} F_{\mathbb{E}_n} \Sigma^k R & \xrightarrow{\bar{0}} & R \\ \bar{0} \downarrow & & \downarrow \\ R & \longrightarrow & F_{\mathbb{E}_n}(\Sigma^{k+1} R) \end{array} \qquad \begin{array}{ccc} F_{\mathbb{E}_n} \Sigma^k R & \xrightarrow{\bar{0}} & R \\ \bar{\chi} \downarrow & & \downarrow \\ R & \longrightarrow & R // \chi. \end{array}$$

Since $F_{\mathbb{E}_n}(\Sigma^k R) \otimes_R A \xrightarrow{\chi \otimes \text{id}_A} A$ is homotopic to 0 if A has characteristic χ , the two diagrams become equivalent after applying the functor $\text{Ind}_R^A: \text{Alg}_R^{\mathbb{E}_n} \rightarrow \text{Alg}_A^{\mathbb{E}_n}$ (see Notation 3.6), which is given on the level of R -modules by smashing with A . \square

Lemma 4.6. *Suppose $A \in \text{Alg}_R^{\mathbb{E}_n}$. If A is of characteristic $\chi: \Sigma^k R \rightarrow R$, then*

$$\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(R //_{\mathbb{E}_n} \chi, A) = \Omega^{\infty+k+1} A;$$

otherwise the mapping space is empty.

Proof. The proof of [Szy14, Cor. 2.9] generalizes effortlessly: By definition of $R // \chi$ as an \mathbb{E}_n -algebra, there exists a pullback square

$$\begin{array}{ccc} \text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(R // \chi, A) & \longrightarrow & \text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(R, A) = * \\ \downarrow & & \downarrow \bar{0}^* \\ * = \text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(R, A) & \xrightarrow{\chi^*} & \text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(F_{\mathbb{E}_n} \Sigma^k R, A) = \Omega^{\infty+k} A \end{array}$$

showing that $\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(R // \chi, A) \xrightarrow{\sim} \Omega^{\infty+k+1} A$ if A is of characteristic χ so that the two points lie in the same component of $\Omega^\infty A$, and empty otherwise. \square

In other words, specializing to the case in which χ is the multiplication by p map we see that $\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(R // p, A)$ is the space of null-homotopies of the composite map $R \xrightarrow{p} R \xrightarrow{\eta} A$ in R -modules, which is equivalent to the space of homotopies $\eta \sim (1-p)\eta$ under the map $\text{Hom}_R(R, A) \rightarrow \text{Hom}_R(R, A)$ given by $\alpha \mapsto \alpha + \eta$. This in turn admits an interpretation in terms of Thom spectra, given in the next section.

4.2. Thom spectra as versal characteristic χ algebras. The goal of this section is to identify the Thom spectrum classified by a map $f: S^{k+1} \rightarrow BGL_1 R$ with the versal characteristic χ algebra, where $\chi = \chi(f): \Sigma^k R \rightarrow R$ is the characteristic corresponding to f as defined below.

Lemma 4.7. *If $k \geq 0$ and A is any unital R -module, that is, an R -module equipped with an R -linear map $\eta: R \rightarrow A$, then there is a natural pullback square of spaces*

$$\begin{array}{ccc} \text{Map}_*(S^{k+1}, BGL_1 R/A) & \longrightarrow & * \\ \downarrow & & \downarrow \eta \\ \text{Map}_*(S^{k+1}, BGL_1 R) & \longrightarrow & \text{Map}_*(S^k, \Omega^\infty A) = \Omega^{\infty+k} A \end{array}$$

where the right vertical map picks out the constant map with value the unit $\eta \in \text{Hom}_R(R, A) = \Omega^\infty A$.

Proof. If we regard $GL_1 R$ as the space of R -linear self-equivalences of R , it acts on $\text{Hom}_R(R, A)$ by pre-composition. This induces a map $\eta_*: GL_1 R \rightarrow \text{Hom}_R(R, A)$ defined by $g \mapsto \eta \circ g$, where $\eta: R \rightarrow A$ is the unit of A . Let $GL_1 R/A$ be the fiber of this map at the point $\eta \in \text{Hom}_R(R, A)$, so that $GL_1 R/A$ can be thought of as the space of automorphisms g of R compatible with η — in the sense that $\eta \circ g$ is homotopic to η . This $GL_1 R/A$ can alternatively be described as the space of endomorphisms of the object η in the ∞ -groupoid $BGL_1 R/A$. In other words, $\Omega_\eta(BGL_1 R/A) = GL_1 R/A$.

This shows that $\text{Map}_*(S^k, GL_1R) = \text{Map}_*(S^{k+1}, BGL_1R)$ and similarly for GL_1R/A . Therefore, applying $\text{Map}_*(S^k, -)$ to the pullback square

$$\begin{array}{ccc} GL_1R/A & \longrightarrow & * \\ \downarrow & & \downarrow \eta \\ GL_1R & \xrightarrow{\eta_*} & \text{Hom}_R(R, A) \end{array}$$

gives the desired result. \square

Definition 4.8. Suppose given a map $f: S^{k+1} \rightarrow BGL_1R$ and let $\tilde{f}: \Sigma^k R \rightarrow R$ be the associated homotopy class. The *characteristic* $\chi(f): \Sigma^k R \rightarrow R$ associated with f is then defined by

$$\chi(f) = \begin{cases} \tilde{f} - 1 & \text{if } k = 0 \\ \tilde{f} & \text{if } k > 0. \end{cases}$$

Proposition 4.9. *If $f: S^{k+1} \rightarrow BGL_1R$ is a based map and $\bar{f}: \Omega^n \Sigma^n S^{k+1} \rightarrow BGL_1R$ is the corresponding n -fold loop map, then for any $A \in \text{Alg}_R^{\mathbb{E}_n}$, there is an equivalence of spaces*

$$\text{Map}_{\text{Alg}_R^{\mathbb{E}_n}}(M\bar{f}, A) = \Omega^{\infty+k+1}A$$

if A has characteristic $\chi(f)$; otherwise, the mapping space is empty.

Proof. By Corollary 3.10, it suffices to prove that $\text{Alg}_R^{\mathbb{E}_0}(Mf, A) = \Omega^{\infty+k+1}A$. To this end, let A be a unital R -module and consider the following commutative diagram of spaces

$$\begin{array}{ccccc} \text{Alg}_R^{\mathbb{E}_0}(Mf, A) & \longrightarrow & \text{Map}_*(S^{k+1}, BGL_1R/A) & \longrightarrow & * \\ \downarrow & & \downarrow & & \downarrow \underline{\eta} \\ * & \xrightarrow{f} & \text{Map}_*(S^{k+1}, BGL_1R) & \longrightarrow & \text{Map}_*(S^k, A) = \Omega^{\infty+k}A. \\ & & \searrow \eta_* \circ f & & \end{array}$$

The left square is a pullback by the universal property of Thom spectra (Theorem 3.5), while the right square is one due to Lemma 4.7. Hence the outer rectangle is a pullback square.

The right vertical arrow $\underline{\eta}$ is induced by $* \rightarrow \Omega^\infty A$ corresponding to the unit η of A , so

$$[\underline{\eta}] = \begin{cases} \eta \in \pi_0(\Omega^\infty A) & \text{if } k = 0 \\ 0 \in \pi_k(\Omega^\infty A) & \text{if } k > 0. \end{cases}$$

This explains the distinction of the two cases appearing in Definition 4.8: the pullback is non-empty if $k = 0$ and $\eta_* f \simeq \eta$, or if $k > 0$ and $\eta_* f \simeq 0$. \square

We thus obtain:

Theorem 4.10. *For any $n \geq 0$ and $f: S^{k+1} \rightarrow BGL_1R$ with corresponding n -fold loop map $\bar{f}: \Omega^n \Sigma^n S^{k+1} \rightarrow BGL_1R$ and with associated characteristic $\chi = \chi(f)$, there is an equivalence $M\bar{f} = R //_{\mathbb{E}_n} \chi$ of \mathbb{E}_n R -algebras.*

Proof. In view of Proposition 4.9 and Lemma 4.6, $M\bar{f}_{\mathbb{E}_n}$ and $R // \chi$ corepresent the same functor on $\text{Alg}_R^{\mathbb{E}_n}$, hence they are equivalent as \mathbb{E}_n R -algebras. \square

Remark 4.11. An alternative proof of Theorem 4.10 starts with Corollary 3.10 to first reduce the claim to the \mathbb{E}_0 case, and then identifies the Thom spectrum with the versal characteristic χ \mathbb{E}_0 -algebra. The latter statement admits a direct and easy computational proof, as can be found for example in [MRS01, Lemma 3.3] for the case $f = (p - 1)$.

5. APPLICATIONS

5.1. The Hopkins–Mahowald theorem I: $H\mathbb{F}_p$. In this section we will assume that all spaces and spectra are implicitly p -complete for some prime p . Let $f_p: S^1 \rightarrow BGL_1(S^0)$ be the map corresponding to the element $1 - p \in \mathbb{Z}_p^\times \cong \pi_1 BGL_1(S^0)$ and \bar{f}_p the extension of f_p to a double loop map, making the following diagram commute:

$$\begin{array}{ccc} S^1 & \xrightarrow{f_p} & BGL_1 S^0 \\ \downarrow & \nearrow \bar{f}_p & \\ \Omega^2 \Sigma^2 S^1 & & \end{array}$$

Theorem 5.1. *If $f_p: S^1 \rightarrow BGL_1(S^0)$ is the map corresponding to the element $1 - p \in \mathbb{Z}_p^\times \cong \pi_1 BGL_1(S^0)$, then the following three spectra are equivalent as \mathbb{E}_2 -algebras:*

- (1) *The Thom spectrum $M\bar{f}_p$,*
- (2) *the versal characteristic p \mathbb{E}_2 -algebra $S^0 //_{\mathbb{E}_2} p$, and*
- (3) *the Eilenberg–Mac Lane spectrum $H\mathbb{F}_p$ viewed as an \mathbb{E}_2 -algebra.*

Proof. The equivalence of $M\bar{f}_p$ and $S^0 // p$ as \mathbb{E}_2 -algebras is a special case of Theorem 4.10. In order to prove that these algebras are also equivalent to $H\mathbb{F}_p$ viewed as an \mathbb{E}_2 -algebra, we observe that there exists a (canonical) map of connective \mathbb{E}_2 -algebras

$$\phi: S^0 // p \longrightarrow H\mathbb{F}_p$$

witnessing the homotopy $p \sim 0$, because $H\mathbb{F}_p$ is of characteristic p and $\pi_1 H\mathbb{F}_p = 0$. Taking $A = H\mathbb{F}_p$ in Lemma 4.5 and using the computation [CLM76] of the homology of free \mathbb{E}_2 -algebras shows that

$$\pi_*(S^0 // p \otimes H\mathbb{F}_p) \cong \pi_*(F_{\mathbb{E}_2} S^1 \otimes H\mathbb{F}_p) \cong H_*(F_{\mathbb{E}_2} S^1, \mathbb{F}_p) \cong \mathcal{A}_p,$$

the dual of the mod p Steenrod algebra. Moreover, the $H\mathbb{F}_p$ \mathbb{E}_2 -algebra morphism

$$F_{\mathbb{E}_2} S^1 \otimes H\mathbb{F}_p \xrightarrow{\sim} S^0 // p \otimes H\mathbb{F}_p \xrightarrow{\phi \otimes H\mathbb{F}_p} H\mathbb{F}_p \otimes H\mathbb{F}_p$$

is adjoint to $S^1 \rightarrow H\mathbb{F}_p \otimes H\mathbb{F}_p$. Unwinding the construction, we see that this map picks out the fundamental class, so the equivalence of (2) and (3) follows. \square

Remark 5.2. The equivalence between the versal \mathbb{E}_2 -algebra of characteristic p and $H\mathbb{F}_p$ has been proven independently in the second version of [MNN15].

Remark 5.3. The reader might wonder why the above argument does not apply to the unique (up to contractible choice) map $\psi: F_{\mathbb{E}_2} S^1 \rightarrow H\mathbb{F}_p$ to imply that $F_{\mathbb{E}_2} S^1 \simeq S^0 // p$. This conclusion is false since the two algebras corepresent different functors: $F_{\mathbb{E}_2} S^1$ corepresents $\Omega^{\infty+1}$, while the functor corepresented by $S^0 // p$ only agrees with that on \mathbb{E}_n -algebras of characteristic p by Lemma 4.6.

The reason the argument does not apply is that the algebra map $\psi \otimes \mathbb{F}_p: F_{\mathbb{E}_2} S^1 \otimes H\mathbb{F}_p \rightarrow H\mathbb{F}_p \otimes H\mathbb{F}_p$ does not correspond to a non-zero multiple of the fundamental class, so it is not an equivalence. This also shows that, in general, the equivalence in Lemma 4.5 does not arise from a morphism of algebras $F_{\mathbb{E}_n}(\Sigma^{k+1}R) \rightarrow R // \chi$.

The Hopkins–Mahowald theorem has the following well-known application. Recall that the Morava K -theory spectrum $K(n)$ of height n is an \mathbb{E}_1 -ring spectrum with coefficients $K(n)_* = \mathbb{F}_{p^n}[v_n^{\pm 1}]$, where the degree of v_n is $2p^n - 2$.

Corollary 5.4. *There is no \mathbb{E}_2 -refinement of the \mathbb{E}_1 -ring structure on $K(n)$.*

Proof. Assume that $K(n)$ has the structure of an \mathbb{E}_2 -ring spectrum. Since p acts trivially on the coefficients, the universal property of $H\mathbb{F}_p$ induces a map of \mathbb{E}_2 -ring spectra

$$H\mathbb{F}_p \longrightarrow K(n).$$

However, since $H\mathbb{F}_p$ is $K(n)$ -acyclic for all finite n by [Rav84, Theorem 2.1(i)], this map must be null, which yields a contradiction. \square

5.2. The Hopkins–Mahowald theorem II: $H\mathbb{Z}$. The goal of this section is to deduce an integral analogue of Theorem 5.1, using the framework of this paper. Our approach is different from the arguments given in [Mah79], [CMT81], and [Blu10], as it does not rely on any further explicit homology computations. Unless otherwise stated, we will implicitly work in the p -complete category of spaces and spectra.

We start with a folklore result of independent interest, which shows that the \mathbb{E}_n -ring structure on Eilenberg–MacLane ring spectra are essentially unique, for all $0 \leq n \leq \infty$. Since we do not know a published reference for this result, we sketch an argument we learned from Tyler Lawson.

Lemma 5.5. *If R is a commutative ring, then the space of \mathbb{E}_∞ -structures on the Eilenberg–MacLane spectrum HR that induce the given ring structure on $\pi_* HR = R$ is contractible.*

Proof. This follows from the observation that the endomorphism operad of HR in spectra is discrete; indeed, for all $m \geq 0$:

$$\begin{aligned} \text{Map}((HR)^{\otimes m}, HR) &\simeq \text{Map}(\tau_{\leq 0}(HR)^{\otimes m}, HR) \\ &\simeq \text{Map}(H(R^{\otimes m}), HR) \\ &\simeq \text{Hom}(R^{\otimes m}, R). \end{aligned}$$

In order to construct a coherent multiplication on HR , we have to pick out the union of path components corresponding to the iterated multiplication maps. This is a suboperad which is levelwise contractible, and hence we can replace it by an \mathbb{E}_∞ -operad. \square

To construct $H\mathbb{Z}$ as a Thom spectrum, we will combine the construction of $H\mathbb{F}_p$ as a Thom spectrum given in Theorem 5.1 with the idea of intermediate Thom spectra from [Bea17]. Consider the fiber sequence

$$S^3\langle 3 \rangle \longrightarrow S^3 \longrightarrow K(\mathbb{Z}, 3)$$

which realizes the bottom of the Whitehead tower for S^3 . Looping twice gives another fiber sequence

$$(6) \quad \Omega^2(S^3\langle 3 \rangle) \longrightarrow \Omega^2 S^3 \xrightarrow{\pi} S^1 \simeq \Omega^2 K(\mathbb{Z}, 3).$$

As in the previous section, let $f_p: \Omega^2 S^3 \rightarrow BGL_1(S_p^0)$ be the free \mathbb{E}_2 -map classifying the element $(1-p) \in \pi_0 S_p^0$ and let g_p be the composite

$$g_p: \Omega^2(S^3\langle 3 \rangle) \longrightarrow \Omega^2 S^3 \xrightarrow{f_p} BGL_1 S_p^0.$$

For clarity, we will indicate the composite of a map $h: X \rightarrow BGL_1 R$ with the inclusion $BGL_1 R \rightarrow \text{Mod}_R$ by the corresponding capital letter H ; in particular, Mh is the colimit of H .

Theorem 5.7. *There is an equivalence $Mg_p \simeq H\mathbb{Z}_p$ of \mathbb{E}_2 -ring spectra.*

Proof. Following ideas of Beardsley [Bea17], we will first construct an intermediate Thom spectrum $M\phi_p$ associated to a map $\phi_p: S^1 \rightarrow BGL_1(Mg_p)$ from the base of the fiber sequence (6). The properties of this spectrum allow us then to determine the homotopy groups of Mg_p , from which the claim will follow.

Let Φ_p be the operadic Kan extension of F_p along π with respect to the \mathbb{E}_2 -operad. By [Bea17], Φ_p factors canonically through a map $\phi_p: S^1 \rightarrow BGL_1(Mg_p)$, so we obtain the following (non-commutative) diagram

$$\begin{array}{ccccc} & & g_p & & \\ & \curvearrowright & & \curvearrowleft & \\ \Omega^2(S^3\langle 3 \rangle) & \longrightarrow & \Omega^2 S^3 & \xrightarrow{f_p} & BGL_1 S_p^0 \longrightarrow \text{Sp} \\ & & \downarrow \pi & & \nearrow \\ & & S^1 & \xrightarrow[\phi_p]{} & BGL_1 Mg_p. \end{array}$$

We will now identify the Thom spectrum of ϕ_p in two different ways. On the one hand, by [Bea17], there is an equivalence $M\phi_p \simeq Mf_p$ of \mathbb{E}_1 -ring spectra, which in turn is equivalent to $H\mathbb{F}_p$ by Theorem 5.1. On the other hand, Theorem 4.10 shows that $M\phi_p$ is also the versal \mathbb{E}_0 - Mg_p -algebra of characteristic χ_p , where $\chi_p \in \pi_0(Mg_p)^\times$ is the element corresponding to ϕ_p . Combining these two descriptions, we obtain a fiber sequence

$$(8) \quad Mg_p \xrightarrow{1-\chi_p} Mg_p \longrightarrow H\mathbb{F}_p$$

of spectra. By construction, Mg_p is a connected p -complete \mathbb{E}_2 -ring spectrum.

We now claim that Mg_p is of finite type. To this end, first observe that it follows from Lemma 3.15 that Mg_p is $Mf_p \simeq H\mathbb{F}_p$ -oriented. Therefore, we get that

$$H_*(Mg_p; \mathbb{F}_p) \cong H_*(\Omega^2(S^3\langle 3 \rangle); \mathbb{F}_p).$$

In particular, the homology of Mg_p is finitely generated in each degree, so an argument with Serre classes shows that $\pi_*((Mg_p)_p) \cong \pi_*(Mg_p)$ is also finitely generated over \mathbb{Z}_p in each degree. Furthermore, the Hurewicz theorem implies that

$$\mathbb{F}_p \otimes \pi_0 Mg_p \cong H_0(\Omega^2(S^3\langle 3 \rangle); \mathbb{F}_p) \cong \mathbb{F}_p,$$

because $\Omega^2(S^3\langle 3 \rangle)$ is connected. In particular, $\pi_0 Mg_p$ is either isomorphic to \mathbb{Z}_p or \mathbb{F}_p as it is finitely generated over \mathbb{Z}_p .

Next, consider the long exact sequence of homotopy groups associated to (8), which degenerates to a short exact sequence

$$0 \longrightarrow \pi_0 Mg_p \xrightarrow{1-\chi_p} \pi_0 Mg_p \longrightarrow \mathbb{F}_p \longrightarrow 0$$

as well as isomorphisms $1 - \chi_p: \pi_i M g_p \xrightarrow{\sim} \pi_i M g_p$ for $i \geq 1$. Using that $M g_p$ is of finite type once more, one easily sees that this forces $1 - \chi_p = p$ and

$$\pi_* M g_p \cong \begin{cases} \mathbb{Z}_p & i = 0 \\ 0 & i \neq 1, \end{cases}$$

hence $M g_p \simeq H\mathbb{Z}_p$ as spectra. By virtue of Lemma 5.5, we conclude that this equivalence is also one of \mathbb{E}_2 -ring spectra. \square

As in [Blu10, Section 9.3], we may glue these maps together to construct $H\mathbb{Z}$ as a Thom spectrum as well. Here, we work in the category of all spectra, not just p -complete ones.

Corollary 5.9. *There is a map $g: \Omega^2 S^3 \rightarrow BGL_1 S^0$ whose associated Thom spectrum is equivalent to $H\mathbb{Z}$ as \mathbb{E}_2 -ring spectra.*

5.3. Topological Hochschild homology and the cotangent complex. Theorem 4.10 allows to transport results proven for Thom spectra to versal algebras and vice versa. We illustrate this idea with two examples, computing the topological Hochschild homology and the cotangent complex of these algebras. These results generalize previous computations of Szymik [Szy14]. In this section, R is an \mathbb{E}_∞ -ring spectrum.

Proposition 5.10. *Let $R //_{\mathbb{E}_n} \chi$ be the versal \mathbb{E}_n -algebra of characteristic χ corresponding to a map $\bar{f}: \Omega^n \Sigma^n S^{k+1} \rightarrow BGL_1 R$. If either $n \geq 3$ or $n \geq 2$ and the versal \mathbb{E}_n -algebra $R //_{\mathbb{E}_n} \chi$ of characteristic χ admits an \mathbb{E}_∞ -refinement, then there is an equivalence*

$$B_+ \Omega^n \Sigma^n S^{k+1} \otimes R //_{\mathbb{E}_n} \chi \xrightarrow{\sim} \mathrm{THH}(R //_{\mathbb{E}_n} \chi).$$

Proof. Using Theorem 4.10, it is enough to compute the topological Hochschild homology of $M\bar{f}$. If $n \geq 3$, the claim thus follows from [BCS10, Thm. 3], while the $n \geq 2$ case is covered by [Blu10, 1.6]. \square

The deformation theory of an \mathbb{E}_n R -algebra A is captured by its \mathbb{E}_n -cotangent complex $L_A^{(n)}$, considered as an object in the \mathbb{E}_n -monoidal category $\mathrm{Mod}_A^{\mathbb{E}_n} = \mathrm{Sp}((\mathrm{Alg}_R^{\mathbb{E}_n})/A)$ of \mathbb{E}_n A -modules, see [Fra13] or [Lur17].

Proposition 5.11. *If $n \geq 1$ and $R //_{\mathbb{E}_n} \chi$ a the versal \mathbb{E}_n -algebra of characteristic $\chi: \Sigma^k R \rightarrow R$, then there is an equivalence*

$$\Sigma^{k+1} F_{\mathbb{E}_1}(\Sigma^{n-1} R) \otimes R //_{\mathbb{E}_n} \chi \xrightarrow{\sim} L_{R //_{\mathbb{E}_n} \chi}^{(n)}$$

of \mathbb{E}_n $R //_{\mathbb{E}_n} \chi$ -modules.

Proof. Firstly, the cotangent complex of the free \mathbb{E}_n -algebra $F_{\mathbb{E}_n} \Sigma^k R$ can be computed as an \mathbb{E}_n $F_{\mathbb{E}_n} \Sigma^k R$ -module using [Fra13, 2.12, 2.25] as follows:

$$L_{F_{\mathbb{E}_n} \Sigma^k R}^{(n)} \simeq \Sigma^k \int_{S^{n-1}} F_{\mathbb{E}_n} \Sigma^k R \simeq \Sigma^k F_{\mathbb{E}_1}(\Sigma^{n-1} R) \otimes F_{\mathbb{E}_n} R,$$

where \int denotes factorization homology. Secondly, the natural cofiber sequence for computing the relative cotangent complex [Fra13, 2.11] specializes to give a cofiber sequence

$$L_{F_{\mathbb{E}_n} R}^{(n)} \otimes_{F_{\mathbb{E}_n} R} R \longrightarrow L_R^{(n)} \longrightarrow L_{R/F_{\mathbb{E}_n} R}^{(n)}.$$

Since $L_R^{(n)}$ is contractible, we get

$$\begin{aligned} L_{R/F_{\mathbb{E}_n}R}^{(n)} &\simeq \Sigma L_{F_{\mathbb{E}_n}R}^{(n)} \otimes_{F_{\mathbb{E}_n}R} R \\ &\simeq \Sigma^{k+1} F_{\mathbb{E}_1}(\Sigma^{n-1}R) \otimes_{F_{\mathbb{E}_n}R} R \\ &\simeq \Sigma^{k+1} F_{\mathbb{E}_1}(\Sigma^{n-1}R). \end{aligned}$$

The base-change formula [Lur17, 7.3.3.7] applied to the defining pushout diagram

$$\begin{array}{ccc} F_{\mathbb{E}_n}R & \xrightarrow{\bar{0}} & R \\ \bar{\chi} \downarrow & & \downarrow \psi \\ R & \longrightarrow & R // \chi \end{array}$$

now gives the desired equivalence

$$L_{R//\chi}^{(n)} \xrightarrow{\sim} \psi! L_{R/F_{\mathbb{E}_n}R}^{(n)} \simeq \Sigma^{k+1} F_{\mathbb{E}_1}(\Sigma^{n-1}R) \otimes R // \chi$$

of $\mathbb{E}_n R // \chi$ -modules. \square

By Theorem 4.10, we can translate this result immediately into a statement about the \mathbb{E}_n -cotangent complex of certain Thom spectra.

Corollary 5.12. *Let $n \geq 1$ and $f: S^{k+1} \rightarrow BGL_1(R)$ with corresponding n -fold loop map $\bar{f}: \Omega^n \Sigma^n S^{k+1} \rightarrow BGL_1(R)$, then there is an equivalence*

$$\Sigma^{k+1} F_{\mathbb{E}_1}(S^{n-1}) \otimes M\bar{f} \xrightarrow{\sim} L_{M\bar{f}}^{(n)}$$

of $\mathbb{E}_n M\bar{f}$ -modules.

REFERENCES

- [ABG11] Matthew Ando, Andrew J. Blumberg, and David Gepner. Parametrized spectra, multiplicative Thom spectra, and the twisted Umkehr map. <http://arxiv.org/abs/1112.2203>, 2011.
- [ABG⁺14a] Matthew Ando, Andrew J. Blumberg, David Gepner, Michael J. Hopkins, and Charles Rezk. An ∞ -categorical approach to R -line bundles, R -module Thom spectra, and twisted R -homology. *J. Topol.*, 7(3):869–893, 2014.
- [ABG⁺14b] Matthew Ando, Andrew J. Blumberg, David Gepner, Michael J. Hopkins, and Charles Rezk. Units of ring spectra, orientations and Thom spectra via rigid infinite loop space theory. *J. Topol.*, 7(4):1077–1117, 2014.
- [Bak] Andrew Baker. Characteristics for E_∞ ring spectra. <http://arxiv.org/abs/1405.3695>, final version to appear in the Proceedings of the Alpine Algebraic and Applied Topology Conference.
- [BCS10] Andrew J. Blumberg, Ralph L. Cohen, and Christian Schlichtkrull. Topological Hochschild homology of Thom spectra and the free loop space. *Geom. Topol.*, 14(2):1165–1242, 2010.
- [Bea17] Jonathan Beardsley. Relative Thom spectra via operadic Kan extensions. *Algebr. Geom. Topol.*, 17(2):1151–1162, 2017. Errata available at <http://msp.org/agt/2017/17-2/p20.xhtml>.
- [Blu10] Andrew J. Blumberg. Topological Hochschild homology of Thom spectra which are E_∞ ring spectra. *J. Topol.*, 3(3):535–560, 2010.
- [CLM76] Frederick R. Cohen, Thomas J. Lada, and J. Peter May. *The homology of iterated loop spaces*. Lecture Notes in Mathematics, Vol. 533. Springer-Verlag, Berlin-New York, 1976.
- [CM15] Steven Greg Chadwick and Michael A. Mandell. E_n genera. *Geom. Topol.*, 19(6):3193–3232, 2015.

- [CMT81] F. R. Cohen, J. P. May, and L. R. Taylor. $K(\mathbf{Z}, 0)$ and $K(\mathbf{Z}_2, 0)$ as Thom spectra. *Illinois J. Math.*, 25(1):99–106, 1981.
- [DHS88] Ethan S. Devinatz, Michael J. Hopkins, and Jeffrey H. Smith. Nilpotence and stable homotopy theory. I. *Ann. of Math. (2)*, 128(2):207–241, 1988.
- [Fra13] John Francis. The tangent complex and Hochschild cohomology of \mathcal{E}_n -rings. *Compos. Math.*, 149(3):430–480, 2013.
- [LMSM86] L. G. Lewis, Jr., J. P. May, M. Steinberger, and J. E. McClure. *Equivariant stable homotopy theory*, volume 1213 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986. With contributions by J. E. McClure.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009. Also available from author’s website as <http://www.math.harvard.edu/~lurie/papers/croppedtopoi.pdf>.
- [Lur17] Jacob Lurie. *Higher Algebra*. 2017. Draft available from author’s website as <http://www.math.harvard.edu/~lurie/papers/HigherAlgebra.pdf>.
- [Mah77] Mark Mahowald. A new infinite family in $2\pi_*^s$. *Topology*, 16(3):249–256, 1977.
- [Mah79] Mark Mahowald. Ring spectra which are Thom complexes. *Duke Math. J.*, 46(3):549–559, 1979.
- [MNN15] Akhil Mathew, Niko Naumann, and Justin Noel. On a nilpotence conjecture of J. P. May. *J. Topol.*, 8(4):917–932, 2015.
- [MRS01] Mark Mahowald, Douglas C. Ravenel, and Paul Shick. The Thomified Eilenberg-Moore spectral sequence. In *Cohomological methods in homotopy theory (Bellaterra, 1998)*, volume 196 of *Progr. Math.*, pages 249–262. Birkhäuser, Basel, 2001.
- [MS06] J. P. May and J. Sigurdsson. *Parametrized homotopy theory*, volume 132 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2006.
- [MS16] Akhil Mathew and Vesna Stojanoska. The Picard group of topological modular forms via descent theory. *Geom. Topol.*, 20(6):3133–3217, 2016.
- [Pri78] Stewart Priddy. $K(\mathbf{Z}/2)$ as a Thom spectrum. *Proc. Amer. Math. Soc.*, 70(2):207–208, 1978.
- [Rav84] Douglas C. Ravenel. Localization with respect to certain periodic homology theories. *Amer. J. Math.*, 106(2):351–414, 1984.
- [Sul05] Dennis P. Sullivan. *Geometric topology: localization, periodicity and Galois symmetry*, volume 8 of *K-Monographs in Mathematics*. Springer, Dordrecht, 2005. The 1970 MIT notes, Edited and with a preface by Andrew Ranicki.
- [Szy13] Markus Szymik. String bordism and chromatic characteristics. <http://arxiv.org/abs/1312.4658>, 2013.
- [Szy14] Markus Szymik. Commutative \mathbb{S} -algebras of prime characteristics and applications to unoriented bordism. *Algebr. Geom. Topol.*, 14(6):3717–3743, 2014.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, CANADA
E-mail address: oantolin@math.ubc.ca

CENTRE FOR SYMMETRY AND DEFORMATION, UNIVERSITY OF COPENHAGEN, DENMARK
E-mail address: tbarthel@math.ku.dk