

HOMOTOPY THEORY OF DIAGRAMS AND CW-COMPLEXES OVER A CATEGORY

ROBERT J. PIACENZA

Introduction. The purpose of this paper is to introduce the notion of a *CW* complex over a topological category. The main theorem of this paper gives an equivalence between the homotopy theory of diagrams of spaces based on a topological category and the homotopy theory of *CW* complexes over the same base category.

A brief description of the paper goes as follows: in Section 1 we introduce the homotopy category of diagrams of spaces based on a fixed topological category. In Section 2 homotopy groups for diagrams are defined. These are used to define the concept of weak equivalence and J - n equivalence that generalize the classical definition. In Section 3 we adapt the classical theory of *CW* complexes to develop a cellular theory for diagrams. In Section 4 we use sheaf theory to define a reasonable cohomology theory of diagrams and compare it to previously defined theories. In Section 5 we define a closed model category structure for the homotopy theory of diagrams. We show this Quillen type homotopy theory is equivalent to the homotopy theory of J -*CW* complexes. In Section 6 we apply our constructions and results to prove a useful result in equivariant homotopy theory originally proved by Elmendorf by a different method.

1. Homotopy theory of diagrams. Throughout this paper we let Top be the cartesian closed category of compactly generated spaces in the sense of Vogt [10]. Let J be a small topological category over Top with discrete object space and $J\text{-Top}$ the category of continuous contravariant Top valued functors on J . Note that the category $J\text{-Top}$ is naturally enriched in Top . See Dubuc [2] for the framework of enriched category theory. We assume the reader is familiar with the standard constructions in Top as in [10] and the standard functor calculus on $J\text{-Top}$ as in [5, Section 1].

We let I be the unit interval in Top . If X and Y are diagrams then a homotopy from X to Y is a morphism $H: I \times X \rightarrow Y$ of $J\text{-Top}$ where $I \times X$ is the functor defined on objects $j \in |J|$ by $(I \times X)(j) = I \times X(j)$ and similarly for morphisms of J . In the usual way homotopy defines an equivalence relation on the morphisms of $J\text{-Top}$ that gives rise to the quotient homotopy category $hJ\text{-Top}$. We denote the homotopy classes of morphisms

Received by the editors February 3, 1989, revised April 10, 1990.

AMS subject classification: Primary: 55U35.

Key words and phrases: Topological category, homotopy, J -cofibration, weak equivalence, J -*CW* complex, cohomology, equivariant homotopy, fixed point set.


© Canadian Mathematical Society 1991.

from X to Y by $hJ\text{-Top}(X, Y)$ abbreviated $h(X, Y)$. An isomorphism in $hJ\text{-Top}$ is called a J -homotopy equivalence.

A morphism of $J\text{-Top}$ is called a J -cofibration if it has the J homotopy extension property, abbreviated $J\text{-HEP}$. The basic facts about cofibrations in Top apply readily to J -cofibrations. See [5, Section 2].

The following results from [6] apply formally to the category $J\text{-Top}$.

THEOREM 1.1 (INVARIANCE OF PUSHOUTS). *Suppose that we have a commutative diagram:*



in which i and i' are J -cofibrations, f and f' are arbitrary morphisms in $J\text{-Top}$. α , β and γ are homotopy equivalences and the front and back faces are pushouts. Then θ is also a homotopy equivalence (θ being the induced map on pushouts).

THEOREM 1.2 (INVARIANCE OF COLIMITS OVER COFIBRATIONS). *Suppose given a homotopy commutative diagram*

$$\begin{array}{ccccccc}
 X^0 & \xrightarrow{i_0} & X^1 & \xrightarrow{i_1} & \dots & \longrightarrow & X^k & \xrightarrow{i_k} & \dots \\
 \downarrow f^0 & & \downarrow f^1 & & & & \downarrow f^k & & \\
 Y^0 & \xrightarrow{j_0} & Y^1 & \xrightarrow{j_1} & \dots & \longrightarrow & Y^k & \xrightarrow{j_k} & \dots
 \end{array}$$

in $J\text{-Top}$ where the i_k and j_k are J -cofibrations and the f^k are homotopy equivalences. Then the map $\text{colim}_k f^k: \text{colim}_k X^k \rightarrow \text{colim}_k Y^k$ is a homotopy equivalence.

2. Homotopy groups. Let I^n be the topological n -cube and ∂I^n its boundary.

DEFINITION 2.1. By a $J\text{-Top}$ pair (X, Y) , we mean an object X in $J\text{-Top}$ together with a subobject $Y \subseteq X$. Morphisms of pairs are defined in the obvious way. A similar definition will be used for triples, n -ads etc. Let $\varphi: j \rightarrow Y$ be a morphism in $J\text{-Top}$ where $j \in |J|$ is viewed as the representable functor $J\text{-Top}(_, j)$. By Yoneda's theorem φ is completely determined by the point $\varphi(\text{id}_j) = y_0 \in Y(j)$. For each $n \geq 0$, define $\pi_n^j(X, Y, \varphi) = h((I^n, \partial I^n, \{0\}) \times j, (X, Y, Y))$ where $y_0 = \varphi(\text{id}_j) \in Y(j)$ serves as a basepoint, and all homotopies are homotopies of triples relative to φ . The reader may formulate a similar definition for the absolute case $\pi_n^j(X, \varphi)$. For $n = 0$ we adopt the convention that $I^0 = \{0, 1\}$ and $\partial I^0 = \{0\}$ and proceed as above. These constructions extend to covariant functors on $J\text{-Top}$. From now on we shall often drop φ from the notation $\pi_n^j(X, Y, \varphi)$.

The proof of the following proposition follows immediately from Yoneda's lemma.

PROPOSITION 2.2. *There are natural equivalences $\pi_n^j(X) \simeq \pi_n(X(j))$ and $\pi_n^j(X, Y) \simeq \pi_n(X(j), Y(j))$ which preserve the (evident) group structure when $n \geq 1$ (for the absolute case; the relative case requires $n \geq 2$).*

As a direct consequence of 2.2 we obtain the long exact sequences:

PROPOSITION 2.3. *For (X, Y) and j as in 2.1, there exist natural boundary maps ∂ and long exact sequences*

$$\begin{array}{ccccccc} \dots & \longrightarrow & \pi_n^j(X, Y) & \xrightarrow{\partial} & \pi_{n-1}^j(Y) & \longrightarrow & \pi_{n-1}^j(X) \longrightarrow \dots \\ & & & & \longrightarrow & \pi_0^j(Y) & \longrightarrow \pi_0^j(X) \end{array}$$

of groups up to $\pi_1^j(Y)$ and pointed sets thereafter.

DEFINITION 2.4. A map $e: (X, Y) \rightarrow (X', Y')$ of pairs in $J\text{-Top}$ is called a J - n -equivalence if $e(j): (X(j), Y(j)) \rightarrow (X'(j), Y'(j))$ is an n -equivalence in Top for each $j \in |J|$. A map e will be called a *weak equivalence* if e is a J - n -equivalence for each $n \geq 0$. Observe that e is a J - n -equivalence if for every $j \in |J|$ and $\varphi: j \rightarrow Y, e_*: \pi_p^j(X, Y, \varphi) \rightarrow \pi_p^j(X', Y', e\varphi)$ is an isomorphism for $0 \leq p < n$ and an epimorphism for $p = n$. The reader may easily formulate a similar definition for morphisms $e: X \rightarrow X'$ of $J\text{-Top}$ (the absolute case).

3. Cellular theory. In this section we adapt the general treatment of classical homotopy theory and CW -complexes given in [9, Chapter 7] and [6] to develop a good theory of CW -complexes over the topological category J .


Let B^{n+1} be the topological $n + 1$ -ball and S^n the topological n -sphere. Of course, these spaces are homeomorphic to I^{n+1} and ∂I^{n+1} respectively. We shall construct all complexes over J by the process of attaching cells of the form $B^{n+1} \times j$ by attaching morphisms with domain $S^n \times j$. The formal definition goes as follows:

DEFINITION 3.1. A J -complex is an object X of $J\text{-Top}$ with a decomposition $X = \text{colim}_{p \geq 0} X^p$ where $X^0 = \coprod_{\alpha \in A_0} B^{n_\alpha} \times j_\alpha, X^p = X^{p-1} \cup_f \left(\coprod_{\alpha \in A_p} B^{n_\alpha} \times j_\alpha \right)$ for some attaching morphism $f: \coprod_{\alpha \in A_p} S^{n_\alpha-1} \times j_\alpha \rightarrow X^{p-1}$ and for each $p \geq 0, \{j_\alpha \mid \alpha \in A_p\}$ is a collection of objects (representable functors) of J . We call X a J - CW -complex if X is a J -complex as above and for all $p \geq 0$ and all $\alpha \in A_p$ we have $n_\alpha = p$.


A J -subcomplex and a relative J complex are now defined in the obvious way. Without further comment we adopt for J - CW -complexes the standard terminology for CW -complexes. See [9, Chapter 7] and [6].

The following technical lemma and its proof are due to May [6,3.5.1].

LEMMA 3.2. *Suppose that $e: Y \rightarrow Z$ is a J - n -equivalence. Then we can complete the following diagram in $J\text{-Top}$:*



THEOREM 3.3 (J-HELP). *If (X, A) is a relative J -CW complex of dimension $\leq n$ and $e: Y \rightarrow Z$ is a J - n -equivalence then we can complete the following diagram in J -Top:*



PROOF. This follows by induction on $\dim(X, A)$, applying 3.2 cell by cell at each stage.

The proofs of the following Whitehead theorem and cellular approximation theorem are formal modifications of the proofs given in [6].

THEOREM 3.4 (WHITEHEAD). *(i) Suppose X is a J -CW complex, and that $e: Y \rightarrow Z$ is a J - n -equivalence. Then $e_*: h(X, Y) \rightarrow h(X, Z)$ is an isomorphism if $\dim X < n$ and an epimorphism of $\dim X = n$. (ii) If $e: Y \rightarrow Z$ is a weak equivalence, and if X is any J -CW complex, then $e_*: h(X, Y) \rightarrow h(X, Z)$ is an isomorphism.*

THEOREM 3.5 (CELLULAR APPROXIMATION). *Suppose that X is a J -CW complex, and that A is a sub- J -CW complex of X . Then, iff: $X \rightarrow Y$ is a morphism of J -Top which is J -cellular when restricted to A , we can homotope $f, \text{rel } f|_A$ to a J -cellular morphism $g: X \rightarrow Y$.*

Next we discuss the local properties of J -CW-complexes. First we develop some preliminary concepts. Let X be in J -Top and for each $j \in |J|$ let $t_j: X(j) \rightarrow \text{colim}_J X$ be the natural map of $X(j)$ into the colimit. Observe that for each morphism $s: i \rightarrow j$ of J , $t_j = t_i X(s)$. For each subspace $A \subseteq \text{colim}_J X$ we define $\check{A}(j) = t_j^{-1}(A)$ and for a given $s: i \rightarrow j$ we define $\check{A}(s) = X(s)|_{\check{A}(j)}$, the restriction of the continuous map $X(s)$ to the subspace $\check{A}(j)$. We apply the K -ification functor to assure that all spaces defined above are compactly generated. One quickly checks that $\check{A} \in J$ -Top, $\text{colim}_J \check{A} = A$, and there is a natural inclusion morphism $\check{A} \rightarrow X$. To simplify notation from now on we write X/J for $\text{colim}_J X$.

DEFINITION 3.6. By a special pair in J -Top we mean an ordered pair (X, A) where $X \in J$ -Top and $A \subseteq X/J$. We call a special pair (X, A) a J -neighborhood retract pair (abbreviated J -NR) if there exist U an open subset of X/J such that $A \subseteq U$ and there exists a retraction morphism $r: \check{U} \rightarrow \check{A}$. (X, A) is called a J -neighborhood deformation

retract pair (abbreviate J -NDR) if (X, A) is a J -NR and the morphism r is a J -deformation retract.

Let X be a J -CW complex. The functor colim_J sends cells $B^p \times j$ to cells B^p and preserves the cellular decomposition of X . For this reason X/J has the natural structure of a CW-complex in Top with all its attaching maps being images under colim_J of the corresponding attaching morphisms in J -Top. One may also check that if A is a subcomplex of X/J then \check{A} has the natural structure of a subcomplex of X . In particular if A^p is the p -skeleton of X/J then $\check{A}^p = X^p$ is the p -skeleton of X .

THEOREM 3.7 (LOCAL CONTRACTIBILITY). *Let (X, A) be a special pair in J -Top with X a J -CW complex and $A = \{a\}$, $a \in X/J$. Then there exists a unique object $j \in J$ such that $\check{A} \simeq j$ (j viewed as a representable functor) and (X, A) is a J -NDR pair.*

PROOF. Suppose $a \in (X/J)^p \setminus (X/J)^{p-1}$, the p -skeleton minus the $p-1$ skeleton of X/J . Then there is a unique attaching morphism f in J -Top

$$\begin{array}{c} f: S^{p-1} \times j \rightarrow X^{p-1} \\ \downarrow \\ B^p \times j \end{array}$$

with a in the interior of B^p . It follows that $\check{A} \simeq j$ for the unique choice of j given above. To construct the required neighborhood U first take an open ball U_1 contained in the interior of B^p and centered at a . Then U_1 is a neighborhood in $(X/J)^p$ contracting to A . One then extends U_1 inductively cell by cell by a well known procedure to construct the required neighborhood U .

THEOREM 3.8. *Let (X, A) be a special pair in J -Top with X a J -CW complex and A an arbitrary subcomplex of X/J . Then (X, A) is a J -NDR pair.*

PROOF. It follows from 3.3 that $\check{A} \subseteq X$ is a J -cofibration. The result then follows from a well known argument of Puppe. See [5, Lemma 4.3, p. 193].

4. Cohomology. In this section we use sheaf theory to construct a cohomology theory on J -Top satisfying a suitably formulated set of Eilenberg-Steenrod axioms. We refer the reader to Bredon [1] for the basic definitions and terminology of sheaf theory.

DEFINITION 4.1. By a contravariant coefficient system M on J we mean a continuous contravariant functor $M: J \rightarrow \text{Ab}$ where Ab is the category of discrete abelian groups. Observe that every contravariant coefficient system M is a homotopy invariant functor in the following sense. If $f, g: j \rightarrow j'$ are homotopic (as morphisms of representable functors in J -Top) then $M(f) = M(g)$.

Let $X \in J$ -Top and let M be a coefficient system on J . We define a presheaf of abelian groups M^X over X/J as follows: for $A \subseteq X/J$ define $M^X(A) = J\text{-Top}(\check{A}, M)$ equipped with its natural discrete abelian group structure. If $B \subseteq A$ there is a natural restriction homomorphism $M^X(A) \rightarrow M^X(B)$ and one easily checks that M^X is a sheaf of abelian

groups over X/J . Let $f: X \rightarrow Y$ be a morphism in $J\text{-Top}$ with $f/J: X/J \rightarrow Y/J$ the induced map in Top . There is a natural f/J -cohomomorphism of sheaves $\bar{f}: M^Y \rightarrow M^X$ given by the obvious composition with f .

DEFINITION 4.2. Let $X \in J\text{-Top}$, ψ a family of supports on X/J and M a coefficient system on J . We define $H_\psi^n(X; M) = H_\psi^n(X/J; M^X)$ where the right side is sheaf cohomology with supports ψ as defined in [1, Chapter II]. Given a morphism $f: X \rightarrow Y$ in $J\text{-Top}$, we let f^* be the homomorphism induced in cohomology by f . Given a special pair (X, A) we define the relative cohomology $H_\psi^n(X, A; M) = H_\psi^n(X/J, A; M^X)$ where the right side is relative sheaf cohomology.

EXAMPLE 4.3. Let G be an abelian group and define the constant coefficient system M with value G by setting $M(s) = \text{id}_G$ for any morphism s of J . Then for any $X \in J\text{-Top}$ one quickly sees that $H^*(X; M) = H^*(X/J; G)$ where the right side is sheaf cohomology with constant coefficients G . Note that absence of a specified support family always means supports in the family of all closed sets.

DEFINITION 4.4. A special pair (X, A) in $J\text{-Top}$ is called *acceptable* if for each coefficient system M on J the sheaf M^A over A is the restriction of the sheaf M^X to the subspace A . Note that if (X, A) is a $J\text{-NR}$ pair or if X is locally $J\text{-NR}$ then (X, A) is acceptable. In particular any special pair (X, A) where X is a $J\text{-CW}$ complex is acceptable by 3.7.

All special pairs considered in the rest of this section will be assumed acceptable. We impose this condition to obtain a good theory of relative cohomology.

Note that a supports preserving morphism $f: (X, A) \rightarrow (Y, B)$ naturally induces a homomorphism f^* in relative cohomology. Hence $H_\psi^*(\ ; M)$ becomes a candidate for a reasonable cohomology theory on $J\text{-Top}$. The following theorem states and verifies a suitable set of Eilenberg-Steenrod axioms for the theory $H^*(\ ; M)$.

THEOREM 4.5.

- (1) (Dimension) $H^n(j; M) = \begin{cases} M(j) & n = 0 \\ 0 & n > 0 \end{cases}$ for each $j \in J$ viewed as a representable functor.
- (2) For each special pair (X, A) in $J\text{-Top}$ there is induced a suitable long exact sequence in cohomology with arbitrary supports.
- (3) (Excision) If A and B are subsets of X/J with $\bar{B} \subseteq \text{int}A$ then the inclusion $i: (X - \bar{B}, A - B) \rightarrow (X, A)$ induces an isomorphism in cohomology for any support family.
- (4) (Homotopy) If f and g are morphisms of special pairs in $J\text{-Top}$ that are homotopic via a support preserving homotopy then $f^* = g^*$.
- (5) If $(X, A) = \coprod_\alpha (X_\alpha, A_\alpha)$ then there is a natural isomorphism induced by the injections into the coproduct,

$$H^*(X, A; M) \simeq \coprod_\alpha H^*(X_\alpha, A_\alpha; M).$$

PROOF. (1) follows from Yoneda's lemma. (2) follows from [1, Chapter 2, Section 12]. (3) follows from [1, Theorem 12.5, p. 61]. (4) follows from [1, Theorem 11.2, p. 55]. (5) is easy to check directly.

If X is a J -CW complex we define cellular cochains $C^n(X; M) = H^n(X^n, (X^{n-1}/J); M)$. Observe that $C^n(X; M) = \prod_{\alpha} M(j_{\alpha})$ where $B^n \times j_{\alpha}$, $\alpha \in A_n$ is the family of all n -cells of X . In the usual way one makes $C^*(X; M)$ into a cochain complex using the coboundary operator of a triple. This construction yields the cellular cohomology theory $H^*_{\text{cel}}(\quad; M)$ defined for J -CW pairs.

We may adapt the classical proof to show:

PROPOSITION 4.6. $H^*(\quad; M)$ is naturally isomorphic to $H^*_{\text{cel}}(\quad; M)$ on the category of J -CW pairs.

REMARK 4.7. (i) The cellular cohomology theory is useful for developing an obstruction theory in J -Top. (ii) Following a well known argument due to Milnor it is possible to prove a uniqueness theorem for cohomology theories defined on the category of J -CW complexes. (iii) In [11] Vogt defines the singular cohomology on J -Top and shows it satisfies a suitable set of axioms. By the above mentioned uniqueness theorem Vogt's singular cohomology agrees with our sheaf cohomology on the category of J -CW complexes.

5. **Closed model structure on J -Top.** In [8] Quillen defines a closed model structure for homotopy theory in Top. In this section we emulate this construction to define a closed model category structure on J -Top.

DEFINITION 5.1. A morphism $f: X \rightarrow Y$ of J -Top is called a *weak fibration*, abbreviated w -fibration, if for each $j \in J$, $f(j): X(j) \rightarrow Y(j)$ is a Serre fibration in Top. See [9, p. 374] for a discussion of Serre fibrations. Observe that f is a w -fibration if f has the homotopy lifting property for all objects of the form $I^n \times j$. A morphism f is called a *weak equivalence* if f is a *weak equivalence* as defined in Section 2. A morphism $g: A \rightarrow B$ is called a *weak cofibration*, abbreviated w -cofibration if g has the left lifting property (LLP) for each trivial w -fibration $f: X \rightarrow Y$ (a w -fibration that is also a weak equivalence). This means one can always fill in the dotted arrow:

$$\begin{array}{ccc} A & \longrightarrow & X \\ g \downarrow & \dashrightarrow & \downarrow f \\ B & \longrightarrow & Y \end{array}$$

REMARK 5.2. (i) The inclusion of a sub- J -complex into a J -complex is always both a J -cofibration and a w -cofibration. (ii) A w -fibration is trivial iff it has the right lifting property (RLP) for each w -cofibration of the form $S^n \times j \rightarrow B^{n+1} \times j$. [8, 3.2, Lemma 2].

LEMMA 5.3 (QUILLEN'S FACTORIZATION LEMMA). *Any morphism $f: X \rightarrow Y$ of $J\text{-Top}$ may be factored $f = pg$ where g is a w -cofibration and p is a trivial w -fibration.*

PROOF. We construct a diagram

$$\begin{array}{ccccccc} X & \xrightarrow{g_0} & Z^0 & \xrightarrow{g_1} & Z^1 & \longrightarrow & \dots \\ & f \searrow & \downarrow p_0 & \swarrow p_1 & & & \\ & & Y & & & & \end{array}$$

as follows: let $Z^{-1} = X$ and $p_{-1} = f$, and having obtained Z^{n-1} consider the set of all diagrams of the form

$$\begin{array}{ccc} S^{q_\alpha} \times j_\alpha & \xrightarrow{t_\alpha} & Z^{n-1} \\ \downarrow & & \downarrow p_{n-1} \\ B^{q_\alpha+1} \times j_\alpha & \xrightarrow{s_\alpha} & Y \end{array}$$

where we have indexed this set of diagrams by A_n and $\alpha \in A_n$. Define $g_n: Z^{n-1} \rightarrow Z^n$ by the pushout diagram

$$\begin{array}{ccc} \coprod_{\alpha \in A_n} S^{q_\alpha} \times j_\alpha & \xrightarrow{\coprod t_\alpha} & Z^{n-1} \\ \downarrow & & \downarrow g_n \\ \coprod_{\alpha \in A_n} B^{q_\alpha+1} \times j_\alpha & \longrightarrow & Z^n \end{array}$$

Throughout this construction we have included the use of the trivial sphere i.e., $S^{-1} = \emptyset, B^0 = \{\text{pt}\}$. Define $p_n: Z^n \rightarrow Y$ by $p_n g_n = p_{n-1}, p_n i_{n2} = \coprod s_\alpha$, let $Z = \text{colim } Z^n, p = \text{colim } p_n$ and $g = \text{colim } g_n g_{n-1} \cdots g_0$. One may check that g has LLP with respect to each trivial w -fibration and by the small object argument [8, 3.4, Remark] p is a trivial w -fibration.

THEOREM 5.4. *With the structure defined above (Definition 5.1) $J\text{-Top}$ is a closed model category.*

PROOF. One quickly checks the axioms for a closed model category [8, 3.1] using 5.3 or its clone to verify the factorization axiom M2.

We let $\text{Ho } J\text{-Top}$ be $J\text{-Top}$ localized at the weak equivalences. We aim to show that $\text{Ho } J\text{-Top}$ is equivalent to the homotopy theory of $J\text{-CW}$ complexes. First we need the following.

LEMMA 5.5. *Let $X = \text{colim } X_n$ taken over a system of J -cofibrations such that each X_n has the J -homotopy type of a $J\text{-CW}$ -complex. Then X has the J -homotopy type of a $J\text{-CW}$ complex.*

PROOF. Replace the colimit by the telescope [6, 1.26] and use the homotopy invariance of the homotopy colimit (Theorem 1.2).

The following proposition follows easily.

PROPOSITION 5.6. *Each J -complex is of the J -homotopy type of a J -CW-complex.*

THEOREM 5.7 (APPROXIMATION THEOREM). *There is a functor $\Gamma: J\text{-Top} \rightarrow J\text{-Top}$ and natural transformation $p: \Gamma \rightarrow \text{id}$ such that for each $X \in J\text{-Top}$, ΓX is a J -complex, and p_X is a trivial w -fibration.*

PROOF. Using 5.3 factor the map $\phi \subseteq X$ into $\phi \subseteq \Gamma X \rightarrow X$ where ϕ is the empty subfunctor of X . Then by the construction in 5.3 we see that X is a J -complex, p_X is a trivial fibration, Γ is a functor, and p a natural transformation.

The following corollary is immediate from 5.6 and 5.7.

COROLLARY 5.8. *The category $\text{Ho } J\text{-Top}$ is equivalent to the category of J -CW complexes modulo homotopy.*

REMARK 5.9. (i) In [9, Theorem 1, p. 412] Spanier makes use of Brown's representability theorem [9, Theorem 11, p. 410] to construct CW approximations in the category Top . In our construction we do not need Brown's theorem and furthermore we construct the useful approximating functor Γ directly on $J\text{-Top}$. We believe this is an improvement over Spanier's construction. (ii) In [5] Heller describes a somewhat different homotopy structure on $J\text{-Top}$. One may check that Heller's localization $\text{Ho}_w \text{Top}^J$ of [5, Section 7] is equivalent to our $\text{Ho } J\text{-Top}$. It follows that many of the results of [5] (homotopy Kan extensions, etc.) may be applied to $\text{Ho } J\text{-Top}$.

6. Elmendorf's Theorem. The purpose of this section is to prove a useful result in equivariant homotopy theory originally proved by Elmendorf in [4] by a different method.

Let G be a topological group and let $G\text{-Top}$ be the category of right G -spaces in Top . Let O_G be the topological category of canonical right orbits. An object of O_G is a closed subgroup $H \subseteq G$ and $O_G(H, K) = G\text{-Top}(G/H, G/K)$ is given the compact open topology. Observe that there is a natural bijection $G\text{-Top}(G/H, G/K) \simeq [G/K]^H$. Where the right side is the H fixed point set of the right orbit G/K . This bijection is a homeomorphism if we impose (as we always do) the compactly generated topology on all spaces in sight. There is a full and faithful functor $\Phi: G\text{-Top} \rightarrow O_G\text{-Top}$ which views each $X \in G\text{-Top}$ as a continuous diagram $\Phi(X)$ of fixed point sets. $\Phi(X)$ is defined by setting $\Phi(X)(H) = G\text{-Top}(G/H, X)$. That is $\Phi(X)$ is the continuous functor $G\text{-Top}(_, X)$ on O_G . Compare [4, Section 1]. We call $f: X \rightarrow Y$ a G -weak equivalence (G -fibration) if $\Phi(f)$ is a weak equivalence (w -fibration in $O_G\text{-Top}$).

In $G\text{-Top}$ there is a well-known theory of G -complexes (G -CW-complexes) that uses cells of the form $B^n \times G/H$. See [12, Section 3] for a discussion of equivariant cellular theory. Observe that under the functor Φ , $B^n \times G/H$ goes to $B^n \times O_G(_, G/H)$, i.e., B^n cross a representable functor.

We need the following lemma for the argument below.

LEMMA 6.1. *If*

$$\begin{array}{ccc} B & \xrightarrow{i} & C \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X \end{array}$$

is a pushout in $G\text{-Top}$ with i a closed inclusion then

$$\begin{array}{ccc} \Phi B & \xrightarrow{\Phi(i)} & \Phi C \\ \downarrow & & \downarrow \\ \Phi Y & \longrightarrow & \Phi X \end{array}$$

is a pushout in $O_G\text{-Top}$.

PROOF. Stripping away the topology we see this holds on the set level since every G -set is a coproduct of orbits. One may then check that the topologies agree.

THEOREM 6.2. *Each O_G -complex (O_G -CW-complex) $Y \in O_G\text{-Top}$ is isomorphic to ΦX where X is a G -complex (G -CW-complex) in $G\text{-Top}$. It follows that Φ is an isomorphism between the categories of G -complexes (G -CW-complexes) and O_G -complexes (O_G -CW-complexes).*

PROOF. The assertion follows from 6.1 and the fact that Φ is full, faithful and preserves ascending unions.

THEOREM 6.3. *There is a functor $A: O_G\text{-Top} \rightarrow G\text{-Top}$ and natural transformation $t: \Phi A \rightarrow \text{id}$ such that ΦAX is an O_G -complex and t_X is a trivial fibration for each $X \in O_G\text{-Top}$. It follows that there is an equivalence of categories $\text{Ho } O_G\text{-Top} \sim \text{Ho } G\text{-Top}$ where $\text{Ho } G\text{-Top}$ is $G\text{-Top}$ localized at the weak equivalences in $G\text{-Top}$.*

PROOF. We construct A and t using the functor Γ and transformation p given in 5.7. The result follows from 5.8 and 6.2.

COROLLARY 6.4. *Let $Y \in G\text{-Top}$ be G homotopically equivalent to a G -CW complex. Then for any $X \in O_G\text{-Top}$, $hG\text{-Top}(Y, AX) \simeq hO_G\text{-Top}(\Phi Y, X) \simeq \text{Ho } O_G\text{-Top}(\Phi Y, X)$.*

PROOF. This follows from 6.3 and generalities about closed model categories.

REMARK 6.5. (i) In [4] Elmendorf assumes G is a compact Lie group and uses a generalized bar construction to obtain his version of 6.3 and 6.4. Let $C: O_G\text{-Top} \rightarrow G\text{-Top}$ be the functor defined by Elmendorf [4, Theorem 1]. For $X \in O_G\text{-Top}$ there is a natural G weak equivalence $AX \rightarrow CX$ which is a G homotopy equivalence if X is regular in the sense of Elmendorf. Clearly the functors A and C are closely related.

(ii) The importance of having the approximation functor A given above is demonstrated by several applications given by Elmendorf in [4, Section 2]. For example consider the following. Let \mathcal{F} be an orbit family in G and define $T \in O_G\text{-Top}$ by:

$$T(H) = \begin{cases} \text{one point} & \text{if } H \in \mathcal{F} \\ \text{empty} & \text{otherwise.} \end{cases}$$

Then $AT = E\mathcal{F}$ is a universal \mathcal{F} -space and $B\mathcal{F} = \text{Ho colim } T = \text{colim } \Gamma T = E\mathcal{F}/G$ is a classifying space for the orbit family \mathcal{F} . If \mathcal{F} consists of the single trivial subgroup of G then $B\mathcal{F} = BG$ is a classifying space for principal G bundles.

(iii) Let $M: O_G \rightarrow \text{Ab}$ be a coefficient system on O_G . One defines equivariant cohomology with coefficients M denotes $H_G^*(X; M)$ by setting $H_G^*(X; M) = H^*(\Phi X; M)$ for $X \in G\text{-Top}$. The results of Section 4 show this definition gives a reasonable cohomology theory on $G\text{-Top}$. Observe that under suitable conditions this theory agrees with Illman's equivariant singular cohomology. See [7, Theorem 3.11].

REFERENCES

1. Glen E. Bredon, *Sheaf theory*. McGraw-Hill, 1967.
2. Eduardo J. Dubuc, *Kan extensions in enriched category theory*. Springer lecture notes in Math **145**(1970).
3. W. G. Dwyer and D. M. Kan, *Singular functors and realization functors*, Proceed. of the Koninlijke Nederlandse Akademie Van Wetenschappen, Series A., **87**(1984), 147–153.
4. A. D. Elmendorf, *Systems of fixed point sets*, Trans. Amer. Math. Soc. **277**(1983), 275–284.
5. Alex Heller, *Homotopy in functor categories*, Trans. Amer. Math. Soc. **272** (1982), 185–202.
6. Peter May, *The homotopical foundations of algebraic topology*, mimeographed notes, University of Chicago.
7. R. Piacenza, *Cohomology of diagrams and equivariant singular theory*, Pacific J. Math. **91**(1981), 435–444.
8. D. Quillen, *Homotopical algebra*. Springer lecture notes in Math **43**(1967).
9. E. Spanier, *Algebraic topology*. McGraw-Hill, 1966.
10. R. Vogt, *Convenient categories of topological spaces for homotopy theory*, Archiv der Mathematik, **XXII** (1971), 545–555.
11. ———, *Equivariant singular homology and cohomology*, preprint.
12. S. Waner, *Equivariant homotopy theory and Milnor's theorem*, Trans. Amer. Math. Soc. **258**(1980), 351–368.

Department of Mathematical Sciences
University of Alaska Fairbanks
Fairbanks, Alaska 99775-1110
 USA