Thick ideals in equivariant and motivic stable homotopy categories

Ruth Joachimi

March 31, 2015

Contents

1	Intro	oduction	3
2	Thic	k ideals in classical stable homotopy theory	9
3	Thic	k ideals in equivariant stable homotopy theory	12
	3.1	Equivariant stable homotopy theory	12
	3.2	Equivariant Morava K-theories	14
	3.3	Nilpotence and lattices of thick ideals	16
	3.4	Thick ideals and equivariant Morava K-theories	21
	3.5	Thick ideals in $\mathcal{SH}(\mathbb{Z}/2)_f$	24
4	Com	parison functors	26
	4.1	Symmetric $\mathbb{C}P^1$ -spectra	26
	4.2	$\mathbb{Z}/2$ -equivariant symmetric spectra	27
	4.3	Complex and real topological realisation functors	28
	4.4	Realisation functors for other fields	32
	4.5	Constant presheaf functors	32
5	Thic	k ideals discovered by comparison functors	37
	5.1	Consequences of the properties of R_k , R'_k , c_k and c'_k	37
	5.2	Finite motivic spectra	38
	5.3	Motivic thick ideals	42
6	Thic	k ideals associated with cohomology theories	46
	6.1	Equivalence of homology and cohomology theories	46
	6.2	Thick ideals	48
	6.3	Morava K-theories in $\mathcal{SH}(k)$	48
		6.3.1 Construction and properties of $AK(n)$	49
		6.3.2 Thick ideals and Morava K-theories	51
7	$\mathcal{SH}(P)$	$(k)_f$ has more thick ideals than \mathcal{SH}^{fin}	53
	7.1	The motivic Hopf map	53
	7.2	Prime ideals	57
		7.2.1 Prime ideals in the topological categories \mathcal{SH}^{fin} and $\mathcal{SH}(\mathbb{Z}/2)_f$.	58

		7.2.2 Prime ideals in the motivic category $\mathcal{SH}(k)_f$	50
8	Mot	tivic type-n spectra 6	3
	8.1	Universal coefficient and Künneth theorems	54
	8.2	The motivic Steenrod algebra	55
	8.3	The motivic Adams spectral sequence	6
	8.4	Vanishing criterion for motivic Morava K-theory	58
	8.5	Construction of an X such that $H^{**}X$ is free over $\Lambda(Q_s)$	58
		8.5.1 Idempotents for free H ^{**} -modules	;9
		8.5.2 A finite cell spectrum with nontrivial Q_s -action and trivial Q_n -action 7	'1
	8.6	The type of the realisation $R(X)$	'3
	8.7	The constant type-n spectrum	'3
9	Bou	sfield classes 7	6
	9.1	v_n -Torsion	$^{\prime}7$
	9.2	Properties of Bousfield classes	31
	9.3	The action of v_i on $AP(n)$	32
	9.4	Bousfield classes of $AK(n)$ and $AB(n)$)2
	9.5	Decomposition of $\langle AE(n) \rangle$)5
	9.6	AK(n) and $AK(n+1)$.)5

Chapter 1

Introduction

In a tensor triangulated category, a thick ideal is a full subcategory which is closed under exact triangles and retracts and under tensoring with arbitrary elements of the category. The classification of thick ideals in the stable homotopy category of *p*-local finite spectra, $S\mathcal{H}_{(p)}^{fin}$, is given by a famous theorem of Hopkins and Smith, [HS98, Theorem 7], see Section 2. It states that, in $S\mathcal{H}_{(p)}^{fin}$, the thick ideals are given as a chain

$$\mathcal{SH}_{(p)}^{fin} = \mathcal{C}_0 \supseteq \mathcal{C}_1 \supseteq \cdots \supseteq \mathcal{C}_n \supseteq \cdots \supseteq \mathcal{C}_\infty = \{0\},$$

and each thick ideal is characterised by the vanishing of a particular Morava K-theory, that is, $C_n = \{X \in \mathcal{SH}_{(p)}^{fin} \mid K(p, n-1)_*(X) = 0\}$ for $0 < n < \infty$. This theorem is a consequence of the nilpotence theorem [HS98, Theorem 3], the existence of type-*n* spectra for any $n \ge 0$ [Mit85, Theorem 4.8] and the fact that $K(n)_*(X) = 0$ implies $K(n-1)_*(X) = 0$ for $X \in \mathcal{SH}^{fin}$ [Rav84, Theorem 2.11].

In equivariant stable homotopy theory for a finite group G, unpublished work of Strickland [Str10], see Chapter 3, contains a partial classification of thick ideals in the category $\mathcal{SH}(G)_f \subset \mathcal{SH}(G)$, which is the full subcategory of compact objects in the G-equivariant stable homotopy category. This is a generalisation of the above result, which concerns the special case $\mathcal{SH}(\{1\})_f = \mathcal{SH}^{fin}$. In $\mathcal{SH}(G)_f$, any thick ideal is characterised by the vanishing of particular equivariant Morava K-theories, which are indexed by a prime and a nonnegative integer (as the ordinary Morava K-theories) and, additionally, by a conjugacy class of subgroups of G. The set of thick ideals can be mapped to a lattice of such multi-indices and Strickland proves lower and upper bounds for a sublattice onto which this map is bijective.

In this thesis, we study thick ideals in $\mathcal{SH}(k)$, $k \subseteq \mathbb{C}$, and related motivic categories, like $(\mathcal{SH}(k)_f)_{(p)}$, the *p*-localisation of the full subcategory of all compact objects, and $\mathcal{SH}(k)_{(p)}^{fin}$, the category of *p*-localised finite cell spectra. We use different approaches, all of which are, in some sense, motivated by the results about thick ideals in \mathcal{SH}^{fin} .

One approach is to use the comparison functors,

$$\mathcal{SH} \xrightarrow{c_k} \mathcal{SH}(k) \xrightarrow{R_k} \mathcal{SH}$$

for $k \subseteq \mathbb{C}$, and also

$$\mathcal{SH}(\mathbb{Z}/2) \xrightarrow{c'_k} \mathcal{SH}(k) \xrightarrow{R'_k} \mathcal{SH}(\mathbb{Z}/2)$$

for $k \subseteq \mathbb{R}$. We show that, for $k \subseteq \mathbb{C}$, the preimages $R_k^{-1}(\mathcal{C}_n) \subseteq (\mathcal{SH}(k)_f)_{(p)}, n \ge 0$, form a chain of different thick ideals in $(\mathcal{SH}(k)_f)_{(p)}$ (Theorem 5.3.1). For $k \subseteq \mathbb{R}$, we also show that $(R'_k)^{-1}(\mathcal{C}) \subseteq (\mathcal{SH}(k)_f)_{(p)}$ are different thick ideals, where \mathcal{C} runs over all thick ideals in $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$, as studied in [Str10] and in Chapter 3.

The second approach is to use methods of nilpotence theory. The thick subcategory theorem for SH is highly related to the nilpotence theorem, which states that the cobordism spectrum MU detects certain kinds of nilpotence. In this context, the Morava K-theories recover the information from MU, meaning that also the family $\{K(p,n) \mid p \text{ prime}, n \geq 0\}$ detects nilpotence. Since the Morava K-theories have a particularly easy structure (they are field theories satisfying the Künneth formula, see e.g. [Rav92, page 176]), this can be used to show that any thick ideal in SH^{fin} can be uniquely described in terms of the vanishing and non-vanishing of Morava K-theories. This is how [HS98, Theorem 7] is proven. Similar nilpotence arguments are used in [Str10] to classify thick ideals in $SH(G)_f$ for finite groups G. Strickland shows that the equivariant Morava K-theories detect nilpotence (Theorem 3.4.2).

The motivic analog to MU is the algebraic cobordism spectrum MGL. But here, the situation is different, as MGL does not detect nilpotence. There is a notion of motivic Morava K-theories, which are more complicated than the topological ones, but they, too, describe a certain family of thick ideals. In Chapter 7, we show that they do not discover all thick ideals in $(\mathcal{SH}(k)_f)_{(p)}$, and also that not all thick ideals are of the form $R_k^{-1}(\mathcal{C}_n)$.

The third approach is to find different lifts of topological type-*n* spectra to the motivic world and to ask whether they generate the same thick ideals or different ones. We consider two explicit different such lifts to $(\mathcal{SH}(\mathbb{C})_f)_{(p)}$ and show that the motivic Morava K-theories do not distinguish the thick ideals generated by them. The question whether the two lifts generate different thick ideals remains open.

In $\mathcal{SH}_{(p)}^{fin}$, the thick ideals are ordered linearly by inclusion, due to the fact that $K(n+1)_*(X) = 0$ implies $K(n)_*(X) = 0$ for $X \in \mathcal{SH}^{fin}$. This raises the question whether this implication also holds in $\mathcal{SH}(k)_f$. For p > 2, we prove that the analog statement holds for motivic Morava K-theories over \mathbb{C} if X is a finite cellular motivic spectrum, as studied in [DI05]. That is, it holds for $X \in \mathcal{SH}(\mathbb{C})^{fin} \subseteq \mathcal{SH}(\mathbb{C})_f$. On the way, we prove a couple of interesting facts concerning the motivic versions of BP, K(n) and related theories. We prove that the analog of the decomposition of Bousfield classes $\langle E(n) \rangle = \bigvee_{i \leq n} \langle K(i) \rangle$ holds in $\mathcal{SH}(\mathbb{C})$ (for p > 2), as conjectured by Hornbostel in [Hor06, Question 2.17] for arbitrary fields.

Outline.

In Chapter 2, we introduce basic notation concerning thick ideals and the stable

homotopy category SH. We recall the thick subcategory theorem of Hopkins and Smith (Theorem 2.0.10). In SH^{fin} , there is no difference between thick subcategories and thick ideals (Lemma 2.0.8).

Chapter 3 is an account on Strickland's work [Str10]. It contains Strickland's main results on thick ideals in $\mathcal{SH}(G)_f$ and their proofs. The chapter begins with the necessary recollection from equivariant stable homotopy theory, such as compact objects in $\mathcal{SH}(G)$ and geometric fixed point functors. Equivariant Morava K-theories

$$K(n,H) = G/H_+ \wedge E[\geq H] \wedge K(n), \text{ for } H \subseteq G,$$

are introduced in Section 3.2. They are related to the classical Morava K-theories via the geometric fixed point functor (Proposition 3.2.1) and satisfy similar properties, such as the Künneth formula (Corollary 3.2.2). Section 3.3 introduces the terminology of lattices and contains the result of Strickland which establishes a general relation between thick ideals and the detection of nilpotence by some family of homology theories (Theorem 3.3.12). The equivariant analog of the nilpotence theorem [HS98, Theorem 3] is Theorem 3.4.2. We give a reformulation of the thick subcategory theorem [HS98, Theorem 7] in a non-*p*-localised way (Theorem 3.4.5) and prove a similar equivariant result, Theorem 3.4.6, which describes an injective lattice homomorphism from the set of thick ideals in $S\mathcal{H}(G)_f$ to the lattice

$$GQ = \prod_{\text{sub}(G)} \{ u \in \prod_{p} \mathcal{Q}_{p} \mid u_{p} = 1 \,\forall \, p \text{ or } u_{p} \neq 1 \,\forall \, p \},\$$

where $Q_p = \{p^{-n} \mid 0 \le n \le \infty\}$, and gives a lower bound for its image:

Theorem 3.4.6. (Strickland)

The composition

$$\tau: \mathrm{Idl}(\mathcal{SH}(G)_f) \xrightarrow{\mathrm{supp}} \mathcal{P}(GQ') \xrightarrow{\mathrm{max}} GQ$$

is injective. Its image contains all $u \in GQ$ which satisfy: if $H \subseteq H'$, then $u_H \ge u_{H'}$.

Here, $\mathcal{P}(GQ')$ denotes the power set of the set

$$GQ' = \{p^{-n} \mid p \text{ prime}, 0 \le n < \infty\} \times \mathrm{sub}(G).$$

An upper bound is given in Proposition 3.4.7. In Section 3.5, we apply Strickland's results to $\mathcal{SH}(\mathbb{Z}/2)_f$, which will be most interesting to us in our study of thick ideals in $\mathcal{SH}(k)_f$, $k \subseteq \mathbb{R}$. Any thick ideal in $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ is of the form

$$\mathcal{C}_{m,n} = \{ X \mid \phi^{\{1\}}(X) \in \mathcal{C}_m \text{ and } \phi^{\mathbb{Z}/2}(X) \in \mathcal{C}_n \},\$$

where $m, n \in [0, \infty]$ and $\phi^H : S\mathcal{H}(G) \to S\mathcal{H}$ is the geometric *H*-fixed point functor (Corollary 3.5.1). But not all $\mathcal{C}_{m,n}$ are different. Corollary 3.5.3 gives partial information on which ones are.

In **Chapter 4**, we introduce the comparison functors $\mathcal{SH} \xrightarrow{c_k} \mathcal{SH}(k) \xrightarrow{R_k} \mathcal{SH}$ for $k \subseteq \mathbb{C}$, and $\mathcal{SH}(\mathbb{Z}/2) \xrightarrow{c'_k} \mathcal{SH}(k) \xrightarrow{R'_k} \mathcal{SH}(\mathbb{Z}/2)$ for $k \subseteq \mathbb{R}$, which are symmetric monoidal and satisfy $R_k \circ c_k \cong$ id and $R'_k \circ c'_k \cong$ id, respectively. This is mainly a recollection from various other sources. The same results are independently obtained in [HO14, Section 4].

In **Chapter 5**, we apply our knowledge concerning comparison functors to the study of thick ideals, proving the following theorem for any prime p.

Theorem 5.3.1. (Lower bound on the number of motivic thick ideals)

- (1) If $k \subseteq \mathbb{C}$, the category $(\mathcal{SH}(k)_f)_{(p)}$ contains at least an infinite chain of different thick ideals given by $\overline{R}_k^{-1}(\mathcal{C}_n)$, $0 \le n \le \infty$, where \overline{R}_k denotes the p-localisation of the restriction of R_k to $\mathcal{SH}(k)_f$ and $\mathcal{C}_n \subseteq \mathcal{SH}_{(p)}^{fin}$ is as defined in Chapter 1.
- (2) If $k \subseteq \mathbb{R}$, then $(\mathcal{SH}(k)_f)_{(p)}$ contains at least a two-dimensional lattice of different thick ideals given by $(\overline{R}'_k)^{-1}(\mathcal{C}_{m,n})$, for all $(m,n) \in \Gamma_p$ as in Definition 3.5.2.

One ingredient of this theorem is Proposition 5.2.10, where we show that the realisation functors R_k and R'_k preserve compactness. In Section 5.3, we also prove a couple of additional results on the connection between motivic thick ideals and the comparison functors.

Chapter 6 begins with an account of homology and cohomology theories in the category of finite motivic cell spectra, $\mathcal{SH}(k)^{fin}$, as studied by Dugger and Isaksen in [DI05]. We show that, for a cellular ring spectrum E and a finite cellular spectrum X, $E_{**}X = 0$ is equivalent to $E^{**}X = 0$ (Proposition 6.1.1). For $k \subseteq \mathbb{R}$, we use a notion of cellular spectra which is more general than the notion from [DI05], see Definition 5.2.1. This yields another version of Proposition 6.1.1 (Corollary 6.1.4). In Section 6.2, we discuss different ways of defining thick ideals associated with a (ring) spectrum. This is applied to motivic Morava K-theories AK(n) in Section 6.3. For example, we show that the thick ideal $\mathcal{C}_{AK(n)}$ associated with the *n*-th motivic Morava K-theories of the motivic Morava K-theories in Section 6.3.1. The motivic Atiyah Hirzebruch spectral sequence described in [Hoy13, Example 8.13], implies that the *n*-th motivic Morava K-theory as K-theory over the field \mathbb{C} has coefficient ring $H\mathbb{Z}/p_{**} \otimes K(n)_*$ (Lemma 6.3.7), as remarked in [Yag05] below Corollary 3.9.

In **Chapter 7**, we study the thick ideal generated by the cofiber of the motivic Hopf map, $C\eta \cong \mathbb{P}^2_k$, and compare it to the thick ideals $R_k^{-1}(\mathcal{C}_n)$ and $\mathcal{C}_{AK(n)}$ for $k \subseteq \mathbb{C}$. We calculate the type of $R_k(C\eta_{(p)}) \in \mathcal{SH}_{(p)}^{fin}$, which is 1, and the equivariant type of $R'_k(C\eta_{(p)}) \in (\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$, which is (1, 2) for p = 2 and $(1, \infty)$ for odd p (Proposition 7.1.3). In Proposition 7.1.4, we show that $C\eta_{(p)}$ generates a thick ideal of $(\mathcal{SH}(k)_f)_{(p)}$ which is neither of the form $R_k^{-1}(\mathcal{C}_{n+1})$ or $\mathcal{C}_{AK(n)}$ for any $n \ge 0$, nor is it all of $(\mathcal{SH}(k)_f)_{(p)}$ (at least, if p = 2 or $k \subseteq \mathbb{R}$). **Proposition 7.1.4.** For $k \subseteq \mathbb{C}$, let thickid $(C\eta_{(p)}) \subseteq (\mathcal{SH}(k)_f)_{(p)}$ denote the thick ideal generated by the p-localised cofiber of the Hopf map. Then the following hold:

- (1) thickid $(C\eta_{(p)}) \not\subseteq \mathcal{C}_{AK(n)}$ for any $n \ge 0$ and any prime p,
- (2) thickid $(C\eta_{(p)}) \not\subseteq R_k^{-1}(\mathcal{C}_n)$ for any n > 0 and any prime p, (3) thickid $(C\eta_{(p)}) \subsetneq$ thickid $(S_{(p)}^0) = (\mathcal{SH}(k)_f)_{(p)}$ if $k \subseteq \mathbb{R}$ and p is any prime or $k \subseteq \mathbb{C}$ and p = 2.
- (4) For any prime p, the thick ideals thickid $(C\eta_{(p)}) \cap R_k^{-1}(\mathcal{C}_n)$ are distinct for different $n \geq 0$ and in particular nonzero if $n < \infty$.

This proves that $\mathcal{SH}(k)_f$, $k \subseteq \mathbb{C}$, really has "more" thick ideals than its topological counterpart. In Section 7.2, we compare our results to Balmer's work on prime ideals [Bal10]. For the categories \mathcal{SH}^{fin} , $\mathcal{SH}(\mathbb{Z}/2)_f$, $\mathcal{SH}(\mathbb{C})_f$ and $\mathcal{SH}(\mathbb{R})_f$, we recover the information on prime ideals given in [Bal10, Section 10] from a different point of view.

In Chapter 8, we study two preimages under $R_{\mathbb{C}}$ of a type-*n* spectrum $X_n \in \mathcal{SH}_{(p)}^{fin}$. One of them is $c_{\mathbb{C}}(X_n)$ and the other one, \mathbb{X}_n , is constructed by a motivic version of the construction of X_n , as given in [Rav92, Appendix C]. In analogy to Mitchell's result [Mit85, Theorem 4.8], we prove the following vanishing theorem for motivic Morava K-theory.

Theorem 8.4.1. (Vanishing criterion)

Let s > 0 and $X \in \mathcal{SH}(\mathbb{C})^{fin}$ be a finite motivic cell spectrum such that $\mathrm{H}^{**}(X, \mathbb{Z}/p)$ is free over the exterior algebra $\Lambda_{H\mathbb{Z}/p^{**}}(Q_s)$ as a module over the motivic Steenrod algebra. Then $AK(s)_{**}X = 0$.

This is proven with the help of the motivic Adams spectral sequence for $Ak(s) \wedge X$, where Ak(s) is the motivic analog of the connective Morava K-theory spectrum. This spectral sequence is studied in Section 8.3. In Section 8.5, we construct a spectrum X_n satisfying the assumption of the theorem, and we show that this spectrum is indeed of motivic type n (Theorem 8.5.12). In Section 8.7, we show that $c_{\mathbb{C}}(X_n)$ is also of motivic type n, proving that the two thick ideals in $(\mathcal{SH}(\mathbb{C})_f)_{(p)}$ generated by $c_{\mathbb{C}}(X_n)$ respectively X_n cannot be distinguished by the motivic Morava K-theories. However, we do not know whether the ideals are actually equal or not.

Chapter 9 is devoted to the study of the Bousfield classes of AK(n) and related motivic spectra. The main goal in writing this chapter was to prove that $AK(n+1)_{**}(X) =$ 0 implies $AK(n)_{**}(X) = 0$, which we show for $X \in \mathcal{SH}(\mathbb{C})^{fin}$ and p > 2 in Theorem 9.6.4.

Theorem 9.6.4. Let p > 2. If $X \in \mathcal{SH}(\mathbb{C})^{fin}$ satisfies $AK(n+1)_{**}(X) = 0$, then it also satisfies $AK(n)_{**}(X) = 0$.

A lot of results in Chapter 9 hold more generally. In Section 9.1, we prove that v_n -torsion in $ABP_{**}ABP$ is also v_{n-1} -torsion. This holds in any $\mathcal{SH}(k), k \subseteq \mathbb{C}$ (Theorem 9.1.1). The proof uses methods similar to the topological version of the statement, [JY80, Theorem 0.1]. Another ingredient is the map of Hopf algebroids $(BP_*, BP_*BP) \rightarrow (ABP_{**}, ABP_{**}ABP)$, as studied in [NSØ09]. In Section 9.4, we construct certain operations on AP(n) in $\mathcal{SH}(k)$, $k \subseteq \mathbb{C}$ (Theorem 9.4.12), similar to the operations on P(n) constructed by Würgler in [Wür76, Theorem 5.1]. These are used to prove the equality of Bousfield classes $\langle AK(n) \rangle = \langle AB(n) \rangle$ in $\mathcal{SH}(\mathbb{C})$ (Corollary 9.4.18) with methods similar to those of [JW75]. In the proof of Corollary 9.4.18, we assume $k = \mathbb{C}$ because we make use of the explicitly known coefficient rings $H\mathbb{Z}/p_{**}$ and $AK(n)_{**}$. The result is used to prove Theorem 9.5.1, which is the following decomposition of Bousfield classes in $\mathcal{SH}(\mathbb{C})$, as conjectured in [Hor06, Question 2.17] for arbitrary fields.

Theorem 9.5.1. For p > 2,

$$\langle AE(n) \rangle = \bigvee_{i \le n} \langle AK(i) \rangle$$
 in $\mathcal{SH}(\mathbb{C})$.

Acknowledgements.

The current preprint is my PhD thesis. I want to thank my advisor Prof. Jens Hornbostel for introducing me to the exciting topic of motivic homotopy theory and for offering me this project. I am grateful for his steady support during my dissertation and for all the helpful discussions.

I thank Neil Strickland for sharing his preprint on thick ideals in equivariant stable homotopy categories with me and for allowing me to incorporate parts of it as Chapter 3 of this thesis.

I also owe great thanks to my colleagues Marcus Zibrowius and Jeremiah Heller, who carefully read earlier versions of this work and helped me a lot with their comments. Furthermore, I thank Marc Hoyois for answering some of my questions and I thank Kyle Ormsby for pointing out the reference [Bal10] to me.

This work was financed by the Deutsche Forschungsgemeinschaft, grant no. HO 4729/1-1.

Chapter 2

Thick ideals in classical stable homotopy theory

In this chapter, we introduce basic notation concerning thick ideals and the stable homotopy category SH. In SH^{fin} , a thick ideal is the same as a thick subcategory (Lemma 2.0.8). We recall the thick subcategory theorem of Hopkins and Smith in Theorem 2.0.10.

Definition 2.0.1. A tensor triangulated category is a triple (\mathcal{T}, \wedge, S) consisting of a triangulated category \mathcal{T} and a symmetric monoidal product \wedge on \mathcal{T} with unit S, such that for any $A \in \mathcal{T}, A \wedge -$ preserves exact triangles (see e.g. [Bal05, Definition 1.1]).

Example 2.0.2. The stable homotopy category $(\mathcal{SH}, \wedge, S)$ with $S = S^0 = \Sigma^{\infty} S^0$ is a tensor triangulated category.

Definition 2.0.3. Let (\mathcal{T}, \wedge, S) be a tensor triangulated category. A full triangulated subcategory $\emptyset \neq \mathcal{C} \subseteq \mathcal{T}$ is called a

- (1) thick subcategory if it is closed under retracts.
- (2) thick ideal if it is a thick subcategory and in addition satisfies:

if $X \in \mathcal{T}$ and $Y \in \mathcal{C}$ then $X \wedge Y \in \mathcal{C}$.

If \mathcal{X} is an object or a set of objects, we denote the smallest thick ideal containing \mathcal{X} by thickid(\mathcal{X}) and call it the thick ideal generated by \mathcal{X} . This is well defined because the intersection of thick ideals is again a thick ideal. If \mathcal{X} is a finite set of objects, thickid(\mathcal{X}) is called finitely generated.

Remark 2.0.4. Any finitely generated thick ideal is generated by a single element, namely the direct sum of all generators.

Example 2.0.5. If (\mathcal{T}, \wedge, S) is a tensor triangulated category, then

$$\operatorname{thickid}(S) = \mathcal{T},$$

since for any $X \in \mathcal{T}, X \cong X \wedge S$.

More generally, if Z is in the Picard group $Pic(\mathcal{T})$, i.e., if there exists a Z' such that $Z' \wedge Z \cong S$, then thickid $(Z) = \mathcal{T}$. The Picard group of the stable homotopy category \mathcal{SH} consists precisely of the spheres $\Sigma^n S^0$, $n \in \mathbb{Z}$ (see e.g. [HMS94]), the Picard group of the equivariant stable homotopy category $\mathcal{SH}(G)$ is described in [FJM01] and examples for elements in the Picard groups of motivic stable homotopy categories are given in [Hu05].

Definition 2.0.6. The category \mathcal{SH}^{fin} is the smallest full subcategory of \mathcal{SH} that contains all finite desuspensions of suspension spectra of finite CW complexes and is closed under isomorphisms.

Remark 2.0.7. $S\mathcal{H}^{fin}$ is a tensor triangulated subcategory of $S\mathcal{H}$. It can equivalently be defined as the smallest thick subcategory of $S\mathcal{H}$ that contains S^0 , or as the full subcategory of compact objects in $S\mathcal{H}$ (see, e.g. [Sch12, Theorem II.7.4]).

Lemma 2.0.8. In SH^{fin} , any thick subcategory is already a thick ideal.

Proof. Let X be an element of the thick subcategory $\mathcal{C} \subseteq S\mathcal{H}^{fin}$ and let $Y \in S\mathcal{H}^{fin}$. By the definition of $S\mathcal{H}^{fin}$, there is a finite sequence of spectra $\{Y^k\}_{0 \leq k \leq n}$ such that $Y^0 = S^{n_0}, Y \cong Y^n$ and each Y^k is the cofiber of some map $S^{n_k} \to Y^{k-1}, n_k \in \mathbb{Z}$. Any thick subcategory is closed under suspensions and desuspensions because $\Sigma^{\pm 1}X$ lies in an exact triangle with $X \xrightarrow{1} X$. Hence, $X \wedge Y^0 \in \mathcal{C}$. Assume that $X \wedge Y^k \in \mathcal{C}$ for some k. Then $X \wedge S^{n_{k+1}} \to X \wedge Y^k \to X \wedge Y^{k+1}$ is an exact triangle whose first two objects are in \mathcal{C} . Since \mathcal{C} is a thick subcategory, it follows that $X \wedge Y^{k+1} \in \mathcal{C}$, too, and inductively, $X \wedge Y^n \in \mathcal{C}$. Note further that thick subcategories are closed under isomorphisms as these are special cases of retractions. Hence, $X \wedge Y \in \mathcal{C}$, which proves that \mathcal{C} is a thick ideal.

Definition 2.0.9. Let p be a prime number. The p-local categories $\mathcal{SH}_{(p)}$ and $\mathcal{SH}_{(p)}^{fin}$ are defined as the Bousfield localisations of \mathcal{SH} and \mathcal{SH}^{fin} at the p-local Moore spectrum $M\mathbb{Z}_{(p)}$.

It is a common procedure to study spectra *p*-locally for each prime *p*, i.e. instead of $X \in S\mathcal{H}$ one studies its image $X_{(p)}$ under $S\mathcal{H} \to S\mathcal{H}_{(p)}$, and then fits the information together. For example, *n*-th Morava K-theory K(n) is defined for any fixed prime *p*, where it satisfies $K(n)_*(X) = K(n)_*(X_{(p)})$. For the construction and properties of K(n), see e.g. [JW75].

Now we are ready to state the thick subcategory theorem of Hopkins and Smith [HS98, Theorem 7], which was the main motivation for this thesis. It gives a beautiful and complete description of the thick ideals in $\mathcal{SH}_{(p)}^{fin}$ in terms of Morava K-theories.

Theorem 2.0.10. (Hopkins, Smith)

In $\mathcal{SH}_{(p)}^{fin}$, the thick ideals are given as a chain

$$\mathcal{SH}_{(p)}^{fin} = \mathcal{C}_0 \supseteq \mathcal{C}_1 \supseteq \cdots \supseteq \mathcal{C}_n \supseteq \cdots \supseteq \mathcal{C}_{\infty} = \{0\},\$$

with $C_n = \{ X \in S\mathcal{H}_{(p)}^{fin} \mid K(n-1)_*(X) = 0 \}$ for $0 < n < \infty$.

Definition 2.0.11. A spectrum $X \in S\mathcal{H}_{(p)}^{fin}$ is said to be of type n if $K(n-1)_*(X) = 0$ and $K(n)_*(X) \neq 0$. We write type(X) = n.

For any fixed prime p, the type of a spectrum is well-defined by [Rav84, Theorem 2.11]. The thick subcategory theorem implies that any spectrum X of type n generates C_n as a thick ideal.

Chapter 3

Thick ideals in equivariant stable homotopy theory

The contents of this chapter (except for the introductory section and some details) are due to Neil Strickland [Str10]. We state Strickland's main results on thick ideals in $\mathcal{SH}(G)_f$ and their proofs. We start with the necessary recollection from equivariant stable homotopy theory. Thick ideals in $\mathcal{SH}(G)_f$ are classified by equivariant Morava K-theories, $K(n, H) = G/H_+ \wedge \tilde{E}[\not\geq H] \wedge K(n), H \subseteq G$, which are introduced in Section 3.2. They are related to the classical Morava K-theories via the geometric fixed point functor (Proposition 3.2.1) and satisfy similar properties, such as the Künneth formula (Corollary 3.2.2). As in the non-equivariant theory of Hopkins and Smith, the equivariant Morava K-theories detect nilpotence in $\mathcal{SH}(G)_f$ (Theorem 3.4.2). The general relation between the detection of nilpotence by a family of homology theories and thick ideals is described in Theorem 3.3.12. As a corollary, we reformulate the thick subcategory theorem [HS98, Theorem 7] in a non-*p*-localised way (Theorem 3.4.5) and prove a similar equivariant result, Theorem 3.4.6, which describes an injective lattice homomorphism from the set of thick ideals in $\mathcal{SH}(G)_f$ to a particular lattice GQ and gives a lower bound for its image. An upper bound is given in Proposition 3.4.7.

For our study of thick ideals in the motivic stable homotopy categories $\mathcal{SH}(k)_f$, $k \subseteq \mathbb{R}$, we will use the here given knowledge concerning thick ideals in the $\mathbb{Z}/2$ -equivariant stable homotopy category. Therefore, the case $G = \mathbb{Z}/2$ is the interesting one for the rest of this thesis and we will summarise all results on thick ideals in $\mathcal{SH}(\mathbb{Z}/2)_f$ in Section 3.5. Any thick ideal in $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ is of the form $\mathcal{C}_{m,n} = \{X \mid \phi^{\{1\}}(X) \in \mathcal{C}_m \text{ and } \phi^{\mathbb{Z}/2}(X) \in \mathcal{C}_n\}$, where $m, n \in [0, \infty]$ and ϕ^H is the geometric fixed point functor (Corollary 3.5.1). But not all $\mathcal{C}_{m,n}$ are different. Corollary 3.5.3 gives partial information on which ones are.

3.1 Equivariant stable homotopy theory

Let G be a finite group and $\mathcal{SH}(G)$ be the stable homotopy category of genuine G-spectra. This category has quite a few models. We switch between spectra of G-CW

complexes and spectra of G-simplicial sets, depending on which is more convenient in the concrete situation. A good model for $\mathcal{SH}(G)$ as a tensor triangulated category is the category of orthogonal G-spectra, see e.g. [MM02] or [Sch13]. In Section 4.2, we make use of two other models, namely symmetric G-spectra and $G\Sigma_G$ -spectra. The following definition of finite G-spectra, for example, makes sense if we use the model of orthogonal G-spectra with G-CW complexes as the underlying category of spaces. The definition induces a notion of finiteness for any other model for $\mathcal{SH}(G)$.

Definition 3.1.1. For G a finite group, let $\mathcal{SH}(G)^{fin}$ be the smallest full subcategory of $\mathcal{SH}(G)$ that contains all finite desuspensions of suspension spectra of finite G-CW complexes and is closed under isomorphisms. The objects of $\mathcal{SH}(G)^{fin}$ are called finite G-CW spectra. We denote the closure of $\mathcal{SH}(G)^{fin}$ under retracts in $\mathcal{SH}(G)$ by $\mathcal{SH}(G)_f$.

Both $\mathcal{SH}(G)^{fin}$ and $\mathcal{SH}(G)_f$ are tensor triangulated subcategorties of $\mathcal{SH}(G)$ because finite G-CW complexes are closed under cofiber sequences and under smash products and because retracts commute with smash products.

Definition 3.1.2. A spectrum $X \in SH(G)$ is called dualisable, if the canonical map

$$F(X, S^0) \land Y \to F(X, Y)$$

is an isomorphism for any $Y \in S\mathcal{H}(G)$, where F(-,-) denotes the derived function spectrum and $S^0 = S_G^0$ is the shere spectrum in $S\mathcal{H}(G)$ (for possible definitions of S^0 and F(-,-), see e.g. [Sch13, Examples 2.10 and 5.12]). $DX = F(X, S^0)$ is called the dual of X. It satisfies $DDX \cong X$ ([LMS86, Proposition III.1.3]). In the following, we also use the notation S for S_G^0 , since it is the unit in $S\mathcal{H}(G)$.

 $X \in \mathcal{SH}(G)$ is called compact if $[X, -]_{\mathcal{SH}(G)} = \operatorname{Hom}_{\mathcal{SH}(G)}(X, -)$ preserves arbitrary coproducts.

Proposition 3.1.3. The subcategory $SH(G)_f \subseteq SH(G)$ has the following equivalent descriptions:

- (1) It is the full subcategory of retracts of finite G-CW spectra.
- (2) It is the full subcategory of dualisable objects.
- (3) It is the full subcategory of compact objects.

Proof. (1) and (2) are equivalent by [May96, Theorem XVI.7.4].

Furthermore, any dualisable object X is also compact, because

$$\begin{bmatrix} X, \bigvee Y_i \end{bmatrix} = \begin{bmatrix} S, F(X, \bigvee Y_i) \end{bmatrix} = \begin{bmatrix} S, DX \land \bigvee Y_i \end{bmatrix} = \begin{bmatrix} S, \bigvee (DX \land Y_i) \end{bmatrix}$$
$$= \bigoplus \begin{bmatrix} S, DX \land Y_i \end{bmatrix} = \bigoplus \begin{bmatrix} S, F(X, Y_i) \end{bmatrix} = \bigoplus \begin{bmatrix} X, Y_i \end{bmatrix},$$

where we used that F(X, -) is right adjoint to $X \wedge -$ and that the unit S is compact, see e.g. [Sch13, Corollary 3.30(i)]. Since $\pi_n^H X = 0$ for all $H \subseteq G$ and $n \in \mathbb{Z}$ implies $X \cong 0$ in $\mathcal{SH}(G)$ by the definition of $\mathcal{SH}(G)$, $\{\Sigma^n \Sigma^\infty (G/H)_+ \mid H \subseteq G, n \in \mathbb{Z}\}$ is a detecting set and, hence, also a generating set by [MS06, Lemma 13.1.6]. That is, the smallest thick subcategory of $\mathcal{SH}(G)$ which is closed under infinite coproducts and contains $\{\Sigma^n \Sigma^\infty (G/H)_+ \mid H \subseteq G, n \in \mathbb{Z}\}$ is $\mathcal{SH}(G)$ itself. By [LMS86, Corollary II.6.3], $\Sigma^\infty (G/H)_+$ is dualisable and, thus, compact. Hence, $\{\Sigma^\infty (G/H)_+ \mid H \subseteq G\}$ is a set of compact generators for $\mathcal{SH}(G)$. By general theory due to Neeman [Nee96], see e.g. [MS06, Theorem 13.1.14], the full subcategory of compact objects in $\mathcal{SH}(G)$ is the thick subcategory generated by this set (i.e., the smallest thick subcategory of $\mathcal{SH}(G)$ containing this set). Therefore, (3) is also equivalent to (1) and (2).

Let G be a finite group and $H \subseteq G$ a subgroup. There are functors $i : SH \to SH(G)$ and $\phi^H : SH(G) \to SH$, where i maps a nonequivariant spectrum to the corresponding G-spectrum with trivial G-action and ϕ^H is the geometric fixed point functor (as defined in [LMS86, Definition 9.7], [MM02, Definition 4.3] or [Sch13, Section 7.3]) concatenated with the forgetful functor from SH(W(H)) to SH, where W(H) denotes the Weyl group of $H \subseteq G$. We will need the following properties [Str10, Proposition 12.1 and Theorem 12.4].

Proposition 3.1.4. The geometric fixed point functor ϕ^H has the following properties: (1) In SH, $\phi^H(\Sigma^{\infty}X) = \Sigma^{\infty}X^H$ for any suspension spectrum $\Sigma^{\infty}X \in SH(G)$. (2) In SH, $\phi^H(X \wedge Y) = \phi^H(X) \wedge \phi^H(Y)$ for any $X, Y \in SH(G)$. (3) In SH, $\phi^H(i(X)) = X$ for any $X \in SH$.

(4) If $\phi^H(X) = 0$ in SH for all $H \subseteq G$, then X = 0 in SH(G).

Proof. A proof of (1) can be found in [LMS86, Corollary II.9.9], [MM02, Corollary 4.6], or in [Sch13, Example 7.7]. (2) follows from [LMS86, Theorem II.9.8(ii)] and [LMS86, Proposition II.9.12(ii)]. (3) follows directly from the definition of ϕ^H , since H acts trivially on i(X). (4) is proven in [Sch13, Theorem 7.12] and in [Str10, Theorem 12.4].

3.2 Equivariant Morava K-theories

For $H \subseteq G$, $\tilde{E}[\not\geq H]$ denotes a G-space which satisfies:

$$\tilde{E}[\not\geq H]^K \simeq \begin{cases} 0 & \text{if } K \not\geq_G H \\ S^0 & \text{if } K \geq_G H \end{cases}$$

where $K \geq_G H$ means that K contains a subgroup conjugate to H.

The existance of such a space $E[\not\geq H]$ follows from the theory of classifying spaces for families (see e.g. [LMS86, Section II.2]), if one takes \mathcal{F} as the family of all subgroups of G for which H is not subconjugate and then defines $\tilde{E}[\not\geq H]$ by the cofiber sequence

$$E\mathcal{F}_+ \to S^0 \to \tilde{E}[\not\geq H],$$

as in [MM02, Notations 4.14].

Fix a prime number p. Strickland [Str10, Definition 16.2] defines Morava K-theory spectra in $S\mathcal{H}(G)$, G a finite group, by

$$K(n,H) = G/H_+ \wedge E[\geq H] \wedge K(n)$$

for any subgroup $H \subseteq G$. He notes that, as a localisation of S^0 , $\tilde{E}[\not\geq H]$ is a commutative ring spectrum, which together with the ring structure of K(n) and the diagonal map on G/H induces a ring structure on K(n, H), which is commutative for p > 2. We will only be interested in H up to conjugacy, because if H and H' are conjugate, then $K(n, H) \cong K(n, H')$.

The following proposition serves as motivation for this definition of equivariant Morava K-theories [Str10, Remark 16.4 ff].

Proposition 3.2.1.

$$K(n, H)_{*}(X) = K(n)_{*}(\phi^{H}(X))$$

and

$$K(n, H)^{*}(X) = K(n)^{*}(\phi^{H}(X)).$$

Proof. Here, we need the following formula for the geometric fixed point spectrum [LMS86, Theorem II.9.8(ii)]:

$$\phi^H(X) \cong (\tilde{E}[\not\geq H] \wedge X)^H,$$

where $(-)^H$ is the spectrification of the levelwise fixed point functor. In the following, we abbreviate $\tilde{E}[\not\geq H]$ by \tilde{E} . The first equation follows from

$$K(n,H)_*(X) = \pi_* \left((X \wedge G/H_+ \wedge \tilde{E} \wedge K(n))^G \right)$$
$$= \pi_* \left((X \wedge \tilde{E} \wedge K(n))^H \right) = \pi_* \left(\phi^H(X \wedge K(n)) \right)$$
$$= \pi_* \left(\phi^H(X) \wedge K(n) \right) = K(n)_*(\phi^H(X)).$$

For the second equation, we use the fact that G/H_+ is self-dual [LMS86, Corollary II.6.3], hence

$$K(n, H)^{*}(X) = [X, F(G/H_{+}, \tilde{E} \wedge K(n))]_{*}^{G} = [X, \tilde{E} \wedge K(n)]_{*}^{H}.$$

We claim that this is isomorphic to $[\phi^H X, \phi^H K(n)]_* = K(n)^*(\phi^H X)$. To prove the claim, first note that because $\tilde{E} \wedge -$ is a Bousfield localisation functor,

$$[X, \tilde{E} \wedge K(n)]^H_* = [\tilde{E} \wedge X, \tilde{E} \wedge K(n)]^H_*.$$

From here, the H-fixed points yield a map

$$\alpha: [\tilde{E} \wedge X, \tilde{E} \wedge K(n)]^H_* \to [(\tilde{E} \wedge X)^H, (\tilde{E} \wedge K(n))^H]_*$$

and the latter group is isomorphic to $[\phi^H X, \phi^H K(n)]_*$. Assume that X^H is an orbit H/K_+ for $K \subseteq H$. If $K \neq H$, $\alpha : 0 \to 0$ is an isomorphism. If K = H, $[\tilde{E} \land X, \tilde{E} \land K(n)]^H_* = [S^0_G, \tilde{E} \land K(n)]^H_* = [S^0, \phi^H(K(n))]_* = [\phi^H X, \phi^H(K(n))]_*$. That is, α is an isomorphism on all orbit types and it follows that α is an isomorphism for any $X \in \mathcal{SH}(G)$.

From this and the properties of nonequivariant Morava K-theories (see e.g. [HS98, Section 1]), it follows immediately that K(n, H) has coefficients like K(n) and satisfies the Künneth formula [Str10, Section 16].

Corollary 3.2.2. The equivariant Morava K-theories satisfy the following properties: (1) $K(n, H)_*S_G^0 = K(n)_*S^0 = \mathbb{F}_p[v_n^{\pm 1}]$ for any n > 0 and any $H \subseteq G$.

(2) $K(n,H)_*(X \wedge Y) \cong K(n,H)_*(X) \otimes_{K(n)_*} K(n,H)_*(Y)$ for any $X, Y \in \mathcal{SH}(G)$. (3) If X is dualisable, i.e., if $X \in \mathcal{SH}(G)_f$, then

 $K(n,H)_*(DX) \cong \operatorname{Hom}_{K(n)_*}(K(n,H)_*(X), K(n)_*).$

Furthermore, Strickland shows [Str10, Proposition 16.6]:

Proposition 3.2.3. If $p \neq p'$ or $n \neq n'$ or $H \neq_G H'$ (i.e., not conjugate in G), then

 $K(p, n, H) \wedge K(p', n', H') = 0.$

Proof. In the cases $p \neq p'$ and $n \neq n'$, this follows from $K(p,n) \wedge K(p',n') = 0$ (see [Rav84, Theorem 2.1(i)]), as these appear as smash factors in $K(p,n,H) \wedge K(p',n',H')$. Therefore, assume p = p', n = n' and $H \neq_G H'$. Now it suffices to show $G/H_+ \wedge \tilde{E}[\geq H] \wedge G/H'_+ \wedge \tilde{E}[\geq H'] = 0$, which is easily checked on the level of K-fixed points for all $K \subseteq G$.

3.3 Nilpotence and lattices of thick ideals

For a convenient description of the collection of thick ideals in the equivariant homotopy category of dualisable spectra, $SH(G)_f$, [Str10] uses the language of lattices.

Definition 3.3.1. A lattice is a partially ordered set A for which any finite subset $F \subseteq A$ has a greatest lower bound (called meet) $\bigwedge F$ and a smallest upper bound (called join) $\bigvee F$. The largest element in A is $\bigwedge \emptyset$, which we denote by 1 and the smallest element is $0 = \bigvee \emptyset$. A lattice homomorphism is an order preserving map $f : A \to B$ which also preserves all joins and meets.

- **Example 3.3.2.** (1) The collection of thick ideals C in a tensor triangulated category \mathcal{T} , partially ordered by inclusion, is a lattice. Meets are just intersections, whereas the join of a finite collection of thick ideals is the smallest thick ideal which contains all objects of the different ideals. We denote this lattice by $\mathrm{Idl}(\mathcal{T})$.
- (2) The power set of any set is a lattice, partially ordered by inclusion, meets given by intersections and joins by unions.

We introduce a new notation, due to Strickland, which will be useful in the rest of this chapter.

Notation 3.3.3. For a prime p and a nonnegative integer n, let $K(p^{-n})$ denote the n-th Morava K-theory spectrum at the prime p (which above was denoted by K(p, n) or just K(n)).

One advantage of this notation is that there is only one name for the zeroth Morava K-theory spectrum (which is independent of p): $K(1) = H\mathbb{Q}$.

Definition 3.3.4. Let

$$\mathcal{Q}_p = \{ p^{-n} \mid 0 \le n \le \infty \}$$

and

$$\mathcal{Q} = \{ u \in \prod_{p} \mathcal{Q}_{p} \mid u_{p} = 1 \,\forall \, p \text{ or } u_{p} \neq 1 \,\forall \, p \}.$$

The sets Q_p and Q are lattices, with the usual ordering of rational numbers and the componentwise partial ordering of products. We immediately see that Theorem 2.0.10 can be reformulated as follows.

Corollary 3.3.5. Let $Idl(\mathcal{SH}^{fin}_{(p)})$ be as in Example 3.3.2(1). The map

$$au_p: \mathrm{Idl}(\mathcal{SH}_{(p)}^{fin}) \longrightarrow \mathcal{Q}_p,$$

 $\tau_p(\mathcal{C}) = \max\{p^{-n} \mid \operatorname{type}(X) = n \text{ for some } X \in \mathcal{C}\},\$

is a lattice isomorphism.

We will see in Theorem 3.4.5 how to merge the information for different p to a classification of finitely generated thick ideals of \mathcal{SH}^{fin} using the lattice Q. But before we are able to do so, we need some more theory on thick ideals and lattice homomorphisms.

Definition 3.3.6. For $X \in \mathcal{SH}(G)_f$, let $\operatorname{ann}(X)$ denote the fibre of the unit map $S \to F(X, X)$ and define

$$\mathcal{A}_{\operatorname{ann}(X)} = \{A \mid \operatorname{ann}(X)^{\wedge n} \land A \to A \text{ is null for some } n > 0\}.$$

The map here is the *n*-fold smash product of the map $\operatorname{ann}(X) \to S$ from the cofiber sequence, smashed with $A \xrightarrow{1} A$.

The following is [Str10, Proposition 15.6].

Proposition 3.3.7. The smallest thick ideal containing X is

$$\operatorname{thickid}(X) = \mathcal{A}_{\operatorname{ann}(X)}$$

and thickid(X) \subseteq thickid(Y) if and only if the map $\operatorname{ann}(Y)^{\wedge n} \to S$ factors through $\operatorname{ann}(X) \to S$ for some n > 0.

Proof. We first show that $\operatorname{thickid}(X) \subseteq \mathcal{A}_{\operatorname{ann}(X)}$. Since X is a module over $F(X, X) = DX \wedge X$, it is a retract of $DX \wedge X \wedge X$. It follows that $\operatorname{ann}(X) \wedge X$ is the fiber of a map $X \to DX \wedge X \wedge X$ which splits, so $\operatorname{ann}(X) \wedge X \to X$ is zero and hence $X \in \mathcal{A}_{\operatorname{ann}(X)}$. It is easy to see that $\mathcal{A}_{\operatorname{ann}(X)}$ is closed under exact triangles and retracts, as well as under smashing with arbitrary objects. Hence, $\mathcal{A}_{\operatorname{ann}(X)}$ is a thick ideal containing X, which proves $\operatorname{thickid}(X) \subseteq \mathcal{A}_{\operatorname{ann}(X)}$.

Now assume $A \in \mathcal{A}_{\operatorname{ann}(X)}$. We need to show $A \in \operatorname{thickid}(X)$. Consider the cofiber sequence

$$\operatorname{ann}(X)^{\wedge n} \wedge A \to A \to S/(\operatorname{ann}(X)^{\wedge n}) \wedge A.$$

By the assumption, we can choose n such that the first map is zero. Then it follows that A is a retract of $S/(\operatorname{ann}(X)^{\wedge n}) \wedge A$. Therefore, it suffices to show $S/(\operatorname{ann}(X)^{\wedge n}) \in \operatorname{thickid}(X)$. By the definition of $\operatorname{ann}(X)$, $S/\operatorname{ann}(X) = F(X, X) = DX \wedge X \in \operatorname{thickid}(X)$. There is a cofiber sequence

$$\operatorname{ann}(X) \wedge S/(\operatorname{ann}(X)^{\wedge j}) \to S/(\operatorname{ann}(X)^{\wedge j+1}) \to S/(\operatorname{ann}(X)^{\wedge j}),$$

which implies inductively that $S/(\operatorname{ann}(X)^{\wedge n}) \in \operatorname{thickid}(X)$.

The existence of this cofiber sequence follows from Verdier's axiom for triangulated categories, also known as octahedral axiom (see e.g. [Nee01, Proposition 1.4.6]). Applied to the three cofiber sequences $I \wedge J \rightarrow I \wedge S \rightarrow I \wedge S/J$, as well as $I \wedge J \rightarrow S \rightarrow S/(I \wedge J)$ and $I \rightarrow S \rightarrow S/I$, the axiom yields a cofiber sequence $I \wedge S/J \rightarrow S/(I \wedge J) \rightarrow S/I$.

For the second claim, assume that $\operatorname{thickid}(X) \subseteq \operatorname{thickid}(Y)$, which is equivalent to $X \in \operatorname{thickid}(Y) = \mathcal{A}_{\operatorname{ann}(Y)}$. Let n > 0 be such that $\operatorname{ann}(Y)^{\wedge n} \wedge X \to X$ is the zero map. Consider the two cofiber sequences

$$\operatorname{ann}(Y)^{\wedge n} \longrightarrow S \longrightarrow S/(\operatorname{ann}(Y)^{\wedge n})$$
$$\|$$
$$\operatorname{ann}(X) \longrightarrow S \longrightarrow F(X, X).$$

The smash product of the upper sequence with X is

$$\operatorname{ann}(Y)^{\wedge n} \wedge X \xrightarrow{0} X \to S/(\operatorname{ann}(Y)^{\wedge n}) \wedge X,$$

so there is a retraction $S/(\operatorname{ann}(Y)^{\wedge n}) \wedge X \to X$, which then induces a morphism $S/(\operatorname{ann}(Y)^{\wedge n}) \to F(X, X)$ making the diagram commutative. From the axioms for triangulated categories it follows that we can fill in the required map $\operatorname{ann}(Y)^{\wedge n} \to \operatorname{ann}(X)$, as claimed.

On the other hand, if $\operatorname{ann}(Y)^{\wedge n} \to S$ factors through $\operatorname{ann}(X) \to S$ and $\operatorname{ann}(X)^{\wedge m} \wedge A \to A$ is zero then also $\operatorname{ann}(Y)^{\wedge (nm)} \wedge A \to A$ is zero and it follows $\mathcal{A}_{\operatorname{ann}(X)} \subseteq \mathcal{A}_{\operatorname{ann}(Y)}$.

Proposition 3.3.8. Let \mathcal{I}_0 and \mathcal{J}_0 be collections of objects in $S\mathcal{H}(G)_f$ and let \mathcal{I} and \mathcal{J} be the thick ideals which they generate. Then

$$\mathcal{I} \cap \mathcal{J} = \text{thickid} \left(\{ Y \land Z \mid Y \in \mathcal{I}_0, Z \in \mathcal{J}_0 \} \right).$$

Proof. This is [Str10, Proposition 15.8]. Let $\mathcal{K} = \text{thickid}(\{Y \land Z \mid Y \in \mathcal{I}_0, Z \in \mathcal{J}_0\})$. The intersection $\mathcal{I} \cap \mathcal{J}$ is a thick ideal which contains $Y \land Z$ for all $Y \in \mathcal{I}_0$ and $Z \in \mathcal{J}_0$. Therefore, $\mathcal{K} \subseteq \mathcal{I} \cap \mathcal{J}$. Now, let

$$\mathcal{I}' = \{ Y \in \mathcal{I} \mid Y \land Z \in \mathcal{K} \; \forall \, Z \in \mathcal{J}_0 \},\$$

$$\mathcal{J}' = \{ Z \in \mathcal{J} \mid Y \land Z \in \mathcal{K} \; \forall \, Y \in \mathcal{I} \}.$$

It is easy to check that \mathcal{I}' is a thick subideal of \mathcal{I} which contains \mathcal{I}_0 . Hence, $\mathcal{I}' = \mathcal{I}$. It follows that the thick subideal $\mathcal{J}' \subseteq \mathcal{J}$ contains \mathcal{J}_0 , so $\mathcal{J}' = \mathcal{J}$. That is, $Y \wedge Z \in \mathcal{K}$ for all $Y \in \mathcal{I}, Z \in \mathcal{J}$.

Now, let $X \in \mathcal{I} \cap \mathcal{J}$. Since X is an F(X, X)-module, X is a retract of $DX \wedge X \wedge X$. Consider $DX \wedge X$ as an object of \mathcal{I} and the other X as an object of \mathcal{J} . It follows $DX \wedge X \wedge X \in \mathcal{K}$ and hence $X \in \mathcal{K}$.

Remark 3.3.9. Note that Proposition 3.3.8 holds in any tensor triangulated category in which internal hom objects exist and all objects are dualisable. The following definition, theorem and corollary can also be formulated in such a general setting, given a suitable notion of homology theories.

Definition 3.3.10. Let $\{E_i \mid i \in I\}$ be a family of ring spectra in $\mathcal{SH}(G)$. For $X \in \mathcal{SH}(G)_f$, define

$$supp(X) = \{i \in I \mid (E_i)_*(X) \neq 0\}.$$

If $\mathcal{C} \subseteq \mathcal{SH}(G)_f$ is a subcategory, let

$$\operatorname{supp}(\mathcal{C}) = \bigcup_{X \in \mathcal{C}} \operatorname{supp}(X).$$

For a map $f: X \to Y$, we also define

$$supp(f) = \{i \in I \mid (E_i)_*(f) \neq 0\}.$$

Remark 3.3.11. If $(E_i)_*(X) = 0$, then $(E_i)_*(Y) = 0$ for any $Y \in \text{thickid}(X)$ by the following arguments. If $A \to B \to C$ is a cofiber sequence and the $(E_i)_*$ -homology of two of the three objects is zero, then, by the long exact $(E_i)_*$ -sequence, $(E_i)_*(-)$ of the third object is zero, too. If $(E_i)_*(A) = \pi_*(E_i \land A) = 0$, then also $(E_i)_*(A \land B) = \pi_*(E_i \land A \land B) = 0$. And, finally, if $(E_i)_*(A) = 0$ and B is a retract of A, then $(E_i)_*(B) \to 0 \to (E_i)_*(B)$ is the identity map, hence, $(E_i)_*(B) = 0$.

This implies

$$\operatorname{supp}(\operatorname{thickid}(X)) = \operatorname{supp}(X).$$

The following theorem is one of the central results in [Str10], where it is Theorem 15.14.

Theorem 3.3.12. (Strickland) Assume $\{E_i \mid i \in I\}$ is a family of ring spectra in SH(G) satisfying the following properties:

- (1) If $f : X \to Y$, with $X, Y \in \mathcal{SH}(G)_f$, and $(E_i)_*(f) = 0$ for all $i \in I$, then there exists n > 0 such that $f^{\wedge n} = 0$.
- (2) For any $X, Y \in \mathcal{SH}(G)_f$ and any $i \in I$, $(E_i)_*(X \wedge Y) \cong (E_i)_*(X) \otimes_{(E_i)_*} (E_i)_*(Y)$.

(3) For any $i \in I$, $(E_i)_* = (E_i)_*(S^0)$ is concentrated in even degrees and any nonzero homogeneous element in $(E_i)_*$ is invertible.

Then, for any $X, Y \in \mathcal{SH}(G)_f$, thickid $(X) \subseteq$ thickid(Y) if and only if supp $(X) \subseteq$ supp(Y).

In other words, a family $\{E_i \mid i \in I\}$ of spectra detecting nilpotence (see Definition 3.3.13) and satisfying some additional properties can be used to distinguish any two different thick ideals with the help of the support functor $\operatorname{supp}(-)$.

Proof. Consider the cofiber sequence $\operatorname{ann}(Y) \xrightarrow{v} S \xrightarrow{u} F(Y,Y)$. We first show that $\operatorname{supp}(v) = I \setminus \operatorname{supp}(Y)$. Consider the long exact sequence

$$\cdots \to (E_i)_*(F(Y,Y)) \to (E_i)_*(\operatorname{ann}(Y)) \xrightarrow{(E_i)_*(v)} (E_i)_*(S) \xrightarrow{(E_i)_*(u)} (E_i)_*(F(Y,Y)) \to \cdots$$

If $i \in I \setminus \operatorname{supp}(Y)$, then $(E_i)_*(F(Y,Y)) \cong (E_i)_*(DY) \otimes_{(E_i)_*} (E_i)_*(Y) = 0$ and $(E_i)_*(v)$ is an isomorphism. Hence, $i \in \operatorname{supp}(v)$. If, on the other hand, $(E_i)_*(v) \neq 0$, it already has to be surjective because $(E_i)_*(\operatorname{ann}(Y))$ is an $(E_i)_*$ -vector space (by property (3)). It follows that $(E_i)_*(u) = 0$. But u is the unit map of F(Y,Y), so this implies $(E_i)_*(F(Y,Y)) = 0$. As Y is a retract of $F(Y,Y) \wedge Y$, it follows that $(E_i)_*(Y) = 0$. This proves $\operatorname{supp}(v) = I \setminus \operatorname{supp}(Y)$.

Now let $X, Y \in \mathcal{SH}(G)_f$ and $\operatorname{supp}(X) \subseteq \operatorname{supp}(Y)$. With v as above, we have $\operatorname{supp}(v) = I \setminus \operatorname{supp}(Y) \subseteq I \setminus \operatorname{supp}(X)$. Hence,

$$\operatorname{supp}\left(\operatorname{ann}(Y) \xrightarrow{v} S \to F(X, X)\right) = \emptyset.$$

By property (1), this map is smash nilpotent, so there is some m > 0 such that

$$\operatorname{ann}(Y)^{\wedge m} \to S \to F(X,X)^{\wedge m}$$

is the zero map. Concatenation defines a map $F(X,X)^{\wedge m} \to F(X,X)$, over which the unit map $S \to F(X,X)$ factors, so we get a diagram in which the lower row is a cofiber sequence, the composition of the two upper maps is zero and the square commutes:

It follows that the map $\operatorname{ann}(Y)^{\wedge m} \to S$ factors over $\operatorname{ann}(X)$. By Proposition 3.3.7, this is equivalent to $\operatorname{thickid}(X) \subseteq \operatorname{thickid}(Y)$.

For the other direction, assume thickid(X) \subseteq thickid(Y). Then by Remark 3.3.11,

$$\operatorname{supp}(X) = \operatorname{supp}(\operatorname{thickid}(X)) \subseteq \operatorname{supp}(\operatorname{thickid}(Y)) = \operatorname{supp}(Y).$$

Definition 3.3.13. We say that a family $\{E_i \mid i \in I\}$ detects nilpotence, if for any $f: X \to Y$ in $\mathcal{SH}(G)_f$, $\operatorname{supp}(f) = \emptyset$ implies $f^{\wedge n} = 0$ for some n > 0.

Corollary 3.3.14. (Strickland) Under the assumptions of the above theorem, the map from the collection of thick ideals in $SH(G)_f$ to the set of subsets of I,

$$\operatorname{Idl}(\mathcal{SH}(G)_f) \longrightarrow \mathcal{P}(I),$$

 $\mathcal{C}\mapsto \mathrm{supp}(\mathcal{C}),$

is a lattice homomorphism (see Definition 3.3.1). It is injective on the collection of finitely generated thick ideals, $\operatorname{FIdl}(\mathcal{SH}(G)_f)$ (see Definition 2.0.3).

Proof. This is [Str10, Corollary 15.15]. It is clear from the definition of $\operatorname{supp}(\mathcal{C})$, that $\operatorname{supp}(-)$ is order preserving. The map preserves meets by Proposition 3.3.8 and by the Künneth formula for E_i . As $\operatorname{supp}(\mathcal{C})$ is the support of any set of generators for \mathcal{C} and the join of thick ideals \mathcal{C}_i is generated by the collection of generators of the individual thick ideals, it is also clear that $\operatorname{supp}(-)$ preserves joins. Recall that any finitely generated thick ideal is already generated by a single element (Remark 2.0.4). By the above theorem, $\operatorname{thickid}(X) = \operatorname{thickid}(Y)$ if and only if $\operatorname{supp}(X) = \operatorname{supp}(Y)$, which is the same as $\operatorname{supp}(\operatorname{thickid}(X)) = \operatorname{supp}(\operatorname{thickid}(Y))$. This proves the injectivity on FIdl.

3.4 Thick ideals and equivariant Morava K-theories

Definition 3.4.1. For a finite group G, let sub(G) denote the set of equivalence classes of conjugate subgroups of G. Let

$$\mathcal{Q}' = \{p^{-n} \mid p \text{ prime}, 0 \le n < \infty\}$$

and $GQ' = \mathcal{Q}' \times \mathrm{sub}(G).$

The following theorem shows that the family of equivariant Morava K-theories $\{K(p^{-n}, H) \mid (p^{-n}, H) \in GQ'\}$ (see Section 3.2) detects nilpotence, as required in the assumptions of Theorem 3.3.12 and Corollary 3.3.14. As in [DHS88, Theorem 1], there are different kinds of nilpotence, which are all detected by the Morava K-theories. Although we mainly work with smash nilpotence, the theorem, which is [Str10, Theorem 16.7], considers all three definitions.

Theorem 3.4.2. (Strickland)

- (1) Let $R \in S\mathcal{H}(G)$ be a ring spectrum. Then $\alpha \in \pi^G_* R$ is nilpotent if and only if for all $v \in GQ'$, $K(v)_*(\alpha)$ is nilpotent as an element of $K(v)_* R$.
- (2) A self-map $f: \Sigma^k W \to W, W \in \mathcal{SH}(G)_f$, is nilpotent if and only if for all $v \in GQ'$, $K(v)_*(f)$ is nilpotent.
- (3) A map $f: W \to X$, with $W \in \mathcal{SH}(G)_f$ and $X \in \mathcal{SH}(G)$, is smash nilpotent if and only if for all $v \in GQ'$, $K(v)_*(f)$ is nilpotent.
- Proof. (1) Let $\alpha: S_G^d \to R$ be such that $K(v)_*(\alpha)$ is nilpotent for all $v \in GQ'$. For any $H \subseteq G, \phi^H(S_G^d)$ is a non-equivariant sphere, so $\phi^H \alpha \in \pi_*(\phi^H R)$. By Proposition 3.2.1, $K(u)_*(\phi^H \alpha) = K(u, H)_*(\alpha)$, so this is nilpotent for all $u \in Q'$. Furthermore, $\phi^H R$ is a ring spectrum and, by [HS98, Theorem 3(i)], it follows that $\phi^H(\alpha)$

is nilpotent, so $(\phi^H R)[\phi^H \alpha^{-1}] = 0$ (for the definition of this mapping telescope, see [DHS88, page 212]). By [Sch13, Remark 7.15], ϕ^H preserves telescopes, hence, $\phi^H(R[\alpha^{-1}]) \cong (\phi^H R)[\phi^H \alpha^{-1}] = 0$. This holds for all H, which by Proposition 3.1.4(4) implies $R[\alpha^{-1}] = 0$, and, hence, $\pi_*^G R[\alpha^{-1}] = 0$. Thus, α is nilpotent.

- (2) The adjoint of f is an element $\alpha \in \pi_d^G F(W, W)$, and $K(v)_*(\alpha)$ is nilpotent for all $v \in GQ'$. So, the claim follows from (1).
- (3) Let $R = \bigvee_{k \ge 0} F(W, X)^{\wedge k} \in \mathcal{SH}(G)$ be the free associative ring spectrum generated by F(W, X). The map f is adjoint to $\alpha \in \pi_d^G F(W, X) \subset \pi_d^G R$ such that $K(v)_*(\alpha)$ is nilpotent for all $v \in GQ'$, and the claim follows from (1).

Definition 3.4.3. Let $GQ = \prod_{\text{sub}(G)} \mathcal{Q}$, where \mathcal{Q} is as in Definition 3.3.4 and let

$$\max: \mathcal{P}(GQ') \longrightarrow GQ, \ I \mapsto \left\{ (H, p) \mapsto \max\{p^{-n} \mid (H, p^{-n}) \in I\} \right\},\$$

with the convention $\max(\emptyset) = 0$. Let

$$\tau = \max \circ \operatorname{supp} : \operatorname{Idl}(\mathcal{SH}(G)_f) \longrightarrow GQ.$$

Corollary 3.4.4. The functor

$$\mathrm{Idl}(\mathcal{SH}(G)_f) \longrightarrow \mathcal{P}(GQ'),$$

$$\mathcal{C} \mapsto \operatorname{supp}(\mathcal{C}) = \left\{ v \in GQ' \mid K(v)_*(X) \neq 0 \text{ for some } X \in \mathcal{C} \right\},\$$

is a lattice homomorphism. Furthermore, it is injective.

Proof. This is [Str10, Corollary 16.8]. It follows from Corollary 3.3.14 applied to the family of equivariant Morava K-theories. The assumptions are satisfied by Theorem 3.4.2(3) and Proposition 3.2.2.

Corollary 3.3.14 states that the restriction of supp to finitely generated thick ideals $\operatorname{FIdl}(\mathcal{SH}(G)_f)$ is injective. To show that supp is injective on arbitrary thick ideals, it suffices to show that τ is injective. Note that τ is injective on $\operatorname{FIdl}(\mathcal{SH}(G)_f)$ because, by [Rav84, Theorem 2.11], $K(p^{-n})_*(\phi^H X) = 0$ implies $K(p^{-m})_*(\phi^H X) = 0$ for all $m \leq n$.

Now let $\tau(\mathcal{I}) = \tau(\mathcal{J})$ for some thick ideals \mathcal{I} and \mathcal{J} in $\mathcal{SH}(G)_f$. Thus, $\tau(\mathcal{I})_{H,p} = \tau(\mathcal{J})_{H,p}$ for all $H \subseteq G$ and all primes p. Assume $X \in \mathcal{I}$. Then $\tau(X)_{H,p} \leq \tau(\mathcal{I})_{H,p} = \tau(\mathcal{J})_{H,p}$. Let $I = \{(H,p) \mid \tau(X)_{H,p} \neq 0, 1\}$. We claim that I is finite. Since G is finite, there are only finitely many possibilities for H. If $(H,p) \in I$, then $K(p^{-0})_*(\phi^H X) = H_*(\phi^H X, \mathbb{Q}) = 0$. Since $\phi^H X$ is a finite spectrum, this implies $H_*(\phi^H X, \mathbb{F}_q) = 0$ for all but finitely many q, so I is finite. For any $u \in I$, there exists by assumption a $Y_u \in \mathcal{J}$ with $\tau(Y_u)_u \geq \tau(X)_u$. Similarly, let $I' = \{H \mid \tau(X)_H = 1\}$ and pick $Y_H \in \mathcal{J}$ with $\tau(Y_H) = 1$ for each $H \in I'$. Then, by the injectivity of τ on finitely generated thick ideals, X is contained in thickid $(\{Y_u \mid u \in I\} \cup \{Y_H \mid H \in I'\})$, which is a finitely generated subideal of \mathcal{J} . It follows $\mathcal{I} \subseteq \mathcal{J}$ and, similarly, $\mathcal{J} \subseteq \mathcal{I}$.

As promised, we now state a reformulation of the thick subcategory theorem from [HS98], which is no longer in *p*-local form. We do not claim that all the notation from above was necessary to state this. But it is helpful for generalising the result to $\mathcal{SH}(G)_f$ or maybe also other categories.

Theorem 3.4.5. The composition

$$\tau: \mathrm{Idl}(\mathcal{SH}^{fin}) \xrightarrow{\mathrm{supp}} \mathcal{P}(\mathcal{Q}') \xrightarrow{\mathrm{max}} \mathcal{Q}$$

is bijective. Its restriction to the finitely generated thick ideals maps $\operatorname{FIdl}(\mathcal{SH}^{fin})$ bijectively to

$$\operatorname{Fin}(\mathcal{Q}) = \{ u \in \mathcal{Q} \mid u = 1 \text{ or } u_p = 0 \text{ for almost all } p \}.$$

Proof. This is [Str10, Proposition 19.14]. The theorem states that supp maps injectively onto its image, which is isomorphic to \mathcal{Q} . The map supp is injective by the nonequivariant version of Corollary 3.4.4. The image of supp maps injectively to \mathcal{Q} because $K(p^{-n})_*X = 0$ implies $K(p^{-m})_*X = 0$ for all $m \leq n$ by [Rav84, Theorem 2.11]. We show that τ is surjective: By [Mit85, Theorem B(b)] or [Rav92, Section C.3], for any number n > 0 there is a spectrum $X_n \in S\mathcal{H}_{(p)}^{fin}$ of type n. Thus, the p-localisation of τ is surjective onto \mathcal{Q}_p (see Definition 3.3.4). Let $u \in \mathcal{Q}$. If u = 1, then $\mathcal{C} = \text{thickid}(S^0)$ satisfies $\tau(\mathcal{C}) = u$. Assume $u \neq 1$, so $u_p = p^{-n_p}$ with $0 < n_p \leq \infty$. Let X_{n_p} be a p-local spectrum of type n_p if $0 < n_p < \infty$ and $X_{n_p} = 0$ if $n_p = \infty$. Then $\mathcal{C} = \text{thickid}(X_{n_2}, X_{n_3}, X_{n_5}, \cdots)$ satisfies $\tau(\mathcal{C}) = u$. Hence, τ is surjective. If u = 1 or $u_p = 0$ for almost all p, then \mathcal{C} as above is finitely generated. As in the proof of Corollary 3.4.4, all finitely generated thick ideals are mapped to Fin(\mathcal{Q}).

This theorem identifies the thick ideals with a sublattice of $\mathcal{P}(\mathcal{Q}')$ that is bijective to \mathcal{Q} . Similarly, thick ideals in the equivariant category $\mathcal{SH}(G)_f$ are mapped injectively into GQ. Unlike its nonequivariant version, the equivariant version of τ is not surjective. However, we can state the following:

Theorem 3.4.6. (Strickland) The composition

$$r : \mathrm{Idl}(\mathcal{SH}(G)_f) \xrightarrow{\mathrm{supp}} \mathcal{P}(GQ') \xrightarrow{\mathrm{max}} GQ$$

is injective. Its image contains all $u \in GQ$ which satisfy: If $H \subseteq H'$ for some $H, H' \in sub(G)$, then $u_H \ge u_{H'}$.

Proof. We already know that supp is an injective lattice homomorphism. The injectivity of τ follows as in the nonequivariant case, by considering *H*-fixed points for each $H \in \text{sub}(G)$ separately.

Now let $u \in GQ$ be such that $H' \subseteq H$ implies $u_{H'} \geq u_H$. Let X_{u_H} be a finite spectrum in $S\mathcal{H}$ with $\tau(X_{u_H}) = u_H$. Recall that any nonequivariant spectrum maps to a *G*-spectrum with trivial *G*-action through the functor $i : S\mathcal{H} \to S\mathcal{H}(G)$. Let $Y_H = i(X_{u_H}) \wedge G/H_+$. This is a finite *G*-spectrum satisfying $\phi^{H'}Y_H = X_{u_H} \wedge G/H_+$ if H' is subconjugate to H and $\phi^{H'}Y_H = 0$ if H' is not subconjugate to H. Hence, $\tau_{H'}(Y_H) = u_H$ if $H' \subseteq H$ and $\tau_{H'}(Y_H) = 0$ if $H' \not\subseteq H$. Let $Y = \bigvee_{H \in \mathrm{sub}(G)} Y_H$. Then $\tau_H(Y) = \max\{u_{H'} \mid H \subseteq H'\} = u_H$ by the assumption. This theorem gives a lower bound on the set of thick ideals in $\mathcal{SH}(G)_f$. The whole set GQ is an upper bound. Strickland was able to show that $\tau(\mathrm{Idl}(\mathcal{SH}(G)_f))$ lies in a certain proper subset of GQ (if G is nontrivial). For a cyclic group $G = \mathbb{Z}/p$, his result is the following [Str10, Proposition 16.9]:

Proposition 3.4.7. (Strickland) If $X \in S\mathcal{H}(\mathbb{Z}/p)_f$ and $K(p^{-(n+1)})_*\phi^{\{1\}}X = 0$, then $K(p^{-n})_*\phi^{\mathbb{Z}/p}X = 0$. (Note that the same prime p appears in two different roles.)

Proof. Again, the proof is taken from [Str10]. X is of the form $X = \Sigma^{\infty-V}T$, where T is a retract of a finite \mathbb{Z}/p -CW complex and V is a representation of \mathbb{Z}/p . Then, by Proposition 3.1.4, $\phi^{\mathbb{Z}/p}X = \Sigma^{\infty-\dim(V^{\mathbb{Z}/p})}T^{\mathbb{Z}/p}$. The assumption is equivalent to $K(p^{-(n+1)})_*T = 0$ and the claim is equivalent to $K(p^{-n})_*T^{\mathbb{Z}/p} = 0$. By [Rav84, Theorem 2.11], we can formulate this in terms of Johnson-Wilson theories, namely: We know $E(n+1)_*T = 0$ and have to show $E(n)_*T^{\mathbb{Z}/p} = 0$. The proof uses the Greenlees-May theory of Tate spectra [GM95]. For any G-spectrum Y, let $t_GY = F(EG_+, Y) \wedge \tilde{E}G$ and $P_GY = (t_GY)^G$. They have the following properties:

- (1) The functors t_G and P_G preserve exact triangles.
- (2) We have $t_G(X \wedge Y) = X \wedge t_G Y$ for finite *G*-spectra *X*, and $P_G(X \wedge Y) = X \wedge P_G Y$ for finite spectra *X* with trivial *G*-action.
- (3) If Y is a free G-spectrum, then $t_G Y = 0$ and so $P_G Y = 0$.
- (4) If Y is nonequivariantly contractible, then $t_G Y = 0$ and so $P_G Y = 0$.
- (5) If p divides the order of G, then the spectrum $P_G E(n+1)$ has Bousfield class $\langle P_G E(n+1) \rangle = \langle E(n) \rangle$ [HS96, Theorem 1.1].

As $E(n+1)_*T = 0$, we see that the \mathbb{Z}/p -spectrum $E(n+1) \wedge T$ is nonequivariantly contractible, so $t_G(E(n+1) \wedge T) = 0$ by (4). We also have $t_G(E(n+1) \wedge T/T^{\mathbb{Z}/p}) = 0$ by (3), so (1) implies $t_G(E(n+1) \wedge T^{\mathbb{Z}/p}) = 0$. Hence, $(P_G E(n+1)) \wedge T^{\mathbb{Z}/p} = P_G(E(n+1) \wedge T^{\mathbb{Z}/p}) = 0$ and (5) gives $E(n) \wedge T^{\mathbb{Z}/p} = 0$, as required.

This result on cyclic groups can be used to derive some restriction on the types of G-spectra that can occur for a general group G. This is done by applying the result to quotients of subgroups of G, see [Str10, Corollary 16.10]. We do not state this result here, because it needs additional notation and because we will only be interested in the case $G = \mathbb{Z}/2$ later on.

3.5 Thick ideals in $\mathcal{SH}(\mathbb{Z}/2)_f$

We apply the results of this chapter to $G = \mathbb{Z}/2$.

Recall that, for G a finite group, the thick ideals in $(\mathcal{SH}(G)_f)_{(p)}$ are characterised by their equivariant types, i.e., by the vanishing or non-vanishing of the different equivariant Morava K-theories.

Corollary 3.5.1. By the injectivity of τ (Theorem 3.4.6), any thick ideal in the category $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ is of the form

$$\mathcal{C}_{m,n} = \{ X \mid \phi^{\{1\}}(X) \in \mathcal{C}_m \text{ and } \phi^{\mathbb{Z}/2}(X) \in \mathcal{C}_n \},\$$

where $m, n \in [0, \infty]$. By Proposition 3.2.1, $X \in C_{m,n}$ is equivalent to $K(m-1, \{1\})_*(X) = 0$ and $K(n-1, \mathbb{Z}/2)_*(X) = 0$ if $0 < m, n < \infty$.

Definition 3.5.2. We say that $X \in (\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ has type $(m, n), 0 \leq m, n \leq \infty$, if $X \in \mathcal{C}_{m,n} \setminus (\mathcal{C}_{m+1,n} \bigcup \mathcal{C}_{m,n+1}).$

Let $\Gamma_p \subseteq (\mathbb{Z}_{\geq 0} \cup \{\infty\}) \times (\mathbb{Z}_{\geq 0} \cup \{\infty\})$ be the sublattice of all (m, n) such that a *p*-local spectrum of type (m, n) exists.

- **Corollary 3.5.3.** (1) Any type-(m, n) spectrum generates $C_{m,n}$ as a thick ideal (by Theorem 3.4.6).
- (2) Not all pairs (m, n) occur as the type of some spectrum. For example, if p = 2, m cannot be greater than n + 1 (by Proposition 3.4.7).
- (3) If $m \leq n$, then a type-(m, n) spectrum exists, namely $X_{m,n} = (X_m \wedge \mathbb{Z}/2_+) \vee X_n$ with $X_k \in S\mathcal{H}_{(p)}^{fin}$ a type-k spectrum and with $\mathbb{Z}/2$ acting nontrivially only on $\mathbb{Z}/2$ (by Theorem 3.4.6). Thus, $m \leq n$ implies $(m, n) \in \Gamma_p$.

These results of Strickland give upper and lower bounds for the set of thick ideals in $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ (which is bijective to Γ_p).

Remark 3.5.4. The results on thick ideals in $\mathcal{SH}(G)_f$ presented in this chapter also hold in the category of finite *G*-CW spectra, $\mathcal{SH}(G)^{fin} \subseteq \mathcal{SH}(G)_f$, as we did not use the closure under retracts in any argument.

Chapter 4

Comparison functors

Notation 4.0.1. For k a field, we write $\mathcal{SH}(k)$ for the motivic stable homotopy category over k. The standard spheres are denoted by $S^{p,q} = S_s^{\wedge (p-q)} \wedge \mathbb{G}_m^{\wedge q}$ and their \mathbb{P}^1 -suspension spectra are also denoted by $S^{p,q}$ if no confusion can arise. $\mathcal{SH}(k)$ is a tensor triangulated category, whose unit is the sphere spectrum $S = \Sigma_{\mathbb{P}^1}^{\infty} S^{0,0}$, which we also denote by S^0 . For any $E \in \mathcal{SH}(k)$, $\pi_{p,q}(E) = [S^{p,q}, E]_{\mathcal{SH}(k)}$ denotes the set of maps from $S^{p,q}$ to E in $\mathcal{SH}(k)$. For E and X in $\mathcal{SH}(k)$, let $E_{p,q}(X) = \pi_{p,q}(X \wedge E)$ and $E^{p,q}(X) = [X, S^{p,q} \wedge E]$.

For our study of thick ideals in motivic categories $S\mathcal{H}(k)$, $k \subseteq \mathbb{C}$, we want to use the given knowledge on thick ideals in the classical stable homotopy category and the $\mathbb{Z}/2$ -equivariant stable homotopy category. For this purpose, we will make use of the functors $S\mathcal{H} \xrightarrow{c_k} S\mathcal{H}(k) \xrightarrow{R_k} S\mathcal{H}$ for $k \subseteq \mathbb{C}$ and $S\mathcal{H}(\mathbb{Z}/2) \xrightarrow{c'_k} S\mathcal{H}(k) \xrightarrow{R'_k} S\mathcal{H}(\mathbb{Z}/2)$ for $k \subseteq \mathbb{R}$. The functors $c_k, R_{\mathbb{C}}$ and $R'_{\mathbb{R}}$ appear in various places in the literature but none of the sources conveniently covers all of the constructions. Most details on $R_{\mathbb{C}}$ can be found in [PPR09, Appendix] and the unstable functor $R'_{\mathbb{R}}$ is studied in [DI04, Section 5]. Other important references are [Voe96, Section 3.4], [MV99, Section 3.3] and [Ayo10]. The stable functors R'_k and c'_k are constructed and studied by Heller and Ormsby in [HO14, Section 4], which was written independently and at the same time as this chapter. Since our approach is slightly different, we give another, mostly self-contained construction of all these functors.

We start with the construction of $R_{\mathbb{C}}$ and $R'_{\mathbb{R}}$.

4.1 Symmetric $\mathbb{C}P^1$ -spectra

Objects of the motivic stable homotopy category $\mathcal{SH}(\mathbb{C})$ are spectra with respect to suspension by $\mathbb{P}^1_{\mathbb{C}} \cong S^{2,1}$. The corresponding analytic space $\mathbb{P}^1_{\mathbb{C}}(\mathbb{C})$ is $\mathbb{C}P^1$. We want that the topological realisation R(X) of a spectrum $X \in \mathcal{SH}(\mathbb{C})$ is a spectrum again. Therefore, we work with $\mathbb{C}P^1$ -spectra. The category of symmetric $\mathbb{C}P^1$ -spectra, $Sp_{\mathbb{C}P^1}^{\Sigma}$, is described in [PPR09, Theorem A.44] and is a model for the stable homotopy category. A symmetric $\mathbb{C}P^1$ -spectrum is defined in the same way as a usual symmetric spectrum with S^1 replaced by $\mathbb{C}P^1 \cong S^2$. The stable model structure is constructed analogously as for symmetric S^1 -spectra in [HSS99]. Hence, the following results also hold for symmetric $\mathbb{C}P^1$ -spectra [HSS99, Lemmas 3.4.5, 3.4.12, 3.4.13].

Proposition 4.1.1. 1. The stable trivial fibrations are the level trivial fibrations.

2. A map of $\mathbb{C}P^1$ -spectra $f: X \to Y$ is a stable fibration if and only if it is a level fibration and



is homotopy cartesian for all n, where the horizontal maps are the adjoints of the structure maps.

3. The fibrant objects are Ω -spectra, i.e., the adjoints of their structure maps are weak equivalences.

4.2 $\mathbb{Z}/2$ -equivariant symmetric spectra

For the $\mathbb{Z}/2$ -equivariant stable homotopy category we use the model constructed in [Man02]. We now recall Mandell's definitions and results. Let G be a finite group. We only need the case $G = \mathbb{Z}/2$.

Definition 4.2.1. Let S(G) denote the based simplicial *G*-set obtained by smashing together copies of the simplicial circle S^1 indexed on the elements of *G*, where the *G*-action permutes the smash factors according to the multiplication in *G*.

For example, $S(\{1\}) = S^1$ and $S(\mathbb{Z}/2) \cong S^2$, where $\mathbb{Z}/2$ acts via an orientation reversing map of degree one.

Definition 4.2.2. A symmetric *G*-spectrum consists of a based $G \times \Sigma_n$ -simplicial set T(n) for each $n \in \mathbb{N}$ and structure maps $T(n) \wedge S(G) \to T(n+1)$ such that the *m*-th iterated structure maps are $G \times \Sigma_n \times \Sigma_m$ -equivariant. Morphisms of spectra are defined in the usual way. We denote the category of symmetric *G*-spectra by $Sp^{\Sigma}(G)$.

Mandell replaces $Sp^{\Sigma}(G)$ by the isomorphic category $Sp(G\Sigma_G)$ of $G\Sigma_G$ -spectra, which is defined as follows.

Definition 4.2.3. Let Σ be the category whose objects are nonnegative integers $\underline{\mathbf{n}} = \{1, \dots, n\}$ and whose morphisms are bijections. Let Σ_G be the category Σ together with the diagram S indexed on Σ taking $\underline{\mathbf{n}}$ to the *n*-fold smash product of S(G) and with arrows permuting these smash factors.

A $G\Sigma_G$ -spectrum T is a functor from Σ to based simplicial G-sets together with natural transformations $\sigma_{n,m}: T(\underline{n}) \wedge S(\underline{m}) \to T(\underline{n+m})$ satisfying certain associativity and unitality conditions [Man02, Definition 1.3]. A morphism of $G\Sigma_G$ -spectra is a natural transformation commuting with the structure maps.

Mandell defines Ω -spectra in $Sp(G\Sigma_G)$ and constructs a projective level model structure with level equivalences, level fibrations and projective cofibrations. The resulting homotopy category is denoted by Ho^l.

Definition 4.2.4. A morphism of $G\Sigma_G$ -spectra $f: T \to U$ is called a stable equivalence if it induces bijections $\operatorname{Ho}^l(U, E) \to \operatorname{Ho}^l(T, E)$ for all Ω -spectra E.

The following theorem is [Man02, Theorem 4.1].

Theorem 4.2.5. The category $Sp(G\Sigma_G)$ has a symmetric monoidal model structure with stable equivalences as weak equivalences and projective cofibrations as cofibrations. A morphism $f: T \to U$ is a fibration if and only if it is a level fibration and



is homotopy cartesian for all $\underline{m}, \underline{n} \in \Sigma$, where the horizontal maps are adjoint to the structure maps.

Using this model structure, Mandell shows [Man02, Theorem 2]:

Theorem 4.2.6. The category $Sp^{\Sigma}(G)$ is Quillen equivalent to the stable G-equivariant category indexed on a complete G-universe.

4.3 Complex and real topological realisation functors

The aim of this section is to recall the construction of the stable topological realisation functors

$$R = R_{\mathbb{C}} : \mathcal{SH}(\mathbb{C}) \to \mathcal{SH},$$
$$R' = R'_{\mathbb{R}} : \mathcal{SH}(\mathbb{R}) \to \mathcal{SH}(\mathbb{Z}/2).$$

Various unstable and stable versions of these functors were constructed in [Voe96, Section 3.4], [MV99, Section 3.3], [Ayo10], [DI04], [PPR09, Appendix] and [HO14, Section 4].

For $k \subseteq \mathbb{C}$, let \mathbf{Sm}/k be the category of smooth schemes of finite type over k and let $\mathbf{sPre}(\mathbf{Sm}/k)$ be the category of simplicial presheaves on the Nisnevich site \mathbf{Sm}/k , see e.g. [MV99] or [Jar00, Appendix B].

We begin with the definition of $R_{\mathbb{C}}$. In the next section we will define R_k and R'_k also for subfields k of \mathbb{C} and \mathbb{R} respectively.

The functor

$$R: \mathbf{sPre}(\mathbf{Sm} \ / \mathbb{C}) \to \mathbf{sSet}$$

is defined in the following way: Any simplicial presheaf A can be written as

$$\operatorname{colim}_{X \times \Delta^n \to A} (X \times \Delta^n) \xrightarrow{\cong} A,$$

where the colimit is taken over the over-category of A, in which X runs over representable presheaves, see [PPR09, Section A.4]. We set

$$R(A) = \operatorname{colim}_{X \times \Delta^n \to A} (X(\mathbb{C}) \times \Delta^n) \in \mathbf{sSet} \,.$$

By $X(\mathbb{C})$ we actually mean the simplicial set $\operatorname{Sing}(X(\mathbb{C})^{an})$, where $X(\mathbb{C})^{an}$ denotes the set of complex points of X with the analytic topology.

Now let $k = \mathbb{R}$. Let $\mathbf{sSet}(\mathbb{Z}/2)$ denote the category of $\mathbb{Z}/2$ -simplicial sets. The functor

$$R': \mathbf{sPre}(\mathbf{Sm} / \mathbb{R}) \to \mathbf{sSet}(\mathbb{Z}/2)$$

is still defined by

$$R'(A) = \underset{X \times \Delta^n \to A}{\operatorname{colim}} (X(\mathbb{C}) \times \Delta^n),$$

but now $\mathbb{Z}/2$ acts on $X(\mathbb{C})$ by precomposing with conjugation. This induces an action of $\mathbb{Z}/2$ on R'(A), see e.g. [DI04, Section 5].

If A is pointed, then so are R(A) and R'(A) respectively.

We equip $\mathbf{sPre}(\mathbf{Sm}/k)$ with the projective model structure defined in [DI04, Section 5.1], where this category is denoted by $\operatorname{Spc}'(k)_{\operatorname{Nis}}$. The model structure for \mathbf{sSet} can be found in [Hov99, Section 3.2] and the equivariant model structure on $\mathbf{sSet}(\mathbb{Z}/2)$ is, for example, described in [Gui06, Example 4.2]:

- Weak equivalences are maps f that induce weak equivalences f^H on the fixed point sets for all $H \subseteq \mathbb{Z}/2$.
- The collection $\{(\mathbb{Z}/2)/H \times \partial \Delta^n \to (\mathbb{Z}/2)/H \times \Delta^n \mid H \subseteq \mathbb{Z}/2\}$ is a set of generating cofibrations.
- The collection $\{(\mathbb{Z}/2)/H \times \Lambda_i^n \to (\mathbb{Z}/2)/H \times \Delta^n \mid H \subseteq \mathbb{Z}/2\}$ is a set of generating trivial cofibrations.

Theorem 4.3.1. The functors R and R' and their pointed versions are strict symmetric monoidal left Quillen functors.

Proof. This is [DI04, Theorems 5.2, 5.5] and [PPR09, Theorem A.23].

Now we define the functor

Sing :
$$\mathbf{sSet} \to \mathbf{sPre}(\mathbf{Sm} / \mathbb{C}),$$

as in [PPR09, Theorem A.23]. It maps a simplicial set Z to the simplicial presheaf sending $X \in \mathbf{Sm} / \mathbb{C}$ to the simplicial set which in degree n is the set of maps $\mathbf{sSet}(X(\mathbb{C}) \times \Delta^n, Z)$, that is, $\operatorname{Sing}(Z)(X)$ is defined as an internal hom object in \mathbf{sSet} .

Proposition 4.3.2. The functor Sing is right adjoint to R.

Proof. We have to find a bijection

 $\Phi: \mathbf{sSet}(R(X), Y) \to \mathbf{sPre}(\mathbf{Sm} / \mathbb{C})(X, \operatorname{Sing}(Y))$

for any $X \in \mathbf{sPre}(\mathbf{Sm}/\mathbb{C})$ and $Y \in \mathbf{sSet}$. The general case will follow by passage to colimits after we have shown this for the case that X is representable. So let $X = \mathbf{Sm}/\mathbb{C}(-\times\Delta^{\bullet}, X')$ with $X' \in \mathbf{Sm}/\mathbb{C}$. Let $f : R(X) \to Y$ be an element of the left hand side, which now is $\mathbf{sSet}(X'(\mathbb{C}) \times \Delta^{\bullet}, Y)$. We have to define a natural transformation

 $\Phi(f): X = \mathbf{Sm} / \mathbb{C}(- \times \Delta^{\bullet}, X') \to \mathbf{sSet}(-(\mathbb{C}) \times \Delta^{\bullet}, Y) = \mathrm{Sing}(Y).$

We do this by the following composition:

$$\Phi(f): \mathbf{Sm} / \mathbb{C}(- \times \Delta^{\bullet}, X') \xrightarrow{R} \mathbf{sSet}(-(\mathbb{C}) \times \Delta^{\bullet}, X'(\mathbb{C})) \xrightarrow{f_*} \mathbf{sSet}(-(\mathbb{C}) \times \Delta^{\bullet}, Y).$$

The map Φ is obviously injective. It is also surjective because any morphism from $\operatorname{Sm}/\mathbb{C}(-\times \Delta^{\bullet}, X')$ to $\operatorname{sSet}(-(\mathbb{C}) \times \Delta^{\bullet}, Y)$ factors through the realisation functor. \Box

The $\mathbb{Z}/2$ -version of this functor,

Sing':
$$\mathbf{sSet}(\mathbb{Z}/2) \to \mathbf{sPre}(\mathbf{Sm}/\mathbb{R}),$$

maps an equivariant simplicial set Z to the simplicial presheaf sending $X \in \mathbf{Sm} / \mathbb{R}$ to the internal hom $\mathbf{sSet}(\mathbb{Z}/2)(X(\mathbb{C}), Z)$.

Proposition 4.3.3. The functor Sing' is right adjoint to R'.

Proof. The proof is the same as in the complex case, except that we have to replace \mathbf{Sm}/\mathbb{C} by \mathbf{Sm}/\mathbb{R} and \mathbf{sSet} by $\mathbf{sSet}(\mathbb{Z}/2)$.

We want to define stable versions of the functors R, R' and Sing, Sing'. Therefore, we consider the category of symmetric \mathbb{P}^1 -spectra on $\mathbf{sPre}(\mathbf{Sm}/k)$, denoted by $Sp_{\mathbb{P}^1}^{\Sigma}(k)$. The stable model structure on $Sp_{\mathbb{P}^1}^{\Sigma}(k)$ is constructed in the same way as in [Jar00], except that we start with our different definitions of fibrations and cofibrations on $\mathbf{sPre}(\mathbf{Sm}/k)$. This construction is also described in [PPR09, Section A.5]. We will only need the following information about this model structure.

Let J be a level fibrant replacement functor and let Q be the stabilisation functor defined in [Jar00, Remark 2.4]. A map f in $Sp_{\mathbb{P}^1}^{\Sigma}(k)$ is called a stable equivalence if QJ(f) is a level equivalence.

Proposition 4.3.4. A map of symmetric \mathbb{P}^1 -spectra, $f: X \to Y$, is a stable fibration if and only if it is a level fibration and

$$\begin{array}{c} X_n \longrightarrow QJX_n \\ \downarrow \qquad \qquad \downarrow \\ Y_n \longrightarrow QJY_n \end{array}$$

is homotopy cartesian [Jar00, Lemma 2.7].

A stable trivial fibration is the same as a levelwise trivial fibration [Jar00, Theorem 2.9].

Since $R(\mathbb{P}^1_{\mathbb{C}}) \cong \mathbb{C}P^1$ and $R'(\mathbb{P}^1_{\mathbb{R}}) \cong R'(S^1_s \wedge \mathbb{G}_m) \cong S^1_+ \wedge S^1_- \cong S(\mathbb{Z}/2)$ (the 2-sphere with orientation reversing $\mathbb{Z}/2$ -action) in the homotopy categories, we can define

$$R: Sp_{\mathbb{P}^1}^{\Sigma}(\mathbb{C}) \to Sp_{\mathbb{C}P^1}^{\Sigma}$$

and

$$R': Sp_{\mathbb{P}^1}^{\Sigma}(\mathbb{R}) \to Sp^{\Sigma}(\mathbb{Z}/2)$$

levelwise.

We can also extend Sing and Sing' to the categories of spectra, as follows.

Over \mathbb{C} , the simplicial presheaf \mathbb{P}^1 is equivalent to the simplicial presheaf sending $Z \in \mathbf{Sm}/\mathbb{C}$ to $\mathbf{Sm}/\mathbb{C}(Z \times \Delta^n, \mathbb{P}^1_{\mathbb{C}})$. realisation defines a map from $\mathbf{Sm}/\mathbb{C}(Z \times \Delta^n, \mathbb{P}^1_{\mathbb{C}})$ to $\mathbf{sSet}(Z(\mathbb{C}) \times \Delta^n, \mathbb{C}P^1)$. These can be assembled into a map of simplicial presheaves $\mathbb{P}^1 \to \operatorname{Sing}(\mathbb{C}P^1)$. For $X \in Sp_{\mathbb{C}P^1}^{\Sigma}$ we can, hence, define structure maps

$$\operatorname{Sing}(X_n) \wedge \mathbb{P}^1 \to \operatorname{Sing}(X_n) \wedge \operatorname{Sing}(\mathbb{C}P^1) \cong \operatorname{Sing}(X_n \wedge \mathbb{C}P^1) \xrightarrow{\operatorname{Sing}(\sigma_n)} \operatorname{Sing}(X_{n+1}),$$

so that we get a spectrum $\operatorname{Sing}(X) \in Sp_{\mathbb{P}^1}^{\Sigma}(\mathbb{C})$ defined levelwise.

Over \mathbb{R} , the same argument holds if we consider $\mathbb{Z}/2$ -equivariant maps of simplicial sets. We get a spectrum $\operatorname{Sing}'(X) \in Sp_{\mathbb{P}^1}^{\Sigma}(\mathbb{R})$ defined levelwise.

Corollary 4.3.5. The functors (R, Sing) and (R', Sing') form adjoint pairs between the categories of symmetric spectra.

Theorem 4.3.6. (The stable functors R, R')

The pairs (R, Sing) and (R', Sing') are Quillen adjunctions on the spectrum level and R, R' are strict symmetric monoidal.

Proof. The case of R is covered in [PPR09, Theorem A.45] and the claim for R' is proven in [HO14, Proposition 4.8]. For completeness, we reprove the theorem in our own words.

To show that R and R' are Quillen functors, we only have to prove that their right adjoints preserve stable fibrations between stably fibrant objects and stable trivial fibrations [Dug01, Corollary A.2]. In all model structures we are considering here, the stable trivial fibrations are the levelwise trivial fibrations. From the unstable version of this theorem it follows therefore that Sing and Sing' preserve stable trivial fibrations.

Stable fibrations are, in all of these model structures, levelwise fibrations with some additional homotopy pullback properties and stably fibrant objects are always Ω -spectra. By Propositions 4.3.2 and 4.3.3, the unstable functors Sing and Sing' are right Quillen functors. It follows that the levelwise-defined functors Sing and Sing' preserve Ω -spectra and level fibrations. Let $f: X \to Y$ be a stable fibration between Ω -spectra in $Sp_{\mathbb{C}P^1}^{\Sigma}$

with the model structure from Proposition 4.1.1 (or in $Sp^{\Sigma}(\mathbb{Z}/2)$ with the model structure from Theorem 4.2.5). We have to show that

$$\begin{array}{ccc} \operatorname{Sing}(X)_n & \longrightarrow QJ \operatorname{Sing}(X)_n \\ & & \downarrow \\ & & \downarrow \\ \operatorname{Sing}(Y)_n & \longrightarrow QJ \operatorname{Sing}(Y)_n \end{array}$$

is homotopy cartesian for all n. Since X and Y are in particular level fibrant and Sing preserves level fibrations, $J \operatorname{Sing}(X) \simeq \operatorname{Sing}(X)$ and similarly for Y. Since Q is defined using only the adjoint stucture maps, $Q \operatorname{Sing}(X) \simeq \operatorname{Sing}(X)$ and $Q \operatorname{Sing}(Y) \simeq \operatorname{Sing}(Y)$ for the Ω -spectra $\operatorname{Sing}(X)$ and $\operatorname{Sing}(Y)$. It follows that the above square is in particular homotopy cartesian.

The functors R and R' are strict symmetric monoidal, since this holds unstably and the product of symmetric spectra is defined in the same way in all the categories considered here.

4.4 Realisation functors for other fields

For $k \subseteq K$ a subfield, the canonical map $\operatorname{Spec} K \to \operatorname{Spec} k$ induces a couple of base change functors between the corresponding motivic homotopy categories. These are studied in [MV99, Section 3.1] and also in [Hu01, Section 2]. For the stable version, see [PPR09, Section A.7]. For a more general approach, see also [Ayo07].

Let $f: k \hookrightarrow K$ be the inclusion of the subfield. On the level of unpointed schemes, f^* is given by

$$f^*: \mathbf{Sm} / k \to \mathbf{Sm} / K, \ f^*(X) = X \times_{\operatorname{Spec} k} \operatorname{Spec} K.$$

It induces a functor

$$f^*: \mathbf{sPre}(\mathbf{Sm}/k) \to \mathbf{sPre}(\mathbf{Sm}/K),$$

which has a right adjoint f_* . By [PPR09, Proposition A.47], this adjunction induces a strict symmetric monoidal Quillen adjunction on the level of symmetric spectra, where f^* is given by $f^*(E)_n = f^*(E_n)$.

One can therefore define realisation functors

$$R_k : \mathcal{SH}(k) \xrightarrow{f^*} \mathcal{SH}(\mathbb{C}) \xrightarrow{R} \mathcal{SH} \text{ for } k \xrightarrow{f} \mathbb{C},$$
$$R'_k : \mathcal{SH}(k) \xrightarrow{f^*} \mathcal{SH}(\mathbb{R}) \xrightarrow{R'} \mathcal{SH}(\mathbb{Z}/2) \text{ for } k \xrightarrow{f} \mathbb{R}.$$

which are strict symmetric monoidal.

4.5 Constant presheaf functors

The following construction of the constant presheaf functors $c_k : SH \to SH(k)$ for $k \subseteq \mathbb{C}$ and $c'_k : SH(\mathbb{Z}/2) \to SH(k)$ for $k \subseteq \mathbb{R}$ is close to the one given in [HO14, Section 4]. Let $k \subseteq \mathbb{C}$. For $X \in \mathbf{sSet}$ we define $c_k(X) \in \mathbf{sPre}(\mathbf{Sm}/k)$ by $c_k(X)(Z) = X$ for all $Z \in \mathbf{Sm}/k$. Using $c_k(S^1) = S_s^1$, we extend the functor $c_k : \mathbf{sSet} \to \mathbf{sPre}(\mathbf{Sm}/k)$ levelwise to symmetric S^1 -spectra and get:

$$c_k: Sp_{S^1}^{\Sigma} \to Sp_{S_{\epsilon}^1}^{\Sigma}(k).$$

We postcompose this functor with the \mathbb{P}^1 -suspension functor, yielding a functor to the category of symmetric (S_s^1, \mathbb{P}^1) -bispectra, $c_k : Sp_{S^1}^{\Sigma} \to Sp_{S_s^1, \mathbb{P}^1}^{\Sigma}(k)$. The homotopy category of $Sp_{S_s^1, \mathbb{P}^1}^{\Sigma}(k)$ is equivalent to $\mathcal{SH}(k)$ by [Hov01, Theorem 9.1].

Theorem 4.5.1. (The stable functor c_k)

This induces a functor $c_k : SH \to SH(k)$, which is strict symmetric monoidal. It is right inverse to R_k and hence faithful. Furthermore, by [Lev14, Theorem 1], c_k is full if k is algebraically closed.

Proof. We first show that the unstable functor, $c_k : \mathbf{sSet} \to \mathbf{sPre}(\mathbf{Sm}/k)$, is a left Quillen functor. The generating cofibrations of \mathbf{sSet} are the maps $\partial \Delta^n \hookrightarrow \Delta^n$. The functor c_k takes these maps to the same maps considered as morphisms of constant simplicial presheaves. These are examples of generating cofibrations in the model structure for $\mathbf{sPre}(\mathbf{Sm}/k)$, as described in [PPR09, Section A.3]. The same applies to the generating trivial cofibrations $\Lambda_r^n \hookrightarrow \Delta^n$. The functor c_k preserves colimits by definition, hence it is a left Quillen functor. We denote its right adjoint by r_0 . It satisfies $r_0(S_s^1) = S^1$.

Now we show that $c_k : Sp_{S^1}^{\Sigma} \to Sp_{S_s}^{\Sigma}(k)$ is left Quillen, where the model structure on $Sp_{S_s}^{\Sigma}(k)$ is described in [Jar00, Section 4.5] and satisfies the analogue of Proposition 4.3.4. The right adjoint, r, to c_k is defined by levelwise application of r_0 . Since r_0 , as a right Quillen functor, preserves fibrations and trivial fibrations, r preserves level fibrations and level trivial fibrations. Stable trivial fibrations are the same as level trivial fibrations, hence these are preserved by r. We have to show that r also preserves stable fibrations between stably fibrant objects. Let $f : X \to Y$ be a stable fibration in $Sp_{S_s}^{\Sigma}(k)$ with X and Y level fibrant Ω -spectra. We have to show that r(f) is a level fibration—which we already know—and that the squares

$$\begin{array}{ccc} r(X)_n & \longrightarrow \Omega r(X)_{n+1} \\ & & \downarrow \\ & & \downarrow \\ r(Y)_n & \longrightarrow \Omega r(Y)_{n+1} \end{array}$$

are homotopy pullbacks. This is trivial because r preserves Ω -spectra (it is defined levelwise and commutes with desuspension), so r(X) and r(Y) are Ω -spectra. This proves that $c_k : Sp_{S^1}^{\Sigma} \to Sp_{S^1}^{\Sigma}(k)$ is a left Quillen functor. It is symmetric monoidal by its pointset definition and by the definition of products of symmetric spectra. The \mathbb{P}^1 -suspension functor $Sp_{S_s^1}^{\Sigma}(k) \to Sp_{S_s^1,\mathbb{P}^1}^{\Sigma}(k)$ is also a symmetric monoidal left

The \mathbb{P}^1 -suspension functor $Sp_{S_s^1}^{\Sigma}(k) \to Sp_{S_s^1,\mathbb{P}^1}^{\Sigma}(k)$ is also a symmetric monoidal left Quillen functor if the category of symmetric \mathbb{P}^1 -spectra over $Sp_{S_s^1}^{\Sigma}(k)$ is endowed with the stable model structure (see [Hov01, Theorems 5.1 and 9.1]). It follows that both functors induce a functor on the respective stable homotopy categories, and the concatenation of the induced functors, $c_k : SH \to SH(k)$, is also strict symmetric monoidal by [Hov01, Theorem 8.11].

To show that c_k is right inverse to R_k , first note that, for $f : k \hookrightarrow \mathbb{C}$, we have $f^*(\operatorname{Spec}(k)) = \operatorname{Spec}(\mathbb{C})$, which implies $f^* \circ c_k = c_{\mathbb{C}}$. So, by definition of R_k , $R_k \circ c_k = R_{\mathbb{C}} \circ f^* \circ c_k = R_{\mathbb{C}} \circ c_{\mathbb{C}}$, and it suffices to consider the case $k = \mathbb{C}$. Unstably, for $A \in \mathbf{sSet}$,

$$(R \circ c)(A) = \underset{X \times \Delta^{\bullet} \to cA}{\operatorname{colim}} (X(\mathbb{C}) \times \Delta^{\bullet}) = \underset{\Delta^{\bullet} \to A}{\operatorname{colim}} \Delta^{\bullet} = A.$$

On the level of spectra, we have used different models for constructing R and c. We therefore have to check that the following diagram is commutative, where, by definition, the composition of the upper maps induces $c : SH \to SH(\mathbb{C})$ and the lower map induces $R : SH(\mathbb{C}) \to SH$.



Since $R(S_s^1) = S^1$, the functor $R : Sp_{\mathbb{P}^1}^{\Sigma}(\mathbb{C}) \to Sp_{\mathbb{C}P^1}^{\Sigma}$ induces a functor ΣR on S_s^1 -spectra on $Sp_{\mathbb{P}^1}^{\Sigma}(\mathbb{C})$ by the levelwise definition. This induced functor is drawn as a diagonal in the above diagram and it makes the lower subdiagram commutative by definition. Thus, it suffices to check that the upper diagram is commutative. Let $X = \{X_n\}_n \in Sp_{S^1}^{\Sigma}$. Xis mapped to $\{cX_n\}_n$ in $Sp_{S_s^1}^{\Sigma}(\mathbb{C})$ and to $\{\mathbb{P}^m \wedge cX_n\}_{m,n}$ in $Sp_{S_s^1,\mathbb{P}^1}^{\Sigma}(\mathbb{C})$, which realises to $\{(\mathbb{C}P^1)^m \wedge X_n\}_{m,n}$ in $Sp_{S^1,\mathbb{C}P^1}^{\Sigma}$. This is the same as the image of X under the vertical map $Sp_{S^1}^{\Sigma} \to Sp_{S^1,\mathbb{C}P^1}^{\Sigma}$, which completes the proof that $R \circ c = \mathrm{id}$ on \mathcal{SH} .

For subfields $k \subseteq \mathbb{R}$, we want to define functors $c'_k : S\mathcal{H}(\mathbb{Z}/2) \to S\mathcal{H}(k)$ which are right inverse to R'_k . For a better understanding, we first consider $k = \mathbb{R}$ and then generalise.

To define a functor $c' : S\mathcal{H}(\mathbb{Z}/2) \to S\mathcal{H}(\mathbb{R})$ which is right inverse to R', we first construct $R' : \mathbf{sSet}(\mathbb{Z}/2) \to \mathbf{sPre}(\mathbf{Sm}/\mathbb{R})$. Observe that $R'(\operatorname{Spec} \mathbb{R}) = *$ is the onepoint set and $R'(\operatorname{Spec} \mathbb{C}) = \mathbb{Z}/2$ is the two-point set with non-trivial $\mathbb{Z}/2$ -action. We let $c'(*) = \operatorname{Spec} \mathbb{R}$ and $c'(\mathbb{Z}/2) = \operatorname{Spec} \mathbb{C}$ and extend this to $\mathbb{Z}/2$ -sets M by

$$c'(M) = \left(\prod_{M^{\mathbb{Z}/2}} \operatorname{Spec} \mathbb{R} \right) \prod \left(\prod_{(M \setminus M^{\mathbb{Z}/2})/(\mathbb{Z}/2)} \operatorname{Spec} \mathbb{C} \right).$$

This can be done functorially, just note that $-1: \mathbb{Z}/2 \to \mathbb{Z}/2$ has to be mapped to the morphism Spec $\mathbb{C} \to \text{Spec } \mathbb{C}$ induced by complex conjugation. Furthermore, c' extends to

simplicial $\mathbb{Z}/2$ -sets by $c'(M \times \Delta^n) = c'(M) \times \Delta^n$. This defines the unstable, basepoint preserving functor $c' : \mathbf{sSet}(\mathbb{Z}/2) \to \mathbf{sPre}(\mathbf{Sm}/\mathbb{R})$. We extend this to a functor of spectra by postcomposing the levelwise defined functor

$$c': Sp_{S(\mathbb{Z}/2)}^{\Sigma} \to Sp_{c'(S(\mathbb{Z}/2))}^{\Sigma}(\mathbb{R})$$

with the suspension spectrum functor $\Sigma_{\mathbb{P}^1}^{\infty} : Sp_{c'(S(\mathbb{Z}/2))}^{\Sigma}(\mathbb{R}) \to Sp_{c'(S(\mathbb{Z}/2)),\mathbb{P}^1}^{\Sigma}(\mathbb{R})$. Note that $c'(S(\mathbb{Z}/2)) \cong c'(S_+^1) \wedge c'(S_-^1) \cong S_s^1 \wedge c'(S_-^1)$ and $c'(S_-^1) = F_{\mathbb{C}/\mathbb{R}}(S^V)$ in the notation of [Hu01] (with V the sign representation), where it is shown that this is invertible in $\mathcal{SH}(\mathbb{R})$ [Hu01, Theorem 3.5]. Thus, by [Hov01, Theorem 9.1], $\Sigma_{c'(S(\mathbb{Z}/2))}^{\infty} : Sp_{\mathbb{P}^1}^{\Sigma}(\mathbb{R}) \to Sp_{c'(S(\mathbb{Z}/2)),\mathbb{P}^1}^{\Sigma}(\mathbb{R})$ is a Quillen equivalence.

Theorem 4.5.2. (The stable functor c')

This induces a functor $c' : SH(\mathbb{Z}/2) \to SH(\mathbb{R})$, which is strict symmetric monoidal and right inverse to R'. In particular, c' is faithful.

Proof. As in the previous proof, we start by considering the functor $c' : \mathbf{sSet}(\mathbb{Z}/2) \to \mathbf{sPre}(\mathbf{Sm}/\mathbb{R})$. It preserves colimits. The generating cofibrations of $\mathbf{sSet}(\mathbb{Z}/2)$ are the maps $(\mathbb{Z}/2)/H \times \partial \Delta^n \to (\mathbb{Z}/2)/H \times \Delta^n$, where $H \subseteq \mathbb{Z}/2$ is a subgroup. The images of these maps under c' can be written as pushout products:

$$c'(\mathbb{Z}/2 \times \partial \Delta^n \to \mathbb{Z}/2 \times \Delta^n) = (\emptyset \to \operatorname{Spec} \mathbb{C}) \square (\partial \Delta^n \to \Delta^n)$$
$$c'(\partial \Delta^n \to \Delta^n) = (\emptyset \to \operatorname{Spec} \mathbb{R}) \square (\partial \Delta^n \to \Delta^n).$$

These are examples of generating cofibrations for $\mathbf{sPre}(\mathbf{Sm}/\mathbb{R})$ as described in [PPR09, Section A.3]. The same argument holds for the generating trivial cofibrations $(\mathbb{Z}/2)/H \times \Lambda_r^n \to (\mathbb{Z}/2)/H \times \Delta^n$. The passage to the spectrum level works similarly as in the previous proof. The induced functor $c' : \mathcal{SH}(\mathbb{Z}/2) \to \mathcal{SH}(\mathbb{R})$ is symmetric monoidal by the same arguments as before.

By its definition, c' is right inverse to R' on the level of simplicial $\mathbb{Z}/2$ -sets. On the level of stable homotopy categories, $R' \circ c' = \text{id}$ follows from the commutativity of the following diagram, similarly as in the previous proposition.


Now let $k \subseteq \mathbb{R}$. Then $R'_k(\operatorname{Spec} k) = *$ and $R'_k(\operatorname{Spec}(k[i])) = \mathbb{Z}/2$. Therefore, we let $c'_k(*) = \operatorname{Spec} k$ and $c'_k(\mathbb{Z}/2) = \operatorname{Spec}(k[i])$ and, for a $\mathbb{Z}/2$ -set M,

$$c'_{k}(M) = \left(\coprod_{M^{\mathbb{Z}/2}} \operatorname{Spec} k\right) \coprod \left(\coprod_{(M \setminus M^{\mathbb{Z}/2})/(\mathbb{Z}/2)} \operatorname{Spec}(k[i])\right).$$

For functoriality, note that $-1 : \mathbb{Z}/2 \to \mathbb{Z}/2$ has to be mapped to $\operatorname{Spec}(k[i]) \to \operatorname{Spec}(k[i])$ induced by complex conjugation. As before, c'_k extends to $c'_k : \operatorname{sSet}(\mathbb{Z}/2) \to \operatorname{sPre}(\operatorname{Sm}/k)$ and then to

$$c'_k: Sp^{\Sigma}_{S(\mathbb{Z}/2)} \to Sp^{\Sigma}_{c'_k(S(\mathbb{Z}/2))}(k) \to Sp^{\Sigma}_{c'_k(S(\mathbb{Z}/2)),\mathbb{P}^1}(k),$$

where the first functor is defined levelwise by c'_k and the second one is the \mathbb{P}^1_k -suspension spectrum functor. Here, $c'_k(S(\mathbb{Z}/2)) \cong S^1_s \wedge c'_k(S^1_-)$ and $c'_k(S^1_-) = F_{k[i]/k}(S^V)$ in the notation of [Hu01], which is invertible in $\mathcal{SH}(k)$ by [Hu01, Theorem 3.5].

Theorem 4.5.3. (The stable functor c'_k)

This induces a functor $c'_k : S\mathcal{H}(\mathbb{Z}/2) \to S\mathcal{H}(k)$ which is strict symmetric monoidal and right inverse to R'_k .

Proof. The main claim follows exactly as in the case $k = \mathbb{R}$ considered above. It is also implied by [HO14, Theorem 4.6]. It remains to prove $R'_k \circ c'_k \cong$ id. Again, this follows from $f^* \circ c'_k \cong c'$ (for $k \stackrel{f}{\hookrightarrow} \mathbb{R}$) and $R' \circ c' \cong$ id, where $f^* \circ c'_k \cong c'$ holds because $f^*(\operatorname{Spec} k) \cong \operatorname{Spec} \mathbb{R}$ and $f^*(\operatorname{Spec}(k[i])) \cong \operatorname{Spec} \mathbb{R} \times_{\operatorname{Spec} k} \operatorname{Spec}(k[i]) \cong \operatorname{Spec} \mathbb{C}$. \Box

Remark 4.5.4. In [HO14, Theorem 1.1], Heller and Ormsby prove that if k is a real closed field, then c'_k is full after *p*-completion.

Remark 4.5.5. With similar methods as above, one can show that the functors $\mathbf{sSet} \xrightarrow{c_k} \mathbf{sPre}(\mathbf{Sm} / k) \xrightarrow{R_k} \mathbf{sSet}$ induce functors

$$\mathcal{SH} \xrightarrow{c_k} \mathcal{SH}_{S^1}(k) \xrightarrow{R_k} \mathcal{SH},$$

where $S\mathcal{H}_{S_s^1}(k)$ is the stable motivic homotopy category in which S_s^1 got inverted but \mathbb{G}_m did not.

For $k \subseteq \mathbb{R}$, the definition of the stable functor c'_k relied on the invertibility of $F_{\mathbb{C}/\mathbb{R}}(S^V)$ in $\mathcal{SH}(\mathbb{R}) = \mathcal{SH}_{\mathbb{P}^1}(\mathbb{R})$, as shown in [Hu01, Theorem 3.5]. One can show that the functors $\mathbf{sSet}(\mathbb{Z}/2) \xrightarrow{c'_k} \mathbf{sPre}(\mathbf{Sm}/k) \xrightarrow{R'_k} \mathbf{sSet}(\mathbb{Z}/2)$ induce functors

$$\mathcal{SH}_{S^1}(\mathbb{Z}/2) \xrightarrow{c'_k} \mathcal{SH}_{S^1_s}(k) \xrightarrow{R_k} \mathcal{SH}_{S^1}(\mathbb{Z}/2),$$

where $\mathcal{SH}_{S^1}(\mathbb{Z}/2)$ is the naive equivariant stable homotopy category, in which only the sphere with trivial action got inverted. The functor $R' : \mathcal{SH}_{S_s^1}(\mathbb{R}) \to \mathcal{SH}_{S^1}(\mathbb{Z}/2)$ sends $F_{\mathbb{C}/\mathbb{R}}(S^V)$ to $S^V = S(\mathbb{Z}/2)$, which is not invertible in $\mathcal{SH}_{S^1}(\mathbb{Z}/2)$. Therefore, $F_{\mathbb{C}/\mathbb{R}}(S^V)$ cannot be invertible in $\mathcal{SH}_{S_s^1}(\mathbb{R})$. This shows that it is not possible to extend $c'_k : \mathbf{sSet}(\mathbb{Z}/2) \to \mathbf{sPre}(\mathbf{Sm}/k)$ to a functor from $\mathcal{SH}(\mathbb{Z}/2)$ to $\mathcal{SH}_{S_s^1}(k)$.

Chapter 5

Thick ideals discovered by comparison functors

The aim of this chapter is to draw conclusions concerning thick subcategories and thick ideals in $\mathcal{SH}(k)$, $k \subseteq \mathbb{C}$, and in finite, local versions of this category using the functors from the previous chapter. In the next chapter, we will study thick ideals that are described by motivic Morava K-theories.

5.1 Consequences of the properties of R_k , R'_k , c_k and c'_k

- **Proposition 5.1.1.** (1) If $C \subseteq SH$ is a thick subcategory or a thick ideal, then, for $k \subseteq \mathbb{C}$, $R_k^{-1}(C) \subseteq SH(k)$ is a thick subcategory or a thick ideal, respectively.
- (2) If $\mathcal{C} \subseteq \widetilde{SH}(\mathbb{Z}/2)$ is a thick subcategory or a thick ideal, then, for $k \subseteq \mathbb{R}$, $(R'_k)^{-1}(\mathcal{C}) \subseteq S\mathcal{H}(k)$ is a thick subcategory or a thick ideal, respectively.
- (3) If $f : k \hookrightarrow K$ and $\mathcal{C} \subseteq S\mathcal{H}(K)$ is a thick subcategory or a thick ideal, then $(f^*)^{-1}(\mathcal{C}) \subseteq S\mathcal{H}(k)$ is a thick subcategory or a thick ideal, respectively.

Proof. Any functor preserves retracts, hence $R_k^{-1}(\mathcal{C})$ is closed under retracts whenever \mathcal{C} is. We also have $R_k(S_s^1) = S^1$ and R_k preserves cofibers because it is a left adjoint, hence it preserves exact triangles. Therefore, $R_k^{-1}(\mathcal{C})$ is closed under triangles whenever \mathcal{C} is. Hence, R_k^{-1} preserves thick subcategories. Since R_k is symmetric monoidal (see Chapter 4), $X \in R_k^{-1}(\mathcal{C})$ and $Y \in \mathcal{SH}(k)$ implies that $R_k(X \wedge Y) \cong R_k(X) \wedge R_k(Y)$ is in \mathcal{C} , if \mathcal{C} is a thick ideal. Thus, $X \wedge Y \in R_k^{-1}(\mathcal{C})$. That is, R_k^{-1} preserves thick ideals, too.

The proofs for $(R'_k)^{-1}$ and $(f^*)^{-1}$ are the same.

Proposition 5.1.2. For $k \subseteq \mathbb{C}$, c_k^{-1} preserves thick subcategories and thick ideals. Similarly, for $k \subseteq \mathbb{R}$, $(c'_k)^{-1}$ preserves thick subcategories and thick ideals.

Proof. Since $S_s^1 = c_k(S^1)$ and c_k preserves mapping cones, c_k preserves exact triangles. It also preserves retracts and is strict symmetric monoidal, hence, c_k^{-1} preserves thick subcategories and thick ideals.

5.2 Finite motivic spectra

The thick subcategory theorem of [HS98] concerns the category of finite spectra, \mathcal{SH}^{fin} , as defined in Chapter 1. The functors R_k and c_k can therefore only help us to understand subcategories of $\mathcal{SH}(k)$ which are at most as big as $R_k^{-1}(\mathcal{SH}^{fin})$. There are multiple equivalent possibilities to define \mathcal{SH}^{fin} , using the notions of finite CW-spectra, dualisable objects or compact objects. These notions are not equivalent in the motivic setting, therefore we obtain more than one possible category of "finite" objects in $\mathcal{SH}(k)$.

We will now discuss the various versions of finiteness. Let k be any field.

- **Definition 5.2.1.** (1) The category $\mathcal{SH}(k)^{fin}$ of finite cellular motivic spectra over a field k is the smallest full subcategory of $\mathcal{SH}(k)$ that contains the spheres $S^{p,q}$ for all $p, q \in \mathbb{Z}$ and is closed under exact triangles [DI05, Definition 8.1].
- (2) For $k \subseteq \mathbb{R}$, let $\mathcal{SH}(k)^{fin+}$ be the smallest full subcategory of $\mathcal{SH}(k)$ that contains $S^{p,q} \wedge (\operatorname{Spec} k[i])^{\wedge m}_+$ for all $p,q \in \mathbb{Z}, m \geq 0$ and is closed under exact triangles.
- (3) The closures of $\mathcal{SH}(k)^{fin}$ and $\mathcal{SH}(k)^{fin+}$ under colimits are denoted by $\mathcal{SH}(k)^{cell}$ and $\mathcal{SH}(k)^{cell+}$. Their objects are called cellular, see [DI05, Definitions 2.1 and 2.10].

Remark 5.2.2. Note that these categories are closed under \wedge because so are their sets of generators and because \wedge preserves exact triangles and colimits, as it is a left adjoint.

With this definition, $\mathcal{SH}(k)^{fin}$ is the smallest tensor triangulated full subcategory of $\mathcal{SH}(k)$ that contains $c_k(\mathcal{SH}^{fin})$ and is closed under $-\wedge \mathbb{G}_m^{\pm 1}$, and $\mathcal{SH}(k)^{fin,+}$ is the smallest tensor triangulated full subcategory that contains $c'_k(\mathcal{SH}(\mathbb{Z}/2)^{fin})$ and is closed under $-\wedge \mathbb{G}_m^{\pm 1}$.

The following results can mostly be found in [NSØ09, Section 4].

- **Definition 5.2.3.** (1) Let $\mathcal{D} \subseteq S\mathcal{H}(k)$ be the collection of all (strongly) dualisable objects. That is, all spectra X such that the canonical map $F(X, S) \wedge Y \to F(X, Y)$ is an isomorphism for all $Y \in S\mathcal{H}(k)$, where F(-, -) denotes the derived internal hom in $S\mathcal{H}(k)$ and $S = S^{0,0}$ is the sphere spectrum. F(X, S) is called the dual of X and is also denoted by DX (compare Definition 3.1.2).
- (2) A motivic spectrum $F \in \mathcal{SH}(k)$ is called compact if $\operatorname{Hom}_{\mathcal{SH}(k)}(F, -)$ preserves arbitrary sums. Let $\mathcal{SH}(k)_f \subseteq \mathcal{SH}(k)$ denote the full subcategory of compact objects.

Remark 5.2.4. (1) $SH(k)_f$ is a thick subcategory of SH(k) [NSØ09, Section 4].

- (2) Any dualisable object is also compact, as shown in the proof of Proposition 3.1.3.
- (3) The smash product of two dualisable objects X and Y is again dualisable, because $F(X \land Y, S) \land Z \cong F(X, F(Y, S)) \land Z \cong F(X, S) \land F(Y, S) \land Z \cong F(X, S) \land F(Y, Z) \cong F(X, F(Y, Z)) \cong F(X \land Y, Z).$ Similarly, compact objects are closed under \land .
- (4) By [Hu01, Cor. 2.14 and Thm. 4.1], $F(\operatorname{Spec}(k[i])_+, E) \cong \operatorname{Spec}(k[i])_+ \wedge E$ in $\mathcal{SH}(k)$, $k \subseteq \mathbb{R}$. That is, $\Sigma^{\infty} \operatorname{Spec}(k[i])_+$ is self-dual and in particular compact.

Definition 5.2.5. For \mathcal{R} a collection of objects in $\mathcal{SH}(k)_f$, let $\mathcal{SH}(k)_{\mathcal{R},f} \subseteq \mathcal{SH}(k)_f$ be the smallest thick subcategory containing \mathcal{R} .

Let $\mathcal{T}_k = \{S^{p,q} \mid p,q \in \mathbb{Z}\}$ be the collection of all motivic spheres in $\mathcal{SH}(k)$ and let $\mathcal{T}_k^+ = \{S^{p,q} \land (\operatorname{Spec} k[i])_+^{\land m} \mid p,q \in \mathbb{Z}, m \geq 0\}$ if $k \subseteq \mathbb{R}$. These are sets of compact objects in $\mathcal{SH}(k)$ by Remark 5.2.4, parts (2) and (4).

Comparing the definitions, we see that $\mathcal{SH}(k)_{\mathcal{T}_k,f}$ is the closure of $\mathcal{SH}(k)^{fin}$ under retracts and $\mathcal{SH}(k)_{\mathcal{T}_k^+,f}$ is the closure of $\mathcal{SH}(k)^{fin+}$ under retracts.

Proposition 5.2.6. $SH(k)_{D,f} \subseteq SH(k)_f$ is the full subcategory of dualisable objects of SH(k).

Proof. This is [NSØ09, Lemma 4.2]. By Remark 5.2.4(2), \mathcal{D} is a collection of compact objects. Furthermore, the full subcategory spanned by \mathcal{D} is already a thick subcategory.

Since $\mathcal{SH}(k)_{\mathcal{D},f} \subseteq \mathcal{SH}(k)_f$ is a thick subcategory and $\mathcal{SH}(k)_f \subseteq \mathcal{SH}(k)$ is a thick subcategory, it follows that the strongly dualisable objects form a thick subcategory of $\mathcal{SH}(k)$. Note also that all the categories mentioned above are closed under \wedge .

Proposition 5.2.7. We have $\mathcal{SH}(k)^{fin} \subseteq \mathcal{SH}(k)_{\mathcal{T}_k,f} \subseteq \mathcal{SH}(k)_{\mathcal{D},f} \subseteq \mathcal{SH}(k)_f$ and for $k \subseteq \mathbb{R}$, $\mathcal{SH}(k)^{fin+} \subseteq \mathcal{SH}(k)_{\mathcal{T}_k^+,f} \subseteq \mathcal{SH}(k)_{\mathcal{D},f} \subseteq \mathcal{SH}(k)_f$. In particular, all objects in $\mathcal{SH}(k)^{fin}$ and $\mathcal{SH}(k)^{fin+}$ are strongly dualisable. Furthermore, $\mathcal{SH}(k)^{fin}$ and $\mathcal{SH}(k)^{fin+}$ are closed under taking duals.

Proof. The first line and the case $\mathcal{SH}(k)^{fin}$ are proven in [NSØ09, Section 4]. For $k \subseteq \mathbb{R}$, the only additional input is the self-duality of $\Sigma^{\infty} \operatorname{Spec}(k[i])_+$.

Remark 5.2.8. [NSØ09, Remark 8.2] gives an example for an object in SH(S) that is compact but not dualisable, where S is the spectrum of a discrete valuation ring.

A stronger result holds if k is a field of characteristic 0. It is also proven in [Rio05].

Proposition 5.2.9. Let k be of characteristic 0. Then $\mathcal{SH}(k)_{\mathcal{D},f} = \mathcal{SH}(k)_f$ is the thick subcategory of $\mathcal{SH}(k)$ generated by $\{\Sigma^{2n,n}\Sigma^{\infty}U_+ \mid U \in \mathbf{Sm}/k, n \in \mathbb{Z}\}$. Hence, any object of $\mathcal{SH}(k)$ is dualisable if and only if it is compact.

Proof. By [DI05, Theorem 9.2], $\{\Sigma^{2n,n}\Sigma^{\infty}U_+ \mid U \in \mathbf{Sm} / k, n \in \mathbb{Z}\}\$ is a set of compact generators for $\mathcal{SH}(k)$, which means two things: First, these objects are compact and second, the only full triangulated subcategory of $\mathcal{SH}(k)$ containing this set and being closed under infinite direct sums is $\mathcal{SH}(k)$ itself. Since schemes are locally affine, also $\{\Sigma^{2n,n}\Sigma^{\infty}U_+ \mid U \in \mathbf{Sm} / k$ quasi-projective} is a set of compact generators. General theory [MS06, Theorem 13.1.14] implies that $\mathcal{SH}(k)_f$ is the thick subcategory of $\mathcal{SH}(k)$ generated by $\{\Sigma^{2n,n}\Sigma^{\infty}U_+ \mid U \in \mathbf{Sm} / k$ quasi-projective}. By [RØ08, Theorem 4.9], $\Sigma^{\infty}U_+$ is dualisable for any such U. Since dualisability is preserved by exact triangles and retracts, the thick subcategory generated by $\{\Sigma^{2n,n}\Sigma^{\infty}U_+ \mid U \in \mathbf{Sm} / k$ quasi-projective} is contained in $\mathcal{SH}(k)_{\mathcal{D},f}$. Thus, $\mathcal{SH}(k)_f = \mathcal{SH}(k)_{\mathcal{D},f}$. For $k \subseteq \mathbb{C}$ $(k \subseteq \mathbb{R})$ all these categories are furthermore included in the preimage of compact topological spectra under R_k (R'_k) :

Proposition 5.2.10. *For* $k \subseteq \mathbb{C}$ *,*

$$\mathcal{SH}(k)_f \subseteq R_k^{-1}(\mathcal{SH}^{fin})$$

and for $k \subseteq \mathbb{R}$,

$$\mathcal{SH}(k)_f \subseteq R'^{-1}_k(\mathcal{SH}(\mathbb{Z}/2)_f).$$

Proof. We have to show that R_k and R'_k preserve compact objects. Let $f: k \hookrightarrow \mathbb{C}$. Then f^* restricts to a functor between the categories of compact objects: $f^*: S\mathcal{H}(k)_f \to S\mathcal{H}(\mathbb{C})_f$, because the base change functor preserves smooth schemes and so f^* sends a compact generator $\Sigma^{(2n,n)}\Sigma^{\infty}U_+$, $U \in \mathbf{Sm}/k$, $n \in \mathbb{Z}$, of $S\mathcal{H}(k)$ to a compact generator of $S\mathcal{H}(\mathbb{C})$. Hence, for the first claim it suffices to prove that $R = R_{\mathbb{C}}$ preserves compact objects. Similarly, for the second claim it suffices to show that $R' = R'_{\mathbb{R}}$ preserves compact objects.

Let $X = \Sigma^{(2n,n)} \Sigma^{\infty} U_+$ be a compact generator of $\mathcal{SH}(\mathbb{C})$. As in the proof of the previous proposition, we can assume that U is a smooth quasi-projective scheme. By Jouanolou's trick [Jou73, Lemma 1.5], there exists an affine vector bundle torsor over U. This is a vector bundle $E \to U$, together with a torsor $p: V \to U$ on E with Vaffine. By the definition of a torsor, V is locally isomorphic to E. This implies that for some $m, U \times \mathbb{A}^m$ is locally isomorphic to an affine smooth scheme V. In particular, Uis \mathbb{A}^1 -equivalent to V, and so $X \cong \Sigma^{(2n,n)} \Sigma^{\infty} V_+$ in $\mathcal{SH}(\mathbb{C})$. Now, since V is smooth and affine, its complex realisation has the homotopy type of a finite CW complex by [Laz04, Example 3.1.9]. Hence, $R(X) \in \mathcal{SH}$ is isomorphic to $\Sigma^n \Sigma^{\infty} Y_+$ for some finite CW complex Y and, so, R(X) is compact by Remark 2.0.7.

The proof that for any smooth and affine variety $V, V(\mathbb{C}) \subseteq \mathbb{C}^r$ has the homotopy type of a finite CW-complex, can be summarised as follows [Laz04, Example 3.1.9]: $V(\mathbb{C})$ is a complex submanifold of \mathbb{C}^r without boundary (because V is smooth, see e.g. [Hul12, Section 3.1.2]), which is closed as a subset of \mathbb{C}^r (because the zero locus of any polynomial is closed). For almost any $c \in \mathbb{C}^r$, the squared distance function $\phi_c : V(\mathbb{C}) \to \mathbb{R}, \phi_c(x) = ||x - c||^2$, has only non-degenerate critical points [Mil63, Theorem 6.6]. Furthermore, ϕ_c being real algebraic implies that it has only finitely many critical points, as in [Laz04, Example 3.1.9]. Using Morse theory [Mil63, Theorem 3.5], it follows that $V(\mathbb{C})$ has the homotopy type of a CW complex with one cell of dimension n for each critical point of ϕ_c of index n.

For X a compact generator of $\mathcal{SH}(\mathbb{R})$, we also have $X \cong \Sigma^{(2n,n)} \Sigma^{\infty} V_+$ and now V is a smooth and affine real variety. Its realisation $R'(V) = V(\mathbb{C}) \subseteq \mathbb{C}^r$ is still a complex manifold, which is closed, but with the property that, for any $x \in V(\mathbb{C})$, also its complex conjugate \bar{x} lies in $V(\mathbb{C})$. This was used for the definition of the $\mathbb{Z}/2$ -action on $V(\mathbb{C})$: $\rho(x) = \bar{x}$ for $\rho \in \mathbb{Z}/2$ the generator. As before, we can choose $c \in \mathbb{C}^r$ such that ϕ_c has finitely many critical points, which are all non-degenerate. If $c \in \mathbb{R}^r$, ϕ_c is $\mathbb{Z}/2$ -invariant and the claim follows from equivariant Morse theory: by the proof of [May89, Theorem 2.2], any invariant Morse function can be turned into a special invariant Morse function and, by [May89, Theorem 3.3], a manifold with a special invariant Morse function is equivariantly homotopy equivalent to an equivariant CW complex with one equivariant cell for each critical orbit. Thus, $V(\mathbb{C})$ is equivalent to a finite $\mathbb{Z}/2$ -CW complex, whose suspension spectrum is compact in $\mathcal{SH}(\mathbb{Z}/2)$ by Proposition 3.1.3. For more information on equivariant Morse theory in English language, we refer the reader to [Was69].

Now, assume $c \in \mathbb{C}^r \setminus \mathbb{R}^r$. We would like to take $x \mapsto \min(\phi_c(x), \phi_c(\bar{x}))$ as a Morse function. It is continuous and $\mathbb{Z}/2$ -invariant but it is not differentiable for $x \in \mathbb{R}^r$. Outside of \mathbb{R}^r , the critical points of $\min(\phi_c(x), \phi_c(\bar{x}))$ are a subset of the critical points of ϕ_c and their complex conjugates. The idea is to proceed in two steps: first, to take care of the real part $V(\mathbb{C}) \cap \mathbb{R}^r = V(\mathbb{R}) = V(\mathbb{C})^{\mathbb{Z}/2}$ and, second, to use $\min(\phi_c(x), \phi_c(\bar{x}))$ as a Morse function away from \mathbb{R}^r .

 $V(\mathbb{R}) \subseteq \mathbb{R}^r$ is a manifold, which is closed and without boundary and for which there exists $d \in \mathbb{R}^r$ such that $\phi_d : V(\mathbb{R}) \to \mathbb{R}$ is a Morse function with finitely many critical points (analogously to $V(\mathbb{C}) \subseteq \mathbb{C}^r$). Let B be an open ball around $0 \in \mathbb{C}^r$ which contains a ball D(d) around d containing all critical points of ϕ_d and a ball D(c) around c containing all critical points of ϕ_c and their conjugates. $V(\mathbb{R})$ can be contracted inside \mathbb{R}^r to $V(\mathbb{R}) \cap \overline{B}$, following the orthogonal trajectories of the hypersurfaces on which ϕ_d is constant, until \overline{B} is reached (that is, we glue an infinite sequence of the diffeomorphisms $M^a \cong M^b$ constructed in the proof of [Mil63, Theorem 3.1] to one homotopy equivalence). We extend this homotopy equivalence to a narrow open neighborhood $U \subset \mathbb{C}^r$ of $\mathbb{R}^r \setminus (B \cap \mathbb{R}^r)$ so that $\mathbb{C}^r \simeq \mathbb{C}^r \setminus U$ by a $\mathbb{Z}/2$ -equivariant homotopy equivalence which does not add critical points to $\min(\phi_c(x), \phi_c(\bar{x}))$ outside of B. Using this homotopy equivalence, we can assume that $V(\mathbb{C}) \setminus B$ does not contain any points near the real subspace \mathbb{R}^r . Now we use ϕ_c to contract $V(\mathbb{C})$ to $V(\mathbb{C}) \cap \overline{B}$ with an equivariant homotopy equivalence: any $x \in V(\mathbb{C}) \setminus B$ with $\phi_c(x) < \phi_c(\bar{x})$ gets moved into B along the line of steepest descent of ϕ_c inside $V(\mathbb{C})$. If, however, $\phi_c(\bar{x}) < \phi_c(x)$, x gets moved along the line of steepest descent of $\phi_{\bar{c}}$. Note that $\phi_{\bar{c}}(x) = \phi_c(\bar{x})$ and that the homotopy equivalence is well defined because we removed \mathbb{R}^r before and because all critical points of ϕ_c and $\phi_{\bar{c}}$ are inside B. Hence, this defines a $\mathbb{Z}/2$ -equivariant homotopy equivalence between $V(\mathbb{C})$ and $V(\mathbb{C}) \cap B$.

Thus, $V(\mathbb{C})$ is $\mathbb{Z}/2$ -homotopy equivalent to a $\mathbb{Z}/2$ -manifold (with boundary) which is closed and bounded and, hence, compact in \mathbb{C}^r . Using equivariant Morse theory, this compact manifold is $\mathbb{Z}/2$ -homotopy equivalent to a finite $\mathbb{Z}/2$ -CW complex (again, see [May89, Theorems 2.2, 3.3] or [Was69]). Note that, although most of the literature only studies Morse theory for manifolds without boundary, it still covers this case for the following reason: Since $D(c) \subset B$, the Morse function ϕ_c evaluated on the boundary of $V(\mathbb{C}) \cap \overline{B}$ takes a higher value than on any critical point. Therefore, $V(\mathbb{C}) \cap \overline{B}$ appears at a finite step in the Morse theoretical construction of the CW complex associated with ϕ_c , i.e., $V(\mathbb{C}) \cap \overline{B} \simeq V(\mathbb{C})^m = \phi_c^{-1}([0,m])$ for some $m \in \mathbb{R}$, which is a finite $\mathbb{Z}/2$ -CW complex.

5.3 Motivic thick ideals

Let $k \subseteq \mathbb{C}$ and p be any prime. The following theorem identifies important families of thick ideals in $(\mathcal{SH}(k)_f)_{(p)}$. It is the main result in this chapter.

Theorem 5.3.1. (Lower bound on the number of motivic thick ideals)

- (1) The category $(\mathcal{SH}(k)_f)_{(p)}$ contains at least an infinite chain of different thick ideals, given by $\overline{R}_k^{-1}(\mathcal{C}_n)$, $0 \le n \le \infty$, where \overline{R}_k denotes the p-localisation of the restriction of R_k to $\mathcal{SH}(k)_f$ and $\mathcal{C}_n \subseteq \mathcal{SH}_{(p)}^{fin}$ is as defined in Chapter 1.
- (2) If $k \subseteq \mathbb{R}$, then $(\mathcal{SH}(k)_f)_{(p)}$ contains at least a two-dimensional lattice of different thick ideals, given by $(\overline{R}'_k)^{-1}(\mathcal{C}_{m,n})$, for all $(m,n) \in \Gamma_p$ as in Definition 3.5.2.

Proof. We prove the first part, the second is proven similarly. By Remark 5.2.2, Proposition 5.2.7 and Proposition 5.2.10, $c_k(\mathcal{SH}^{fin}) \subseteq \mathcal{SH}(k)_f \subseteq R_k^{-1}(\mathcal{SH}^{fin})$, hence c_k and R_k restrict to functors $\mathcal{SH}^{fin} \xrightarrow{c_k} \mathcal{SH}(k)_f \xrightarrow{\overline{R}_k} \mathcal{SH}^{fin}$. Since the motivic *p*-local Moore spectrum (the homotopy colimit of a diagram of sphere spectra whose arrows are multiplications by integers prime to *p*) is the image of the topological *p*-local Moore spectrum under c_k , we get induced functors between the localised categories,

$$\mathcal{SH}_{(p)}^{fin} \xrightarrow{c_k} (\mathcal{SH}(k)_f)_{(p)} \xrightarrow{\overline{R}_k} \mathcal{SH}_{(p)}^{fin},$$

which still have the properties that c_k and \overline{R}_k preserve exact triangles and smash products and that $\overline{R}_k \circ c_k = \text{id}$, as proven in Theorem 4.5.1. Similarly to Proposition 5.1.1, it follows that $\overline{R}_k^{-1}(\mathcal{C}_n)$ is a thick ideal. Now let n < m. Then $\mathcal{C}_m \subset \mathcal{C}_n$ in $\mathcal{SH}_{(p)}^{fin}$ and there is some $X \in \mathcal{C}_n \setminus \mathcal{C}_m$. It follows that $\overline{R}_k^{-1}(\mathcal{C}_m) \subseteq \overline{R}_k^{-1}(\mathcal{C}_n)$ and that $c_k(X) \in \overline{R}_k^{-1}(\mathcal{C}_n) \setminus \overline{R}_k^{-1}(\mathcal{C}_m)$. Thus, $\overline{R}_k^{-1}(\mathcal{C}_n)$, $0 \le n \le \infty$, form a chain of pairwise different thick ideals.

For the second part, one needs to consider the functors

$$(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)} \xrightarrow{c'_k} (\mathcal{SH}(k)_f)_{(p)} \xrightarrow{\overline{R}'_k} (\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}.$$

In the following, we will omit the overline and use the notation R_k , R'_k for the (p-localised) restricted functors, too.

Remark 5.3.2. In the above theorem, $\mathcal{SH}(k)_f$ can be replaced by any other tensor triangulated full subcategory $\mathcal{D} \subseteq \mathcal{SH}(k)$ satisfying

$$c_k(\mathcal{SH}^{fin}) \subseteq \mathcal{D} \subseteq R_k^{-1}(\mathcal{SH}^{fin}), \text{ or }$$

$$c'_k(\mathcal{SH}(\mathbb{Z}/2)^{fin+}) \subseteq \mathcal{D} \subseteq (R'_k)^{-1}(\mathcal{SH}(\mathbb{Z}/2)_f)$$
 respectively.

In particular, the theorem applies to any

$$\mathcal{D} \in \{\mathcal{SH}(k)^{fin}, \ \mathcal{SH}(k)^{fin+}, \ \mathcal{SH}(k)_{\mathcal{T}_k, f}, \ \mathcal{SH}(k)_{\mathcal{T}_k^+, f}, \ R_k^{-1}(\mathcal{SH}^{fin}), \ R_k'^{-1}(\mathcal{SH}(\mathbb{Z}/2)_f)\}$$

for the following reasons. Recall from Remark 3.5.4 that Strickland's characterisation of equivariant thick ideals works in $\mathcal{SH}(G)^{fin}$ as well as in its closure under retracts, $\mathcal{SH}(G)_f$. Therefore, we can here take $\mathcal{SH}(k)^{fin(+)}$ as well as its closure under retracts, $\mathcal{SH}(k)_{\mathcal{T}_k^{(+)},f}$. Note also that all categories mentioned here are closed under \wedge because they are generated as thick or triangulated subcategories by classes of objects closed under \wedge : $\mathcal{SH}(k)^{fin(+)}$ and $\mathcal{SH}(k)_{\mathcal{T}_k^{(+)},f}$ are generated by $\{S^{p,q}(\wedge \operatorname{Spec}(k[i])^{\wedge m}_+)\}$ and $\mathcal{SH}(k)_f$ is generated by smooth schemes, which are also closed under smash product. $R_k^{-1}(\mathcal{SH}^{fin})$ and $(R'_k)^{-1}(\mathcal{SH}(\mathbb{Z}/2)_f)$ are closed under \wedge because \wedge commutes with R_k , R'_k .

Remark 5.3.3. The construction of the functors R_k and c_k does not depend on the fact that \mathbb{P}^1 is invertible in $\mathcal{SH}(k)$. They can also be constructed for the category $\mathcal{SH}_{S_s^1}(k)$ in which only $S_s^1 = S^{1,0}$ got inverted and \mathbb{G}_m did not. Therefore, part (1) of the theorem also holds for $(\mathcal{SH}_{S_s^1}(k)_f)_{(p)}$. The construction of c'_k , however, needed the invertibility of \mathbb{P}^1 (see Remark 4.5.5). Thus, part (2) cannot be applied to $(\mathcal{SH}_{S_s^1}(k)_f)_{(p)}$.

Definition 5.3.4. For a full subcategory C of a tensor triangulated category \mathcal{T} , let $\Delta(C)$ denote the smallest thick subcategory of \mathcal{T} that contains C and recall that thickid(C) denotes the smallest thick ideal that contains C.

We state two more observations about thick ideals in the category $(\mathcal{SH}(k)_f)_{(p)}$. They also hold in any of the above categories \mathcal{D} .

Proposition 5.3.5. Let $X_n \in S\mathcal{H}_{(p)}^{fin}$ be any spectrum of type n, i.e. $X_n \in \mathcal{C}_n \setminus \mathcal{C}_{n+1}$ (see Definition 2.0.11) and let $X_{m,n} \in S\mathcal{H}(\mathbb{Z}/2)_{(p)}^{fin}$ be any spectrum of type (m,n) (see Definition 3.5.2). Then

> thickid $(c_k X_n)$ = thickid $(c_k C_n)$ in $(\mathcal{SH}(k)_f)_{(p)}$ if $k \subseteq \mathbb{C}$ and thickid $(c'_k X_{m,n})$ = thickid $(c'_k C_{m,n})$ in $(\mathcal{SH}(k)_f)_{(p)}$ if $k \subseteq \mathbb{R}$.

Proof. It is clear that $\operatorname{thickid}(c_k X_n) \subseteq \operatorname{thickid}(c_k \mathcal{C}_n)$. Since c_k^{-1} preserves thick ideals by Proposition 5.1.2, $c_k^{-1}(\operatorname{thickid}(c_k X_n))$ is a thick ideal containing X_n . Since \mathcal{C}_n is the smallest thick ideal containing X_n , we have $\mathcal{C}_n \subseteq c_k^{-1}(\operatorname{thickid}(c X_n))$ and hence $c_k \mathcal{C}_n \subseteq \operatorname{thickid}(c_k X_n)$, which implies the first claim. The same proof shows the second claim.

Proposition 5.3.6. If $X, Y \in (\mathcal{SH}(k)_f)_{(p)}$ with $\operatorname{type}(R_k X) \neq \operatorname{type}(R_k Y)$ (see Definition 2.0.11) or, if $k \subseteq \mathbb{R}$, $\operatorname{type}(R'_k X) \neq \operatorname{type}(R'_k Y)$ (see Definition 3.5.2), then

$$\operatorname{thickid}(X) \neq \operatorname{thickid}(Y).$$

Proof. Let type $(R_kX) = n >$ type (R_kY) . Then thickid $(X) \subseteq R_k^{-1}(\mathcal{C}_n)$ but $Y \notin R_k^{-1}(\mathcal{C}_n)$. The case of R'_k is similar.

The next proposition gives a description of thick ideals in the categories of finite cellular spectra, $\mathcal{SH}(k)^{fin}$, $k \subseteq \mathbb{C}$, and $\mathcal{SH}(k)^{fin+}$, $k \subseteq \mathbb{R}$, from Definition 5.2.1. In this case, a thick ideal is a thick subcategory that is closed under $-\wedge \mathbb{G}_m^{\pm 1}$ and under $-\wedge \operatorname{Spec}(k[i])_+$ if $k \in \mathbb{R}$.

Proposition 5.3.7. Let $C \subseteq SH(k)^{fin}$ be a subcategory, $k \subseteq \mathbb{C}$. Then

thickid(
$$\mathcal{C}$$
) = $\Delta\left(\bigcup_{n\in\mathbb{Z}}\mathcal{C}\wedge\mathbb{G}_m^{\wedge n}\right)$.

For a subcategory $\mathcal{C} \subseteq \mathcal{SH}(k)^{fin+}$, $k \subseteq \mathbb{R}$, we have

thickid(
$$\mathcal{C}$$
) = $\Delta \left(\bigcup_{n \in \mathbb{Z}} (\mathcal{C} \wedge \mathbb{G}_m^{\wedge n}) \cup \bigcup_{n \in \mathbb{Z}} (\mathcal{C} \wedge \mathbb{G}_m^{\wedge n} \wedge \operatorname{Spec}(k[i])_+) \right)$

Proof. We prove the second claim, the proof of the first claim is slightly shorter. Let $k \subseteq \mathbb{R}$. The subcategory thickid(\mathcal{C}) contains

$$\overline{\mathcal{C}} = \Delta\left(\bigcup_{n \in \mathbb{Z}} (\mathcal{C} \wedge \mathbb{G}_m^n) \cup \bigcup_{n \in \mathbb{Z}} (\mathcal{C} \wedge \mathbb{G}_m^n \wedge \operatorname{Spec}(k[i])_+)\right)$$

because thickid(\mathcal{C}) is closed under $-\wedge \mathbb{G}_m^n$ and $-\wedge \operatorname{Spec}(k[i])_+$ and is a thick subcategory. We have to show that the triangulated subcategory $\overline{\mathcal{C}}$ is already a thick ideal. Let $\mathcal{D} \subseteq S\mathcal{H}(k)^{fin+}$ be the full subcategory consisting of all objects X such that $\overline{\mathcal{C}}$ is closed under $-\wedge X$. We have to show $\mathcal{D} = S\mathcal{H}(k)^{fin+}$. First note that \mathcal{D} contains all spheres $S^{p,q}$ because, as a triangulated subcategory, $\overline{\mathcal{C}}$ is closed under $-\wedge S^{p,0}$ and because we added $S^{q,q}$. If $X \to Y \to Z$ is a triangle with two objects in \mathcal{D} , then $A \wedge X \to A \wedge Y \to A \wedge Z$ is a triangle with two objects in $\overline{\mathcal{C}}$, hence the third object is also in $\overline{\mathcal{C}}$. It follows that \mathcal{D} is closed under exact triangles. Furthermore, \mathcal{D} is closed under \wedge by definition. It remains to show that \mathcal{D} contains $\Sigma_T^{\infty} \operatorname{Spec}(k[i])_+$. By the equivalence

$$\operatorname{Spec}(k[i])_{+} \wedge \operatorname{Spec}(k[i])_{+} \cong \operatorname{Spec}(k[i])_{+} \vee \operatorname{Spec}(k[i])_{+} \cong \operatorname{cof}\left(\operatorname{Spec}(k[i])_{+} \xrightarrow{0} \operatorname{Spec}(k[i])_{+}\right),$$

any triangulated subcategory of $\mathcal{SH}(k)$ containing the spectrum $\Sigma_T^{\infty} \operatorname{Spec}(k[i])_+$ also contains $\Sigma_T^{\infty} \operatorname{Spec}(k[i])_+^{ll}, l \geq 1$. It follows that $\overline{\mathcal{C}}$ contains $\mathcal{C} \wedge \mathbb{G}_m^n \wedge \operatorname{Spec}(k[i])_+^{ll}, l \geq 0$. Hence, \mathcal{D} contains $\Sigma_T^{\infty} \operatorname{Spec}(k[i])_+$ and is thus equal to $\mathcal{SH}(k)^{fin+}$.

For $k \subseteq \mathbb{C}$, we have so far identified the following thick ideals in $(\mathcal{SH}(k)_f)_{(p)}$:

For $k \subseteq \mathbb{R}$, the picture has another dimension and depends on the classification of thick ideals in the equivariant category as described in Chapter 3. There is at least one spot where the inclusion from the lower row into the upper row is actually an equality.

Proposition 5.3.8.

$$R_k^{-1}(\mathcal{C}_0) = \text{thickid}(c_k(\mathcal{C}_0)) = (\mathcal{SH}(k)_f)_{(p)} \text{ and}$$
$$(R_k')^{-1}(\mathcal{C}_{0,0}) = \text{thickid}(c_k'(\mathcal{C}_{0,0})) = (\mathcal{SH}(k)_f)_{(p)}.$$

Proof. This is because all these subcategories contain the sphere spectrum and are closed under smashing with arbitrary elements of $(\mathcal{SH}(k)_f)_{(p)}$.

Chapter 6

Thick ideals associated with cohomology theories

6.1 Equivalence of homology and cohomology theories

We will now concentrate on the categories of finite cellular spectra $\mathcal{SH}(k)^{fin}$, $\mathcal{SH}(k)^{fin+}$, and on (co)homology theories represented by objects in $\mathcal{SH}(k)^{cell}$ or $\mathcal{SH}(k)^{cell+}$ respectively, because these satisfy some useful additional properties. A couple of these are proven in [DI05].

A consequence of the results in [DI05] is the following proposition, which states that cellular homology theories and cohomology theories for $\mathcal{SH}(k)^{fin}$ are exchangeable. We will state the analogous result for $\mathcal{SH}(k)^{fin+}$ below, in Corollary 6.1.4.

Proposition 6.1.1. Let $E \in S\mathcal{H}(k)^{cell}$ (see Definition 5.2.1) be a ring spectrum and $X \in S\mathcal{H}(k)^{fin}$. Then $E_{**}(X) = 0$ if and only if $E^{**}(X) = 0$.

Proof. In the universal coefficient spectral sequence (see [DI05, Proposition 7.7] or Proposition 8.1.1),

$$E_2 = \text{Ext}_{E_{**}}^{a,b,c}(M_{**}, N_{**}) \Rightarrow \pi_{-a-b,-c}F_E(M, N),$$

we set $M = X \wedge E$ and N = E:

$$E_2 = \text{Ext}_{E_{**}}^{a,b,c}(E_{**}(X), E_{**}) \Rightarrow \pi_{-a-b,-c}F_E(X \land E, E).$$

Since $F_E(X \wedge E, E) = F(X, E)$, the spectral sequence converges conditionally to $E^{**}(X)$. If $E_{**}(X) = 0$, the spectral sequence collapses and thus converges strongly to $E^{**}(X)$, which, hence, is 0. For the other direction, we set M = F(X, E) and N = E. Note that this M is cellular because X is dualisable and its dual is again a finite cell spectrum by Proposition 5.2.7. So we get

$$E_2 = \text{Ext}_{E_{**}}^{a,b,c}(E^{**}(X), E_{**}) \Rightarrow \pi_{-a-b,-c}F_E(F(X, E), E).$$

Now $F_E(F(X, E), E) = F_E(D(X) \land E, E) = F(D(X), E) = X \land E$, so the sequence converges conditionally to $E_{**}(X)$. Hence, $E^{**}(X) = 0$ implies $E_{**}(X) = 0$.

One important result of Dugger and Isaksen, [DI05, Proposition 7.1], states that for cellular objects $E \in S\mathcal{H}(k)^{cell}$, $\pi_{**}E = 0$ implies $E \cong 0$. We will show that an adjusted statement holds for $E \in S\mathcal{H}(k)^{cell+}$, $k \subseteq \mathbb{R}$.

We know from equivariant stable homotopy theory that a generalisation of homotopy groups is needed to obtain the corresponding result in $\mathcal{SH}(G)$. In $\mathcal{SH}(\mathbb{Z}/2)$, for example, we have the equivariant homotopy groups

$$\pi_n^{\mathbb{Z}/2}(X) = [S^n, X]_{\mathbb{Z}/2} = [S^n, X^{\mathbb{Z}/2}]_{\{1\}} \text{ and } \pi_n^{\{1\}}(X) = [S^n \wedge \mathbb{Z}/2_+, X]_{\mathbb{Z}/2} = [S^n, X]_{\{1\}}.$$

We can similarly define homotopy groups in $\mathcal{SH}(k)$, $k \subseteq \mathbb{R}$ such that R'_k maps $\pi^k_{p,q}(X)$ to $\pi^{\mathbb{Z}/2}_p(R'_kX)$ and $\pi^{k[i]}_{p,q}(X)$ to $\pi^{\{1\}}_p(R'_kX)$.

Definition 6.1.2. For $k \subseteq \mathbb{R}$ and $X \in \mathcal{SH}(k)$, let

$$\pi_{p,q}^{k}(X) = [S^{p,q}, X]_{\mathcal{SH}(k)} \text{ and } \pi_{p,q}^{k[i]}(X) = [S^{p,q} \land \operatorname{Spec}(k[i])_{+}, X]_{\mathcal{SH}(k)}$$

We write $\pi_{**}^+(X) = 0$ if $\pi_{**}^K(X) = 0$ for both K = k and K = k[i]. Furthermore, $E_{**}^+(X) = 0$ will mean $\pi_{**}^+(E \wedge X) = 0$ and $(E^+)^{**}(X) = 0$ will mean $\pi_{**}^+(F(X, E)) = 0$.

The same arguments as in [DI05, Proposition 7.1] now imply the following.

Proposition 6.1.3. If $X \in S\mathcal{H}(k)^{cell+}$, $k \subseteq \mathbb{R}$ and $\pi^+_{**}(X) = 0$ then $X \cong 0$.

Proof. Assuming X is cofibrant and fibrant, one considers the class \mathcal{D} of all Y such that $\operatorname{Map}(Y^{cof}, X)$ is contractible. By assumption, \mathcal{D} contains $S^{p,q}$ and $S^{p,q} \wedge \operatorname{Spec}(k[i])_+$. Furthermore, \mathcal{D} is closed under isomorphisms, (de-)suspensions and homotopy colimits. As in the proof of Proposition 5.3.7, it follows that it also contains $S^{p,q} \wedge (\operatorname{Spec}(k[i])_+)^{\wedge m}$, $m \geq 0$. As $\mathcal{SH}(k)^{cell+} \subseteq \mathcal{SH}(k)$ is generated by $\{S^{p,q} \wedge (\operatorname{Spec}(k[i])_+)^{\wedge m} \mid p,q \in \mathbb{Z}, m \geq 0\}$ under isomorphisms and homotopy colimits, it follows that $\mathcal{SH}(k)^{cell+} \subseteq \mathcal{D}$, in particular $X \in \mathcal{D}$. Hence, $\operatorname{Map}(X, X)$ is contractible, which implies $X \cong 0$.

The following corollary is a $k \subseteq \mathbb{R}$ -version of Proposition 6.1.1.

Corollary 6.1.4. Let $E \in \mathcal{SH}(k)^{cell+}$ be a ring spectrum and $X \in \mathcal{SH}(k)^{fin+}$. Then $E_{**}^+(X) = 0$ if and only if $(E^+)^{**}(X) = 0$.

Proof. The spectral sequence from [DI05, Proposition 7.7] is derived similarly to the universal coefficient spectral sequence in [Elm+97, Section IV.5]. The crucial point is the convergence, where cellularity is needed to apply [DI05, Proposition 7.1]. The equivariant version of this spectral sequence (a spectral sequence of Mackey functors) is proven in [LM06]. Using [LM06] and Proposition 6.1.3, one can derive the universal coefficient spectral sequence for our motivic homotopy groups over $k \subseteq \mathbb{R}$. The new version of Proposition 6.1.1 can then be deduced from this spectral sequence as before.

6.2 Thick ideals

In $\mathcal{SH}_{(p)}^{fin}$, any thick ideal can be described by some cohomology theory (namely, *n*-th Morava K-theory). Conversely, any cohomology theory defines a thick ideal. We are therefore interested in thick ideals of $\mathcal{SH}(k)_f$, as well as $\mathcal{SH}(k)^{fin}$, $\mathcal{SH}(k)^{fin+}$, described by cohomology theories as follows.

Lemma 6.2.1. Let k be any field and let $\mathcal{T} \subseteq S\mathcal{H}(k)$ be a tensor triangulated subcategory of $S\mathcal{H}(k)$. For any $E \in S\mathcal{H}(k)$, the full subcategory of \mathcal{T} given by

$$\mathcal{C}_E = \{ X \in \mathcal{T} \mid X \land E \cong 0 \}$$

is a thick ideal of \mathcal{T} .

Proof. Since $-\wedge E$ preserves exact triangles, \mathcal{C}_E is closed under these. If Y is a retract of X and $X \wedge E = 0$, then $Y \wedge E \to 0 \to Y \wedge E$ is the identity on $Y \wedge E$, hence $Y \wedge E \cong 0$. Let $X \in \mathcal{C}_E$ and $Y \in \mathcal{T}$. Then $X \wedge Y \wedge E \cong 0$, hence $X \wedge Y \in \mathcal{C}_E$.

Proposition 6.2.2. Let C_E be as defined in the above lemma. If $k \subseteq \mathbb{C}$, $E \in S\mathcal{H}(k)^{cell}$ and $\mathcal{T} = S\mathcal{H}(k)^{fin}$ or $\mathcal{T} = S\mathcal{H}(k)^{cell}$, then

$$\mathcal{C}_E = \{ X \in \mathcal{T} \mid E_{**}(X) = 0 \}$$

If, furthermore, E is a ring spectrum and $\mathcal{T} = \mathcal{SH}(k)^{fin}$, then also

 $\mathcal{C}_E = \{ X \in \mathcal{T} \mid E^{**}(X) = 0 \}.$

If $k \subseteq \mathbb{R}$, $E \in \mathcal{SH}(k)^{cell+}$ and $\mathcal{T} = \mathcal{SH}(k)^{fin+}$ or $\mathcal{T} = \mathcal{SH}(k)^{cell+}$, then

$$\mathcal{C}_E = \{ X \in \mathcal{T} \mid (E^+)_{**}(X) = 0 \}.$$

If, furthermore, E is a ring spectrum and $\mathcal{T} = \mathcal{SH}(k)^{fin+}$, then also

$$\mathcal{C}_E = \{ X \in \mathcal{T} \mid (E^+)^{**}(X) = 0 \}.$$

The same descriptions apply if \mathcal{T} is the p-localisation of any of these categories for some prime p.

Proof. For E and X as in the first claim, we have $E_{**}(X) = 0 \Leftrightarrow E \land X \cong 0$ by [DI05, Proposition 7.1] and for X finite, E a ring spectrum, $E^{**}(X) = 0 \Leftrightarrow E_{**}(X) = 0$ by Proposition 6.1.1. For $k \subseteq \mathbb{R}$, the same arguments hold by Proposition 6.1.3 and Corollary 6.1.4.

6.3 Morava K-theories in $\mathcal{SH}(k)$

Topology suggests that thick ideals described by Morava K-theories are particularly interesting. Therefore, we want to study motivic Morava K-theories and their properties.

6.3.1 Construction and properties of AK(n)

Motivic Morava K-theories were introduced in [Bor03]. The rough idea is as follows. One starts with MGL, the motivic analogue of MU. Since MU_* is a subring of MGL_{**} , elements in MU_* can be used to define maps on MGL and to construct motivic spectra which are analogous to certain other topological spectra constructed from MU.

Definition 6.3.1. Let $k \subseteq \mathbb{C}$. Let MGL be the algebraic cobordism spectrum as constructed in [Voe96, Section 3.5], see also [Voe98, Section 6.3] or [PY02, Section 6.5]. In [Bor03, Theorem 10], elements $a_i \in MGL_{2i,i}$, $i \geq 1$, are defined, whose images under R_k are $a_i^{\text{top}} \in MU_{2i}$. If E is an MGL-module, then E/a_i and $a_i^{-1}E$ can be defined as in [Hor06, Definition 2.10].

Note that the functor $MGL \wedge -$ from $S\mathcal{H}(k)$ to the category of MGL-modules in $S\mathcal{H}(k)$ has F(MGL, -) as right adjoint and has the forgetful functor as left adjoint. As in [Elm+97, Lemma II.1.3] and [Bor03, page 99], it follows that the category of MGL-modules is complete and cocomplete. Thus, E/a_i and $a_i^{-1}E$ are again MGL-modules (see Lemma 9.3.1 for the proof that the action of a_i on E is a map of MGL-modules).

The motivic Brown-Peterson spectrum for a fixed prime p was first constructed in [Vez01, Section 5]. By [Hoy13, Remark 6.20], it is equivalent as an MGL-module to $MGL_{(p)}/I$, where I is the image under $MU_* \to MGL_{**}$ of a regular sequence in the Lazard ring L that generates the vanishing ideal for p-typical formal group laws. We take this as the definition of the motivic Brown Peterson spectrum ABP at the prime p. That is,

$$ABP = MGL_{(p)}/(a_i \mid i \neq p^j - 1).$$

With $v_0 = p$ and $v_i = a_{p^i-1}$ for $i \ge 1$, we define:

$$AP(n) = ABP/(v_0, \dots, v_{n-1}) \text{ and } AB(n) = v_n^{-1}AP(n),$$

$$Ak(n) = ABP/(v_0, \dots, v_{n-1}, v_{n+1}, v_{n+2}, \dots) \text{ and } AK(n) = v_n^{-1}Ak(n),$$

$$AE(n) = v_n^{-1}ABP/(v_{n+1}, v_{n+2}, \dots).$$

We will make use of all these spectra in Chapter 9. For now, we are mostly interested in AK(n).

Lemma 6.3.2. Let Ah be one of the motivic spectra mentioned in the above definition, e.g. Ah = AK(n) with h = K(n). Then $R_k(Ah) = h$.

Proof. This follows from $R_k(MGL) = MU$ [Voe96, Section 3.5] and $R_k(v_i) = v_i^{\text{top}}$ because R_k preserves colimits.

Remark 6.3.3. (1) All spectra mentioned in the above definition are cellular: Theorem 6.4 in [DI05] shows that MGL is cellular. Since the spectra above are constructed from MGL by taking homotopy cofibers and homotopy colimits, they are cellular. The mod-p version of the main result in [Hoy13] states that $H\mathbb{Z}/p$ is the cofiber of $v_n: \Sigma^{2(p^n-1),p^n-1}Ak(n) \to Ak(n)$. Hence, $H\mathbb{Z}/p$ is cellular, too.

- (2) ABP is a homotopy commutative ring spectrum by construction [Vez01, Definition 5.3] and the orientation of MGL induces an orientation on ABP by [PPR08, Theorem 1.1]. The ring structure of ABP induces an ABP-module structure on the quotients of ABP defined above.
- (3) Since MGL is a ring spectrum, ABP is also an $MGL_{(p)}$ -module spectrum and so is ABP/I for $I \subseteq MU_*$ a regular ideal.
- (4) As remarked at the end of the introduction in [NSØ09], ABP is Landweber exact in the sense of [NSØ09] and their results for MGL also hold for ABP.
- (5) MGL_{2*,*} and H_{**} MGL (with coefficients in Z or Z/p) are known (see e.g. [Bor03, Theorem 5] and [Hoy13, Corollary 6.9]). From this, one can compute Ah_{2*,*} and H_{**} Ah if Ah is one of the above spectra ([Bor03, Theorem 12] and [Hoy13, Lemma 6.10]). More can be said about Ah_{**} if H_{**}(Spec k) is known, see Lemma 6.3.7.
- (6) In general, however, Ah_{**} is not known in degrees different from (2i, i). It is also not known whether Ah can be given the structure of a ring spectrum (except for Ah = ABP). Another open question is whether AK(n) satisfies the Künneth formula like K(n).

We will make use of motivic Atiyah-Hirzebruch spectral sequences as discovered by Hopkins and Morel and worked out by Hoyois [Hoy13, Example 8.13]. See also [Lev08, Section 11].

Proposition 6.3.4. Let $h = MU_{(p)}/I$ and $Ah = MGL_{(p)}/I$ for some regular ideal $I \subseteq MU_*$. For $X \in SH(k)_{(p)}^{fin}$, there are strongly convergent spectral sequences:

$$\begin{split} E_2^{p,q,t} &= \mathbf{H}^{p+2t,q+t}(X,h_t) \Rightarrow Ah^{p,q}(X), \\ E_2^{p,q,t} &= \mathbf{H}^{p+2t,q+t}(X,(v_n^{-1}h)_t) \Rightarrow (v_n^{-1}Ah)^{p,q}(X). \end{split}$$

- **Remark 6.3.5.** (1) These are the spectral sequences associated with the slice filtrations of Ah and $v_n^{-1}Ah$. The *n*-th truncation of MGL in the slice filtration is described in the proof of [Spi10, Theorem 4.6] as the colimit of a certain diagram $D_{\deg \geq n}$, meaning that $f_n MGL$ is constructed from MGL by quotienting out all monomials $a_1^{k_1} \cdots a_m^{k_m}$ with $a_i \in MGL_{2*,*}$ and $\sum_{i=1}^m ik_i \geq n$. The same construction with MU instead of MGL yields the Postnikov truncation of MU because $MU_* = \mathbb{Z}[a_1^{\text{top}}, a_2^{\text{top}}, \cdots]$ with $|a_i^{\text{top}}| = 2i$. It follows that, for $k \subseteq \mathbb{C}$, $R_k : S\mathcal{H}(k) \to S\mathcal{H}$ maps the slice filtration of MGL to the Postnikov filtration of MU. By the construction of Ah and $v_n^{-1}Ah$ from MGL, this implies that their slice filtrations also realise to the Postnikov filtrations of h and $v_n^{-1}h$. For the associated spectral sequences, this means that R_k maps the above spectral sequences to the analogous topological Atiyah-Hirzebruch spectral sequences.
- (2) In [Hoy13, Example 8.13], the convergence is stated for $X \in \mathbf{Sm} / S$ and follows from $Ah, v_n^{-1}Ah$ being convergent with respect to $[\Sigma^{0,q}\Sigma^{\infty}X_+, -], X \in \mathbf{Sm} / S$, in the sense of [Hoy13, Section 8.5]. But this implies convergence with respect to [X, -] for all finite cell spectra X by motivic cellular induction. Hence, the sequences converge for all finite cell spectra X. Note that [Hoy13, Example 8.13] shows the convergence for Landweber exact spectra, but his proof holds for quotients of MGL as well.

In Chapters 8 and 9, we will often assume $k = \mathbb{C}$ to be able to prove more results than for general k. The reason is that the coefficients of $H\mathbb{Z}/p$ are particularly simple in this case.

Lemma 6.3.6. For $k = \mathbb{C}$,

$$\mathrm{H}^{**}(\mathrm{Spec}(\mathbb{C}),\mathbb{Z}/p)\cong\mathbb{F}_p[\tau]$$

with $\deg(\tau) = (0, 1)$.

Proof. This is [Voe10, Equation (74)].

This, and the above spectral sequence, can be used to calculate the coefficients of theories like Ak(n).

Lemma 6.3.7. Let $k = \mathbb{C}$ and $h = MU_{(p)}/I$ with $(p) \subseteq I \subseteq MU_*$ a regular ideal. Let $Ah = MGL_{(p)}/I$. Then

$$Ah_{**} \cong \operatorname{H}_{**}(\operatorname{Spec} \mathbb{C}, \mathbb{F}_p) \otimes_{\mathbb{F}_p} h_* \cong h_*[\tau].$$

Proof. This is remarked in [Yag05], below Corollary 3.9. The reason is the following: For $X = \text{Spec } \mathbb{C}$, the motivic Atiyah-Hirzebruch spectral sequence is

$$\mathrm{H}^{p+2t,q+t}(\operatorname{Spec}\mathbb{C},\mathbb{F}_p)\otimes h_t \Rightarrow Ah^{p,q}(\operatorname{Spec}\mathbb{C}).$$

By Lemma 6.3.6, $\mathrm{H}^{**}(\operatorname{Spec} \mathbb{C}, \mathbb{F}_p) \cong \mathbb{F}_p[\tau]$ with $\deg(\tau) = (0, 1)$. Thus, for fixed t, $E_2^{p,q,t} = \mathrm{H}^{p+2t,q+t}(X, h_t)$ can only be nonzero in the column p = -2t, which implies that all differentials vanish. Therefore, the spectral sequence collapses immediately, proving $Ah_{**} \cong \mathrm{H}_{**} \otimes h_*$.

6.3.2 Thick ideals and Morava K-theories

Let K be a cellular spectrum in $\mathcal{SH}(k)$, $k \subseteq \mathbb{C}$, such that $R_k(K) = K(n)$ is the n-th Morava K-theory with respect to a fixed prime p. We call such a K a motivic model for K(n).

Let $C_K = \{X \in (\mathcal{SH}(k)_f)_{(p)} | K \wedge X \cong 0\}$, which is a thick ideal in the *p*-localised category $(\mathcal{SH}(k)_f)_{(p)}$ by Lemma 6.2.1. If $K \wedge X \cong 0$, then $R_k(K \wedge X) \cong K(n) \wedge R_k(X) \cong 0$. Consequently, $R_k(X) \in C_{n+1}$ for any $X \in C_K$. This is content of the following proposition.

Proposition 6.3.8. For K a motivic model for K(n), we have an inclusion of thick ideals in $(\mathcal{SH}(k)_f)_{(p)}$:

$$\mathcal{C}_K \subseteq R_k^{-1}(\mathcal{C}_{n+1}).$$

The same is true in the category $\mathcal{SH}(k)^{fin}_{(p)}$.

Example 6.3.9. The motivic Morava K-theory spectra AK(n), as defined in Definition 6.3.1, satisfy $R_k(AK(n)) = K(n)$ and are cellular. In particular, they satisfy the previous proposition.

Another possibility for such a spectrum K is the constant Morava K-theory spectrum $c_k(K(n))$. In Sections 8.4 - 8.7 we will have a closer look at the thick ideals $\mathcal{C}_{AK(n)}$ and $\mathcal{C}_{c_kK(n)}$ for $k = \mathbb{C}$.

Remark 6.3.10. The spectrum K(n) is not finite. One possibility to see this is by the equivalence [Rav84, Theorem 2.1.(h) and (i)]:

$$K(m) \wedge K(n) \cong 0 \Leftrightarrow m \neq n$$

If K(n) were finite, we would have $K(m)_*K(n) = 0 \Rightarrow K(m-1)_*K(n) = 0$ by [Rav84, Theorem 2.11], which is wrong for n = m - 1. As a consequence, any spectrum K with $R_k(K) = K(n)$ can also not be finite cellular.

Remark 6.3.11. Another corollary of the above equivalence and of [Rav84, Theorem 2.1.(h)] is the following: If K and K' are motivic models for K(n), then $R_k(K \wedge K') = K(n) \wedge K(n)$, which has the same Bousfield class as K(n). In particular, any $X \in \mathcal{C}_{K \wedge K'}$ satisfies $R_k(X) \in \mathcal{C}_{n+1} \setminus \mathcal{C}_{n+2}$. Hence,

$$\mathcal{C}_{K \wedge K'} \subseteq R_k^{-1}(\mathcal{C}_{n+1}) \text{ and } \mathcal{C}_{K \wedge K'} \nsubseteq R_k^{-1}(\mathcal{C}_{n+2}).$$

Chapter 7

$\mathcal{SH}(k)_f$ has more thick ideals than \mathcal{SH}^{fin}

7.1 The motivic Hopf map

In this chapter, we study the cofiber of the motivic Hopf map, which generates a thick ideal that is not of the form $R_k^{-1}(\mathcal{C}_n)$ or $\mathcal{C}_{AK(n)}$. The first part of this claim was proven by Balmer in [Bal10, Proposition 10.4]. We will reprove and extend this result, as well as summarise and apply other results of Balmer's work on prime ideals in tensor triangulated categories.

For $k \subseteq \mathbb{R}$, we have already shown that $\mathcal{SH}(k)_f$ has more thick ideals than \mathcal{SH}^{fin} . This chapter is, therefore, in particular interesting for $k = \mathbb{C}$.

To understand why there cannot be a complete analogy between motivic thick ideals and topological thick ideals, recall the reasoning in the topological case: the thick subcategory (i.e., thick ideal) theorem of Hopkins and Smith [HS98] is derived from the fact that Morava K-theories detect nilpotence, which follows from the theorem that MUdetects nilpotence. This is not the case in the motivic setting.

The Hopf map in $\mathcal{SH}(k)$ is given by $\eta : \mathbb{A}^2 \setminus \{0\} \to \mathbb{P}^1$, $(x, y) \mapsto [x : y]$, defining an element $\eta \in \pi_{1,1}(S)$. Unlike the topological Hopf element, η is not nilpotent [Mor06, Theorem 4.7]. The following lemma is due to Morel, see e.g. [Mor02], but we could not find a proof in the literature. It shows that the spectrum MGL does not detect the non-nilpotence of η .

Lemma 7.1.1.

$$MGL \wedge \eta \cong 0$$

Proof. The unit map u of the ring spectrum MGL factors through $\Sigma^{-2,-1}C\eta$, as in the

proof of [Hoy13, Theorem 3.8]. Consider the diagram:



The upper row is a cofiber sequence and u, m are the structure maps of the ring spectrum MGL. The diagram is commutative. It follows that $1 \wedge \Sigma^{-2,-1}\eta$ factors through its own cofiber, hence it must be zero. Suspending by $S^{2,1}$, we get $MGL \wedge \eta \cong 0$.

It seems likely that also $AK(n) \wedge \eta \cong 0$, but this does not immediately follow from the above lemma. For our interests, it will suffice to know that $AK(n) \wedge C\eta \cong 0$, which we prove differently.

Remark 7.1.2. By [Mor03, Lemma 6.2.1], $C\eta \cong \mathbb{P}^2$.

Proposition 7.1.3. For any $k \subseteq \mathbb{C}$ and any prime p,

thickid
$$(R_k(C\eta_{(p)})) = \mathcal{C}_0 = \mathcal{SH}_{(p)}^{fin}.$$

For $k \subseteq \mathbb{R}$,

thickid
$$(R'_k(C\eta_{(p)})) = \begin{cases} \mathcal{C}_{0,1} \subset (\mathcal{SH}(\mathbb{Z}/2)_f)_{(2)} & \text{if } p = 2\\ \mathcal{C}_{0,\infty} \subset (\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)} & \text{if } p \neq 2, \end{cases}$$

where C_n is the thick ideal defined in Theorem 2.0.10 and $C_{m,n}$ is the thick ideal defined in Corollary 3.5.1.

Proof. Since, by definition, $R_k = R_{\mathbb{C}} \circ f^*$ for $f: k \to \mathbb{C}$, $R'_k = R_{\mathbb{R}} \circ f^*$ for $f: k \to \mathbb{R}$, and $f^*(\eta_k) = \eta_K$ for $f: k \to K$, it suffices to prove the claims for $k = \mathbb{C}$ and $k = \mathbb{R}$. For $k = \mathbb{C}$, we have $R(C\eta) = R(\mathbb{P}^2) = \mathbb{C}P^2$. Since $K(0)_*(\mathbb{C}P^2_{(p)}) = H_*(\mathbb{C}P^2_{(p)}, \mathbb{Q}) \neq 0$,

For $k = \mathbb{C}$, we have $R(C\eta) = R(\mathbb{P}^2) = \mathbb{C}P^2$. Since $K(0)_*(\mathbb{C}P_{(p)}^2) = \mathrm{H}_*(\mathbb{C}P_{(p)}^2, \mathbb{Q}) \neq 0$, $R(C\eta_{(p)})$ has type 0. By [HS98, Theorem 7], any spectrum of type 0 generates $\mathcal{SH}_{(p)}^{fin}$. So, thickid $(R(C\eta_{(p)})) = \mathcal{SH}_{(p)}^{fin}$.

So, thickid $(R(C\eta_{(p)})) = \mathcal{SH}_{(p)}^{fin}$. For $k = \mathbb{R}$, $R'(C\eta)^{\{1\}} = C\eta(\mathbb{C}) = \mathbb{C}P^2$ as before. Therefore, if $R'(C\eta_{(p)}) \in \mathcal{C}_{m,n}$, then m = 0.

Furthermore, $R'(\eta)^{\mathbb{Z}/2} = \eta(\mathbb{R})$ is the quotient map $\mathbb{R}^2 \setminus \{0\} \to \mathbb{R}P^1$, which is isomorphic to $2: S^1 \to S^1$. It follows that $R'(C\eta)^{\mathbb{Z}/2} = S/2$.

If $p \neq 2$, $S \xrightarrow{2} S$ is an isomorphism in $\mathcal{SH}_{(p)}^{fin}$ and, hence, $(S/2)_{(p)} = 0$, which has type ∞ . This proves thickid $(R'(C\eta_{(p)})) = \mathcal{C}_{0,\infty}$ if $p \neq 2$.

Now let p = 2. Since $S \xrightarrow{2} S$ is an isomorphism rationally, we have $K(0)_*(S/2) = H_*(S/2, \mathbb{Q}) = 0$. On the other hand, K(1) is a direct summand of mod 2 topological K-theory, so $K(1)_*(S \xrightarrow{2} S) = 0$ and $K(1)_*(S/2) \neq 0$, as in the proof of [Bal10, Proposition 9.4]. Hence, the type of $S/2 \in S\mathcal{H}_{(2)}^{fin}$ is 1 and, therefore, thickid $(R'_k((C\eta)_{(2)})) = \mathcal{C}_{0,1}$.

In terms of the motivic thick ideals thickid $(C\eta_{(p)}) \subseteq (\mathcal{SH}(k)_f)_{(p)}$, the above proposition states the following: For $k \subseteq \mathbb{C}$, thickid $(C\eta_{(p)}) \subseteq R_k^{-1}(\mathcal{C}_0)$ (which is a trivial statement) and thickid $(C\eta_{(p)}) \not\subseteq R_k^{-1}(\mathcal{C}_n)$ for n > 0. For $k \subseteq \mathbb{R}$ and $p \neq 2$, thickid $(C\eta_{(p)}) \subseteq (R'_k)^{-1}(\mathcal{C}_{0,\infty})$ and thickid $(C\eta_{(p)}) \not\subseteq (R'_k)^{-1}(\mathcal{C}_{m,n})$ for any m > 0, n arbitrary. If p = 2, then thickid $((C\eta)_{(2)}) \subseteq (R'_k)^{-1}(\mathcal{C}_{0,1})$ and thickid $((C\eta)_{(2)}) \not\subseteq (R'_k)^{-1}(\mathcal{C}_{m,n})$ for any m > 0 or n > 1.

The next proposition contains two more results on thickid $(C\eta_{(p)}) \subseteq (\mathcal{SH}(k)_f)_{(p)}$.

Proposition 7.1.4. For $k \subseteq \mathbb{C}$, let thickid $(C\eta_{(p)}) \subseteq (\mathcal{SH}(k)_f)_{(p)}$ denote the thick ideal generated by the p-localised cofiber of the Hopf map. Then the following hold:

- (1) thickid $(C\eta_{(p)}) \not\subseteq \mathcal{C}_{AK(n)}$ for any $n \ge 0$ and any prime p,
- (2) thickid $(C\eta_{(p)}) \not\subseteq R_k^{-1}(\mathcal{C}_n)$ for any n > 0 and any prime p,
- (3) thickid $(C\eta_{(p)}) \subseteq \operatorname{thickid}(S_{(p)}^0) = (\mathcal{SH}(k)_f)_{(p)}$ if $k \subseteq \mathbb{R}$ and p is any prime or $k \subseteq \mathbb{C}$ and p = 2.
- (4) For any prime p, the thick ideals thickid $(C\eta_{(p)}) \cap R_k^{-1}(\mathcal{C}_n)$ are distinct for different $n \ge 0$ and in particular nonzero if $n < \infty$.
- Proof. (1) By Proposition 6.3.8, $\mathcal{C}_{AK(n)} \subseteq R_k^{-1}(\mathcal{C}_{n+1})$. Assuming thickid $(C\eta_{(p)}) \subseteq \mathcal{C}_{AK(n)}$ therefore implies thickid $(C\eta_{(p)}) \subseteq R_k^{-1}(\mathcal{C}_{n+1})$. Hence, (1) will follow from (2).
- (2) This can either be derived from the previous proposition or can be seen by the following argument.

 $R_k(\eta)$ is the topological Hopf map, which is nilpotent. Hence, for $n \ge 1$, $R_k(\eta)_*$ is not surjective in the sequence

$$\cdots \to K(n-1)_*(S^3) \xrightarrow{R_k(\eta)_*} K(n-1)_*(S^2) \to K(n-1)_*(C(R_k\eta)) \to \cdots$$

It follows that $K(n-1)_*(C(R_k\eta)) \neq 0$ and, since $C(R_k\eta) \cong R_k(C\eta), C\eta_{(p)} \notin R_k^{-1}(\mathcal{C}_n).$

(3) Note that Cη ∈ thickid(S⁰) = SH(k)_f, so thickid(Cη_(p)) ⊆ thickid(S⁰_(p)) for any p. We have to show that S⁰_(p) ∉ thickid(Cη_(p)). We consider the sheaf cohomology theory H*(-, K^{MW}_{*}[η⁻¹]), on which η induces an isomorphism as in the proof of [Mor06, Theorem 4.7], where Morel concludes that η cannot be nilpotent. Localising at p, we get that η_(p) induces an isomorphism on H*(S⁰_(p), K^{MW}_{*}[η⁻¹]). Consider H⁰(S⁰_(p), K^{MW}_{*}[η⁻¹]) = K^{MW}_{*}(k)[η⁻¹]_(p). As in the proof of [Mor06, Theorem 4.7], this is K^W₀(k)[η, η⁻¹]_(p), and K^W₀(k) is isomorphic to the Witt ring W(k) by [Mor06, Remark 4.2]. We have W(C) = Z/2 and W(R) = Z, see e.g. [Lam05, p. 34]. Hence, for k = C, H⁰(S⁰_(p), K^{MW}_{*}[η⁻¹]) = (Z[η, η⁻¹])_(p) ≠ 0 for any p. Therefore, the above mentioned isomorphism induced by η_(p) is not the zero map, and, as in [Mor06, Theorem 4.7], it follows that η_(p) is not nilpotent in these cases. Now we apply [Bal10, Theorem 2.15], which states that, for a map f : X → Y between invertible objects X and Y in a tensor triangulated category T, the thick

ideal generated by Cf is equal to the full subcategory consisting of all objects A such that $f^{\wedge n} \wedge A = 0$ for some $n \geq 1$ (this is similar to Proposition 3.3.7). A corollary of this theorem is that $\operatorname{thickid}(Cf) = \mathcal{T}$ if and only if f is nilpotent.

It follows that, since $\eta_{(p)}$ is not nilpotent, $S^0_{(p)} \notin \text{thickid}(C\eta_{(p)})$ and $\text{thickid}(C\eta_{(p)}) \neq \text{thickid}(S^0_{(p)})$ in the cases $k = \mathbb{C}$ and p = 2 or $k = \mathbb{R}$ and p any prime.

Now, let $f: k \to \mathbb{C}$. By Proposition 5.1.1, $(f^*)^{-1}$ preserves thick ideals. Since $f^*(C\eta_k) = C\eta_{\mathbb{C}}$, thickid $((C\eta_k)_{(2)}) \subseteq (f^*)^{-1}$ (thickid $((C\eta_{\mathbb{C}})_{(2)})$). As $f^*((S^0_k)_{(2)}) = (S^0_{\mathbb{C}})_{(2)} \notin$ thickid $((C\eta_{\mathbb{C}})_{(2)})$, it follows $(S^0_k)_{(2)} \notin (f^*)^{-1}$ (thickid $((C\eta_{\mathbb{C}})_{(2)})$) and, hence, $(S^0_k)_{(2)} \notin$ thickid $((C\eta_k)_{(2)})$.

The same argument holds for arbitrary primes p if $f : k \hookrightarrow \mathbb{R}$. Alternatively, the statement for $k \subseteq \mathbb{R}$ can be derived from Proposition 7.1.3 and Theorem 5.3.1:

$$\operatorname{thickid}(C\eta_{(p)}) \subseteq (R'_k)^{-1}(\operatorname{thickid}(R'_k(C\eta_{(p)}))) \neq (R'_k)^{-1}(\mathcal{C}_{0,0}) = \operatorname{thickid}(S^0_{(p)}).$$

(4) Let $X \in R_k^{-1}(\mathcal{C}_n)$ be such that $R_k(X)$ is of type n, i.e., $R_k(X) \in \mathcal{C}_n \setminus \mathcal{C}_{n+1}$. From the proof of (2), we know that $R_k(C\eta_{(p)})$ is of type 0. From the Künneth formulas for K(n) and K(n+1), it follows that $R_k(C\eta_{(p)} \wedge X) \cong R_k(C\eta_{(p)}) \wedge R_k(X)$ is of type n. Therefore,

$$C\eta_{(p)} \wedge X \in \left(\operatorname{thickid}(C\eta_{(p)}) \cap R_k^{-1}(\mathcal{C}_n)\right) \setminus \left(\operatorname{thickid}(C\eta_{(p)}) \cap R_k^{-1}(\mathcal{C}_{n+1})\right)$$

Corollary 7.1.5. For $k \subseteq \mathbb{C}$, thickid $((C\eta)_{(2)})$ is neither of the form $\mathcal{C}_{AK(n)}$ for any $n \geq 0$ (by Proposition 7.1.4(1)) nor of the form $R_k^{-1}(\mathcal{C}_n)$ for any $n \geq 0$ (by Proposition 7.1.4(2) and (3)).

For $k \subseteq \mathbb{R}$, the statements analogous to (2) and (4) of Proposition 7.1.4 read as follows.

Proposition 7.1.6. Let η denote the Hopf map in SH(k), $k \subseteq \mathbb{R}$. In $(SH(k)_f)_{(p)}$ the following inequalities hold.

- (1) $\operatorname{thickid}(C\eta_{(p)}) \not\subseteq (R'_k)^{-1}(\mathcal{C}_{m,n})$ for any m > 0 (p, n arbitrary) or, if p = 2, m = 0and n > 1.
- (2) Let (m, n) and (m', n') be pairs of integers ≥ 0 such that $\mathbb{Z}/2$ -equivariant spectra of types (m, n) and (m', n') exist (see Chapter 3). If p is any prime and $m \neq m'$ or p = 2 and $\max(n, 1) \neq \max(n', 1)$ then

thickid
$$(C\eta_{(p)}) \cap (R'_k)^{-1}(\mathcal{C}_{m,n}) \neq \text{thickid}(C\eta_{(p)}) \cap (R'_k)^{-1}(\mathcal{C}_{m',n'}).$$

Otherwise, the two thick ideals are equal.

Proof. (1) is a reformulation of the second part of Proposition 7.1.3. For (2), take $X = c'_k(X_{m,n})$ or any other spectrum whose realisation is a spectrum of type (m, n). Then $R'_k(C\eta_{(p)} \wedge X) \cong R'_k(C\eta_{(p)}) \wedge R'_k(X)$ is the smash product of a spectrum of type $(0, \infty)$ if p is odd ((0, 1) if p = 2) with a spectrum of type (m, n). By the equivariant

Künneth formula, Corollary 3.2.2, it follows that $R'_k(C\eta_{(p)} \wedge X)$ has type (m, ∞) if p is odd and $(m, \max(n, 1))$ if p = 2. Consequently, thickid $(C\eta_{(p)}) \cap (R'_k)^{-1}(\mathcal{C}_{m,n})$ is equal to thickid $(C\eta_{(p)}) \cap (R'_k)^{-1}(\mathcal{C}_{m,\infty})$ if p is odd and to thickid $(\check{C}\eta_{(p)}) \cap (R'_k)^{-1}(\mathcal{C}_{m,\max(n,1)})$ if p=2. The intersection is not contained in $(R'_k)^{-1}(\mathcal{C}_{m',n'})$ for any m' > m or, if p=2, for any $n' > \max(n, 1)$.

Related to the previous propositions are the following conjectures.

Conjecture 7.1.7. (1) $R_k^{-1}(\mathcal{C}_n) \not\subseteq \text{thickid}(C\eta_{(p)})$ for all $0 \leq n < \infty$. (2) $\mathcal{C}_{AK(n)} \not\subseteq \text{thickid}(C\eta_{(p)})$ for all $n \geq 0$.

- (3) The thick ideals thickid $(C\eta_{(p)}) \cap \mathcal{C}_{AK(n)}$ are distinct for different $n \geq 0$ and in particular nonzero if $n < \infty$.
- (4) For $k \subseteq \mathbb{R}$, thickid $(C\eta_{(p)}) \subsetneq (R'_k)^{-1}(\mathcal{C}_{0,\infty})$ for odd primes p and, for p = 2, thickid $(C\eta_{(p)}) \subsetneq (R'_k)^{-1}(\mathcal{C}_{0,1}).$ (5) For $k \subseteq \mathbb{R}$, $(R'_k)^{-1}(\mathcal{C}_{m,n}) \not\subseteq$ thickid $(C\eta_{(p)})$ for all $0 \le m, n < \infty$.

Remark 7.1.8. A possible approach for proving (1) or (2) might be to choose a motivic spectrum X whose realisation is of type n (e.g. $c_k(X_n)$) or a spectrum which is of motivic type n (e.g. as constructed in Chapter 8), respectively. If one can show that $\mathrm{H}^{*}(X, \underline{K}^{\tilde{M}\tilde{W}}_{*}[\eta^{-1}]) \neq 0$, this proves (1) respectively (2).

An idea to prove (3) is to choose a spectrum X of motivic type n and to show that $C\eta \wedge X$ is again of motivic type n for this particular choice.

For (4), p odd, one has to find a spectrum $X \in (R'_k)^{-1}(\mathcal{C}_{0,\infty})$, which is not contained in thickid $(C\eta_{(p)})$. $X = (\Sigma^{\infty} \operatorname{Spec} \mathbb{C}_+)_{(p)}$ satisfies the first condition and it might be possible to show $\mathrm{H}^*(X, \underline{K}^{MW}_*[\eta^{-1}]) \neq 0$, which would imply the second. Since $\mathcal{C}_{0,\infty} \subseteq$ $C_{0,1}$, the same X could be used for the case p = 2.

Remark 7.1.9. If and only if the first conjecture is true, the ideals in Proposition 7.1.4(4) are different from the ideals $R_k^{-1}(\mathcal{C}_n)$ for all $n \geq 0$.

7.2Prime ideals

Let us recall some results of [Bal10], where Balmer studies prime ideals in tensor triangulated categories.

Definition 7.2.1. A prime ideal is a proper thick ideal \mathcal{C} with the additional property that $X \wedge Y \in \mathcal{C}$ implies $X \in \mathcal{C}$ or $Y \in \mathcal{C}$. The set of prime ideals of a tensor triangulated category \mathcal{T} is denoted by $\operatorname{Spc}(\mathcal{T})$.

In [Bal10], the endomorphism ring of the unit object of \mathcal{T} is denoted by $R_{\mathcal{T}}$. To avoid confusion with the realisation functors, we use the notation $\pi_0^{\mathcal{T}}$ instead of $R_{\mathcal{T}}$. In [Bal10, Corollary 5.6], Balmer defines a functor

$$\rho_{\mathcal{T}} : \operatorname{Spc}(\mathcal{T}) \to \operatorname{Spc}(\pi_0^{\mathcal{T}}),$$
$$\rho_{\mathcal{T}}(\mathcal{P}) = \{ f \in \pi_0^{\mathcal{T}} \mid C(f) \notin \mathcal{P} \},$$

which he proves to be surjective if \mathcal{T} is connective [Bal10, Theorem 7.13], where connectivity means that $\operatorname{Hom}_{\mathcal{T}}(S, \Sigma^i S) = 0$ for all i > 0 and S the unit object of \mathcal{T} . Furthermore, this functor is natural for tensor triangulated functors $f : \mathcal{T} \to \mathcal{T}'$ by [Bal10, Theorem 5.3(c) and Corollary 5.6(b)].

7.2.1 Prime ideals in the topological categories \mathcal{SH}^{fin} and $\mathcal{SH}(\mathbb{Z}/2)_f$

Applied to $\mathcal{SH}_{(p)}^{fin}$, this yields the following [Bal10, Proposition 9.4]:

All proper thick ideals $C_n \subset S\mathcal{H}^{fin}_{(p)}$, $0 < n \leq \infty$ are prime ideals, as can be seen either from the Künneth formula for K(n) or by the linear ordering of the C_n . The functor

$$\rho_{\mathcal{SH}_{(p)}^{fin}} : \operatorname{Spc}(\mathcal{SH}_{(p)}^{fin}) \to \operatorname{Spec}(\mathbb{Z}_{(p)})$$

has the following values:

$$\rho_{\mathcal{SH}_{(p)}^{fin}}(\mathcal{C}_n) = \begin{cases} p\mathbb{Z}_{(p)} & \text{if } n > 1, \\ 0 & \text{if } n = 1. \end{cases}$$

Note that C_0 is not proper, so it is not in $\operatorname{Spc}(\mathcal{SH}_{(p)}^{fin})$. Note also that [Bal10] uses a different indexing convention, in which the indices are shifted by one.

The prime ideals in the non-localised category \mathcal{SH}^{fin} are given as follows.

Proposition 7.2.2. A subcategory $C \subset S\mathcal{H}^{fin}$ is a prime ideal if and only if there exist a prime p and a number $1 \leq n \leq \infty$ such that

$$\mathcal{C} = \mathcal{C}_{p,n} = \{ X \in \mathcal{SH}^{fin} \mid K(p, n-1)_* X = 0 \}$$

 $\textit{if } n < \infty, \textit{ or }$

$$\mathcal{C} = \mathcal{C}_{p,\infty} = \{ X \in \mathcal{SH}^{fin} \mid X_{(p)} = 0 \}$$

if $n = \infty$. Here, $C_{p,1} = C_{q,1} = C_1$ for any primes p and q. Except for n = 1, the $C_{p,n}$ are pairwise different.

The functor $\rho_{\mathcal{SH}^{fin}}$: $\operatorname{Spc}(\mathcal{SH}^{fin}) \to \operatorname{Spec}(\mathbb{Z})$ maps $\mathcal{C}_{p,n}$ to $p\mathbb{Z}$ for any n > 1 and it maps \mathcal{C}_1 to 0.

Proof. This is [Bal10, Corollary 9.5]. We give another proof of the first statement. By Theorem 3.4.5, any thick ideal of \mathcal{SH}^{fin} is an intersection

$$\mathcal{C} = \bigcap_{p \in P} \{ X \in \mathcal{SH}^{fin} \mid X_{(p)} \in \mathcal{C}_{n_p} \subseteq \mathcal{SH}^{fin}_{(p)} \}$$

for some set of primes P and some numbers $1 \leq n_p \leq \infty$. If |P| = 0, then $\mathcal{C} = \mathcal{SH}^{fin}$, which is not proper, and, thus, is no prime ideal. Assume $|P| \geq 1$. By Proposition 3.3.8, the intersection of two thick ideals \mathcal{I} and \mathcal{J} satisfying $\mathcal{I} \not\subseteq \mathcal{J}$ and $\mathcal{J} \not\subseteq \mathcal{I}$ is never a prime ideal, because it contains all $X \wedge Y$ with $X \in \mathcal{I}$ and $Y \in \mathcal{J}$. It follows that, if \mathcal{C} is a prime ideal, then |P| = 1. This proves that any prime ideal of \mathcal{SH}^{fin} is of the form $\mathcal{C} = \{X \in \mathcal{SH}^{fin} \mid X_{(p)} \in \mathcal{C}_{n_p}\}$ for some prime p and some $1 \leq n_p \leq \infty$.

On the other hand, the Künneth formula implies that $K(p, n)_*(X \wedge Y)$ can only be zero if $K(p, n)_*X = 0$ or $K(p, n)_*Y = 0$, proving that $\mathcal{C}_{p,n}$ as above is indeed a prime ideal. Note that, for $n = \infty$, we have $X \in \mathcal{C}_{p,n}$ if and only if $K(m)_*(X) = 0$ for all $m \ge 0$, so $\mathcal{C}_{p,\infty}$ is also a prime ideal by the Künneth formula (using that $K(m+1)_*(X) = 0$ implies $K(m)_*(X) = 0$).

Now we turn to the $\mathbb{Z}/2$ -equivariant category, $\mathcal{T} = \mathcal{SH}(\mathbb{Z}/2)_f$. By [Seg70, Corollary 1], the map

$$\pi_0^{\mathcal{T}} \to A(\mathbb{Z}/2), \ [f] \mapsto (\deg(f^{\{1\}}), \deg(f^{\mathbb{Z}/2}))$$

is an isomorphism to the Burnside ring $A(\mathbb{Z}/2) \cong \mathbb{Z} \oplus \mathbb{Z}$. Hence, [Bal10, Theorem 7.13] yields surjective functors

$$\rho_{\mathcal{SH}(\mathbb{Z}/2)_f} : \operatorname{Spc}(\mathcal{SH}(\mathbb{Z}/2)_f) \to \operatorname{Spec}(\mathbb{Z} \oplus \mathbb{Z})$$

and

$$\rho_{(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}} : \operatorname{Spc}((\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}) \to \operatorname{Spec}(\mathbb{Z}_{(p)} \oplus \mathbb{Z}_{(p)})$$

Proposition 7.2.3. A thick ideal $C_{m,n} \subseteq (\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ is a prime ideal if and only if $C_{m,n} = C_{m,0}, \ 0 < m \leq \infty, \text{ or } C_{m,n} = C_{0,n}, \ 0 < n \leq \infty.$

Proof. From Section 3.5, we know that any thick ideal in $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ is of the form

$$\mathcal{C}_{m,n} = \{X \mid \phi^{\{1\}}(X) \in \mathcal{C}_m \text{ and } \phi^{\mathbb{Z}/2}(X) \in \mathcal{C}_n\} = \mathcal{C}_{m,0} \cap \mathcal{C}_{0,n}.$$

By Proposition 3.3.8, this is equal to $\{X \land Y \mid X \in \mathcal{C}_{m,0}, Y \in \mathcal{C}_{0,n}\}$. It follows that $\mathcal{C}_{m,n}$ being a prime ideal implies $\mathcal{C}_{m,n} = \mathcal{C}_{m,0}$ or $\mathcal{C}_{m,n} = \mathcal{C}_{0,n}$. On the other hand, if either m or n is 0 then $\mathcal{C}_{m,n}$ is a prime ideal by the Künneth formula for $K(m, \{1\})$ or $K(n, \mathbb{Z}/2)$. Note that $\mathcal{C}_{0,0} = (\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ is not a prime ideal since it is no proper subcategory.

Remark 7.2.4. As a consequence of Strickland's results [Str10], $C_{m,n} = C_{0,n}$ implies m = 0 but $C_{m,n} = C_{m,0}$ does not always imply that n = 0. This follows from Corollary 3.5.3: for $m \le n$, a type (m, n)-spectrum $X_{m,n}$ always exists, and $X_{0,n} \in C_{0,n} \setminus C_{m,n}$ for all m > 0. On the other hand, Proposition 3.4.7 implies that, for p = 2, $C_{m,0} = C_{m,n}$ for any $0 \le n \le m - 1$.

Recall from Definition 3.5.2 that the thick ideals in $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ are in bijection with a lattice $\Gamma_p \subseteq (\mathbb{Z}_{\geq 0} \cup \{\infty\}) \times (\mathbb{Z}_{\geq 0} \cup \{\infty\})$ containing all (m, n) such that a spectrum of type (m, n) exists. In the case just considered, (m, 0) would not be in Γ_2 (at least for m > 1). In terms of the bijection to Γ_p , the above proposition instead reads as follows:

Corollary 7.2.5. For any $(m,n) \in \Gamma_p$, the thick ideal $\mathcal{C}_{m,n}$ is a prime ideal if and only if one of the following conditions holds: (1) (m,n) = (0,n), with $0 < n \le \infty$, (2) $0 < m \le \infty, \ 0 \le n \le \infty$ and $(m, n-1) \notin \Gamma_p$.

For evaluating the functor $\rho_{(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}}$, we use the first description of the prime ideals in $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$, given by Proposition 7.2.3.

Proposition 7.2.6. The functor

$$\rho_{(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}} : \operatorname{Spc}((\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}) \to \operatorname{Spec}(\mathbb{Z}_{(p)} \oplus \mathbb{Z}_{(p)})$$

has the following values:

$$\rho_{(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}}(\mathcal{C}_{m,0}) = \begin{cases} 0 \oplus \mathbb{Z}_{(p)} & \text{if } m = 1\\ p\mathbb{Z}_{(p)} \oplus \mathbb{Z}_{(p)} & \text{if } m > 1, \end{cases}$$
$$\rho_{(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}}(\mathcal{C}_{0,n}) = \begin{cases} \mathbb{Z}_{(p)} \oplus 0 & \text{if } n = 1\\ \mathbb{Z}_{(p)} \oplus p\mathbb{Z}_{(p)} & \text{if } n > 1. \end{cases}$$

Proof. By [Bal10, Corollary 5.6(b)], $\rho_{\mathcal{T}}$ is natural in \mathcal{T} . Since $X \in \mathcal{C}_{m,0}$ is only a condition on $\phi^{\{1\}}X$ and $X \in \mathcal{C}_{0,n}$ is only a condition on $\phi^{\mathbb{Z}/2}X$, the fixed point functors can be used to derive this result from the nonequivariant case. Note that the induced functor between the endomorphism rings,

$$\pi_0(\phi^H): \pi_0^{(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}} \to \pi_0^{\mathcal{SH}_{(p)}^{fin}}$$

is the projection onto the first summand of $\mathbb{Z} \oplus \mathbb{Z}$ if $H = \{1\}$ and onto the second summand of $\mathbb{Z} \oplus \mathbb{Z}$ if $H = \mathbb{Z}/2$ [Seg70, Corollary 1].

The generalisation to the non-localised category $\mathcal{SH}(\mathbb{Z}/2)_f$ works similarly as in the non-equivariant case. In particular, any prime ideal in $\mathcal{SH}(\mathbb{Z}/2)_f$ is the preimage under the *p*-localisation functor of a prime ideal in $(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}$ for some prime *p*.

Summarising the result, we can say that all information on thick ideals in $\mathcal{SH}(\mathbb{Z}/2)_f$ given in [Bal10]—namely, that they map surjectively to $\operatorname{Spec}(\mathbb{Z} \oplus \mathbb{Z})$ —is recovered in the results of [Str10] as presented in Chapter 3. Furthermore, [Str10] does not only specify a preimage of any element in $\operatorname{Spec}(\mathbb{Z} \oplus \mathbb{Z})$ but gives a complete list of all possible such preimages.

7.2.2 Prime ideals in the motivic category $SH(k)_f$

[Bal10, Section 10] studies $\mathcal{T} = \mathcal{SH}(F)_f$ for F a perfect field, in which case the map $\rho_{\mathcal{T}} : \operatorname{Spc}(\mathcal{T}) \to \operatorname{Spec}(GW(F))$ is surjective. That is, $\operatorname{Spec}(GW(F))$ gives a lower bound on the thick ideals of $\mathcal{SH}(F)_f$.

For $F = \mathbb{C}$, the naturality of $\rho_{\mathcal{T}}$ yields a commutative diagram:

$$\begin{array}{c|c} \operatorname{Spc}(\mathcal{SH}(\mathbb{C})_f) & \xrightarrow{\rho} \operatorname{Spc}(\pi_0^{\mathcal{SH}(\mathbb{C})_f}) \\ & & & & & \\ \operatorname{Spc}(R) & & & & \\ & & & & \\ \operatorname{Spc}(\mathcal{SH}^{fin}) & \xrightarrow{\rho} \operatorname{Spcc}(\pi_0^{\mathcal{SH}^{fin}}). \end{array}$$

The map $\operatorname{Spc}(R)$ takes a prime ideal \mathcal{C} of \mathcal{SH}^{fin} to its preimage under $R = R_{\mathbb{C}}$. The endomorphism ring $\pi_0^{\mathcal{SH}^{fin}}$ is isomorphic to \mathbb{Z} , generated by the identity $S^0 \to S^0$. By [Mor06, Corollary 4.11], the same holds for $\pi_0^{\mathcal{SH}(\mathbb{C})_f} \cong \mathbb{Z}$. Since $R(\operatorname{id} : S_{\mathbb{C}}^0 \to S_{\mathbb{C}}^0) =$ (id : $S^0 \to S^0$), it follows that the map $\operatorname{Spec}(\pi_0(R))$ is isomorphic to the identity map $\operatorname{Spec}(\mathbb{Z}) \to \operatorname{Spec}(\mathbb{Z})$. Thus:

Corollary 7.2.7. For any $0 < n \leq \infty$, the thick ideal $R^{-1}(\mathcal{C}_n) \subseteq (\mathcal{SH}(\mathbb{C})_f)_{(p)}$ is a prime ideal, and

$$\rho_{(\mathcal{SH}(\mathbb{C})_f)_{(p)}}(R^{-1}(\mathcal{C}_n)) = \begin{cases} 0 & \text{if } n = 1, \\ p\mathbb{Z}_{(p)} & \text{if } n > 1. \end{cases}$$

Remark 7.2.8. From Proposition 7.1.4(2) or [Bal10, Proposition 10.4], we know that $\operatorname{Spc}(R)$ is not surjective, since $C\eta$ lies in some prime ideal which is not of the form $R^{-1}(\mathcal{C}_n)$.

For $F = \mathbb{R}$, there is also a commutative diagram:

$$\operatorname{Spc}(\mathcal{SH}(\mathbb{R})_{f}) \xrightarrow{\rho} \operatorname{Spc}(\pi_{0}^{\mathcal{SH}(\mathbb{R})_{f}})$$

$$\operatorname{Spc}(R') \bigwedge^{f} \operatorname{Spc}(\pi_{0}(R'))$$

$$\operatorname{Spc}(\mathcal{SH}(\mathbb{Z}/2)_{f}) \xrightarrow{\rho} \operatorname{Spc}(\pi_{0}^{\mathcal{SH}(\mathbb{Z}/2)_{f}}).$$

Lemma 7.2.9. In the above diagram, the right map $\operatorname{Spec}(\pi_0(R'))$ is an isomorphism.

Proof. The realisation functor *R'* maps generators of $\pi_0^{S\mathcal{H}(\mathbb{R})} \cong GW(\mathbb{R}) \cong \mathbb{Z} \oplus \mathbb{Z}$ to generators of $\pi_0^{S\mathcal{H}(\mathbb{Z}/2)}$. This follows from [Mor06, Section 4] and is explained on [Dug12, Slides 15-16 and 22-24]: $\pi_0^{S\mathcal{H}(\mathbb{R})}$ is generated by 1 and $\epsilon = -1 - \rho_{-1}\eta$ (in *GW*(ℝ)), ϵ corresponds to the quadratic form $q(x) = -x^2$). The element $\epsilon \in \pi_0^{S\mathcal{H}(\mathbb{R})}$ is also represented by the twist map $S^{1,1} \land S^{1,1} \to S^{1,1} \land S^{1,1}$ [Dug12, Slide 22]. Now, $S^{1,1} = \mathbb{G}_m = \mathbb{A}^1 \setminus \{0\}$ is mapped by *R'* to the circle with $\mathbb{Z}/2$ acting by involution, which is also denoted by $S^{1,1}$. Hence, $R'(\epsilon) = \epsilon : S^{1,1} \land S^{1,1} \to S^{1,1} \land S^{1,1}$, the twist map in $S\mathcal{H}(\mathbb{Z}/2)$. By [Dug12, Slide 16], 1 and ϵ are generators for $\pi_0^{S\mathcal{H}(\mathbb{Z}/2)}$. Thus, $\pi_0(R') : \pi_0^{S\mathcal{H}(\mathbb{R})} \to \pi_0^{S\mathcal{H}(\mathbb{Z}/2)}$ is an isomorphism.

Corollary 7.2.10. For any prime ideal $C_{m,n} \subseteq (\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}, (R')^{-1}(C_{m,n})$ is a prime ideal of $(\mathcal{SH}(\mathbb{R})_f)_{(p)}$ and $\rho_{(\mathcal{SH}(\mathbb{R})_f)_{(p)}}((R')^{-1}(C_{m,n}))$ can be identified with the same prime ideal of $\mathbb{Z}_{(p)} \oplus \mathbb{Z}_{(p)}$ as $\rho_{(\mathcal{SH}(\mathbb{Z}/2)_f)_{(p)}}(C_{m,n})$.

Remark 7.2.11. Conjecture 7.1.7(4) would imply that in the above diagram, $(R')^{-1}$ is not surjective, as thickid $(C\eta_{(p)})$ would have to lie in some prime ideal which is not of the form $(R')^{-1}(\mathcal{C}_{m,n})$.

Remark 7.2.12. Since the Künneth formula might not hold for AK(n), we do not know whether $\mathcal{C}_{AK(n)}$ or $\mathcal{C}_{AK(m,n)}$ are prime ideals.

Remark 7.2.13. If $k \subset \mathbb{C}$ or $k \subset \mathbb{R}$ is a field such that $\operatorname{Spec}(GW(\mathbb{C})) \to \operatorname{Spec}(GW(k))$ is not surjective or $\operatorname{Spec}(GW(\mathbb{R})) \to \operatorname{Spec}(GW(k))$ is not surjective respectively then [Bal10, Corollary 10.1] already implies that there is an infinite family of thick ideals in $\mathcal{SH}(k)_f$ which are not of the form $R_k^{-1}(\mathcal{C})$ or $(R'_k)^{-1}(\mathcal{C})$ respectively.

Chapter 8

Motivic type-n spectra

Definition 8.0.1. Let AK(n) be the motivic Morava K-theory spectrum as defined in Definition 6.3.1. We say that $X \in (\mathcal{SH}(k)_f)_{(p)}$ has motivic type n if $AK(n-1)_{**}(X) = 0$ and $AK(n)_{**}(X) \neq 0$.

A priori, the motivic type of X might not be unique, as we do not know whether $\mathcal{C}_{AK(n-1)} \subseteq \mathcal{C}_{AK(s)}$ for all s < n. In Section 9.6, we will prove that any $X \in \mathcal{SH}(\mathbb{C})_{(p)}^{fin}$, p > 2, has a unique motivic type. For any prime p, the motivic type n spectra that we are going to consider in this chapter satisfy $AK(s)_{**}(X) = 0$ for all s < n.

Remark 8.0.2. In the topological category $\mathcal{SH}_{(p)}^{fin}$, the notion of types is equivalent to the notion of thick ideals by the thick subcatgory theorem [HS98, Theorem 7]. In Chapter 3, we have seen that, in equivariant homotopy theory, a more general notion of types is required. Also in the motivic world, not every thick ideal can be described in the language of types, as defined above. For example, as shown in Chapter 7, the motivic Morava K-theories AK(n) do not distinguish between the nonequal thick ideals thickid $(C\eta_{(2)})$ and thickid $(S_{(2)}^0)$, as both are generated by a spectrum of motivic type 0.

In this chapter, we will often assume $k = \mathbb{C}$, because we will need explicit knowledge of H^{**} and of $AK(n)_{**}$ for some of our arguments. The results might hold in greater generality, but this seems to require different methods of proof.

If $X_n \in \mathcal{SH}_{(p)}^{fin}$ has type *n* (see Definition 2.0.11), then $c(X_n) \in \mathcal{SH}(\mathbb{C})_{(p)}^{fin}$ has motivic type *n* (Section 8.7). That is, $c(X_n) \in \mathcal{C}_{AK(n-1)}$ and $c(X_n) \notin \mathcal{C}_{AK(n)}$. For any given *n*, we show how to construct a spectrum $\mathbb{X}_n \in (\mathcal{SH}(\mathbb{C})_f)_{(p)}$ with motivic type *n* that is not in the image of the functor *c*. The construction will be similar to topological constructions given by [Mit85, Section 4] and [Rav92, Appendix C]. We stick to the approach in [Rav92] but we believe that a motivic version of Mitchell's spectrum would give another spectrum of motivic type *n*.

This chapter is organised as follows:

We first discuss some foundations that are needed to apply the motivic Adams spectral sequence and obtain a vanishing result for motivic Morava K-theory (Theorem 8.4.1). Afterwards, we construct a spectrum satisfying the conditions of the theorem and we prove that it has indeed motivic type n (Section 8.5). Finally, we compare our findings to a constant type-*n* spectrum, $c(X_n)$ (Section 8.7), realising that motivic Morava K-theory does not distinguish the thick ideal generated by \mathbb{X}_n from the one generated by cX_n , meaning that both spectra have motivic type *n*. We partly calculate their types with respect to the cohomology theories c(K(s)), as well. However, we do not have the answer to the question whether thickid(\mathbb{X}_n) equals thickid($c(X_n)$).

8.1 Universal coefficient and Künneth theorems

This section states some general results, which hold in SH(k), k any field, and which will be used later on. In the whole chapter, we use the notation $H = H\mathbb{Z}/p$. Recall from Remark 6.3.3(1) that H is cellular.

Proposition 7.7 in [DI05] describes the following universal coefficient spectral sequences:

Proposition 8.1.1. Let $E \in S\mathcal{H}(k)$ be a motivic ring spectrum, M a right E-module and N a left E-module. Furthermore, assume that $E, M \in S\mathcal{H}(k)^{cell}$ (see Definition 5.2.1).

(1) There is a strongly convergent spectral sequence

$$E^2 = \operatorname{Tor}_{a,b,c}^{E_{**}}(M_{**}, N_{**}) \Rightarrow \pi_{a+b,c}(M \wedge_E N).$$

(2) There is a conditionally convergent spectral sequence

$$E_2 = \text{Ext}_{E_{**}}^{a,b,c}(M_{**}, N_{**}) \Rightarrow \pi_{-a-b,-c}F_E(M, N),$$

where $F_E(-,-)$ denotes the E-function spectrum.

We apply (2) to the case $E = H = H\mathbb{Z}/p$, $M = A \wedge H$ and N = H for A a cell spectrum and get:

$$\operatorname{Ext}_{\operatorname{H}_{**}}(\operatorname{H}_{**}A, \operatorname{H}_{**}) \Rightarrow \pi_{**}F_H(A \wedge H, H) = \operatorname{H}^{**}A.$$

If A is a finite cell spectrum, we can also apply the spectral sequence to the case E = H, M = F(A, H), N = H:

$$\operatorname{Ext}_{\operatorname{H}^{**}}(\operatorname{H}^{**}A, \operatorname{H}^{**}) \Rightarrow \pi_{**}F_H(F(A, H), H) = \pi_{**}F(F(A, S^0), H) = \operatorname{H}_{**}A.$$

The first equality holds because $F(A, H) = F(A, S^0) \wedge H$ for finite A and the second holds because taking the dual of A twice gives A again (see [LMS86, Proposition III.1.3]).

In the case of vanishing higher Ext-groups, the spectral sequence collapses and we get the following result:

Corollary 8.1.2. If $A \in S\mathcal{H}(k)^{cell}$ is any cell spectrum such that $H_{**}A$ is free over H_{**} , then $H^{**}A \cong \operatorname{Hom}_{H_{**}}(H_{**}A, H_{**})$. If A is a finite cell spectrum with $H^{**}A$ free over H^{**} , then $H_{**}A \cong \operatorname{Hom}_{H^{**}}(H^{**}A, H^{**})$.

The universal coefficient spectral sequence also implies the following Künneth theorem [DI05, Remark 8.7]: **Proposition 8.1.3.** Let A and B be motivic spectra such that A is a finite cell spectrum. If $H^{**}A$ is free over H^{**} , then

$$\mathrm{H}^{**}(A) \otimes_{\mathrm{H}^{**}} \mathrm{H}^{**}(B) \cong \mathrm{H}^{**}(A \wedge B).$$

8.2 The motivic Steenrod algebra

In this section, too, we let $H = H\mathbb{Z}/p$. Let $k \subseteq \mathbb{C}$.

The motivic mod-*p* Steenrod algebra, $\mathcal{A} = \mathcal{A}^{**}$ has first been defined in [Voe03b, Section 11] as the algebra of certain bistable natural transformations $H^{**}(-) \rightarrow H^{**}(-)$. By [Voe10, Theorem 3.49] and [Hoy13, Lemma 5.7], \mathcal{A} is the algebra of all such operations and, as an H^{**} -module,

$$\mathcal{A} \cong \mathrm{H}^{**} \otimes_{\mathbb{F}_p} \mathcal{A}_{\mathrm{top}},$$

where \mathcal{A}_{top} is the topological mod-*p* Steenrod algebra. Thus, as H^{**}-modules,

$$\mathcal{A} \cong \mathrm{H}^{**} \otimes_{\mathbb{F}_p} RP \otimes_{\mathbb{F}_p} \Lambda_{\mathbb{F}_p}(Q_0, Q_1, \cdots),$$

where RP is the \mathbb{F}_p -module generated by certain products of reduced powers P^i : $\mathrm{H}^{**}(-) \to \mathrm{H}^{*+2i(p-1),*+i(p-1)}(-), i \geq 0$, and $\Lambda_{\mathbb{F}_p}(Q_0, Q_1, \cdots)$ denotes the exterior algebra over \mathbb{F}_p generated by $Q_i : \mathrm{H}^{**}(-) \to \mathrm{H}^{*+2p^i-1,*+p^i-1}, i \geq 0$, as defined in [Voe03b, Section 13]. See also [Bor03, Corollaries 3 and 4] or [Yag05, Equation (2.18)].

Borghesi [Bor03, Theorem 12] computes the cohomology of the motivic connective Morava K-theory spectrum, which can be expressed by the same formula as in topology.

Proposition 8.2.1.

$$\mathbf{H}^{**}(Ak(s)) = \mathcal{A}/\mathcal{A}Q_s = \mathbf{H}^{**} \otimes RP \otimes \Lambda_{\mathbb{F}_n}(Q_0, \cdots, Q_{s-1}, Q_{s+1}, \cdots).$$

If X is a finite cell spectrum such that $H^{**}(X)$ is free over H^{**} , we can apply the Künneth theorem to $Ak(s) \wedge X$.

Corollary 8.2.2. Let $X \in S\mathcal{H}(k)^{fin}$ with $H^{**}(X)$ free over H^{**} . Then

$$\mathrm{H}^{**}(Ak(s) \wedge X) = \mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} \mathrm{H}^{**}(X).$$

Writing $\Lambda(Q_s)$ for the exterior algebra over H^{**} generated by Q_s , we have $\mathcal{A} = \mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} \Lambda(Q_s)$. Any resolution of $\mathrm{H}^{**} X$ by projective $\Lambda(Q_s)$ -modules P_i yields a resolution $\mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} P_i$ of $\mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} \mathrm{H}^{**} X$ by projective \mathcal{A} -modules. This implies the following change of rings isomorphism.

Corollary 8.2.3. For any finite cell spectrum X with $H^{**}(X)$ free over H^{**} , we have

$$\operatorname{Ext}_{\mathcal{A}}(\operatorname{H}^{**}(Ak(s) \wedge X), \operatorname{H}^{**}) \cong \operatorname{Ext}_{\Lambda(Q_s)}(\operatorname{H}^{**}(X), \operatorname{H}^{**})$$

This isomorphism will later be applied to the motivic Adams spectral sequence for $Ak(s)_{**}(X)$.

8.3 The motivic Adams spectral sequence

Recall the notation $H = H\mathbb{Z}/p$. Let $k \subseteq \mathbb{C}$.

Our aim in this section is to show the existence and convergence of the following Adams spectral sequence:

$$E_2^{s,t,u} = \operatorname{Ext}_{\mathcal{A}}^{s,t,u}(\operatorname{H}^{**}(Ak(s) \wedge X), \operatorname{H}^{**}) \Rightarrow Ak(s)_{**}(X)$$

for finite cell spectra $X \in \mathcal{SH}(k)^{fin}$ (see Definition 5.2.1), with $H^{**}X$ free over H^{**} .

Motivic Adams spectral sequences were first described in [Mor99]. [HKO11, Corollary 3] shows that over fields of characteristic 0, the Adams spectral sequence for a motivic cell spectrum X of finite type converges to the homotopy groups of the completion $X_{p,\eta}^{\wedge}$. In [DI10], calculations are made for the case p = 2, X = S. More details and explanations can be found in [Sta12]. The convergence of the Adams spectral sequence for $Ak(s) \wedge X$ will follow from [HKO11]. However, we have to start again from the Adams resolution, to get the limit term $Ak(s)_{**}(X)$ using arguments from [Rav86, Section 2.1], and to get a module structure on the sequence.

Definition 8.3.1. An Adams resolution $(Y_s, g_s, K_s, f_s)_{s \ge 0}$ for a motivic cell spectrum $Y \in \mathcal{SH}(k)^{cell}$ is a diagram



where each K_s is a wedge of suspensions of H, f_s is surjective on motivic mod-p cohomology and Y_{s+1} is the fiber of f_s .

Such an Adams resolution exists whenever $H^{**}Y$ is a free module over H^{**} of motivically finite type [HKO11], [Sta12, Section 2.5.4]. The finite type condition is defined in [D110, Definition 2.12]. For our purposes, it suffices to know that, if the generators of $H^{**}Y$ are located in degrees $(i_{\alpha}, j_{\alpha})_{\alpha}$ with i_{α} bounded below and, for each α , there are only finitely many β 's with $i_{\alpha} = i_{\beta}$ and $j_{\alpha} \geq j_{\beta}$, then $H^{**}Y$ is of motivically finite type. This holds, for example, if $Y \in S\mathcal{H}(k)^{fin}$.

Corollary 8.3.2. Let X be a finite cell spectrum such that $H^{**}X$ is free over H^{**} and let Ak(s) be the motivic connective Morava K-theory spectrum (see Definition 6.3.1). Then $Y = Ak(s) \wedge X$ satisfies the finite type condition as described above.

Proof. By Corollary 8.2.2, $\mathrm{H}^{**}(Y) \cong \mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} \mathrm{H}^{**} X$. The motivic Steenrod algebra \mathcal{A} is of motivically finite type, since, for the bidegrees (i_{α}, j_{α}) of its H^{**} -generators, i_{α} and j_{α} are nonnegative and there are only finitely many generators of a fixed bidegree. The same holds for $\mathcal{A}/\mathcal{A}Q_s$. Furthermore, $\mathrm{H}^{**} X$ is of motivically finite type, since X is finite. It follows that the tensor product $\mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} \mathrm{H}^{**} X$ is also of motivically finite type.

Corollary 8.3.3. (Convergence of the ASS)

Let $k = \mathbb{C}$ (or any other field satisfying the assumptions of [HKO11, Theorem 1]) and let X be a finite cell spectrum such that $H^{**}(X)$ is a free H^{**} -module. Then the Adams spectral sequence for $Y = Ak(s) \wedge X$, s > 0, strongly converges to $Ak(s)_{**}(X)$.

Proof. Since Y is of motivically finite type by the previous corollary, strong convergence follows from [HKO11, Corollary 3]. By [HKO11, Theorem 1], the spectral sequence converges to the *p*-completion of $\pi_{**}Y$. Since Ak(s) is a quotient of ABP/(p) by definition, $Y_p^{\wedge} = Y$, whence the limit term is $Ak(s)_{**}(X)$.

Remark 8.3.4. In Corollary 8.3.3, X does not necessarily need to be finite. All we need to know is that $Ak(s) \wedge X$ is of motivically finite type.

In Theorem 8.4, it will be crucial that this particular spectral sequence is a spectral sequence of $(MGL_{(p)})_{**}$ -modules.

Since Ak(s) is an $MGL_{(p)}$ -module spectrum by Remark 6.3.3(3), $Ak(s)_{**}X$ and $H^{**}(Ak(s) \wedge X)$ are $(MGL_{(p)})_{**}$ -modules. Before we can prove compatibility of the Adams spectral sequence with the module structure, we need to determine $H_{**}Ak(s)$. Borghesi [Bor03, Theorem 5 and Remark 2.2] shows:

Lemma 8.3.5.

$$\mathrm{H}_{**}(MGL) \cong \mathrm{H}_{**}[m_1, m_2, \cdots] \cong \mathrm{H}_{**} \otimes_{\mathbb{F}_p} \mathrm{H}_{*}(MU, \mathbb{Z}/p).$$

Since $\pi_{**}(H \wedge MGL_{(p)}) = \pi_{**}(H_{(p)} \wedge MGL)$ and $H_{(p)} = H$, it follows that also $H_{**}(MGL_{(p)})$ is a free H_{**} -module with basis elements $m_I = m_{i_1}^{k_1} m_{i_2}^{k_2} \cdots m_{i_t}^{k_t}$. Here, m_i is defined as the Hurewicz image of $a_i \in MGL_{2i,i}$ and a_i is the image of a polynomial generator of MU_* . The motivic connective Morava K-theory spectra are defined as homotopy colimits of spectra E_i , which are defined by successively taking cofibers of maps $\Sigma^{2(i-1),i-1}E_{i-1} \xrightarrow{a_{i-1}} E_{i-1}$, starting from $MGL_{(p)}$ (see Definition 6.3.1). Passing from $H_{**}E_{i-1}$ to $H_{**}E_i$, the H_{**} -basis changes but the fact that the homology is a free H_{**} -module remains true for each i.

Corollary 8.3.6. $H_{**}(Ak(s))$ is a free H_{**} -module. Hence,

$$H_{**}(Ak(s)) \cong Hom_{H^{**}}(H^{**}(Ak(s)), H^{**})$$

Proof. The second statement follows from Corollary 8.1.2. For the cellularity of Ak(s), see Remark 6.3.3(1).

Proposition 8.3.7. Let k and X be as in Corollary 8.3.3. The Adams spectral sequence for $Ak(s) \wedge X$ is a spectral sequence of $(MGL_{(p)})_{**}$ -modules.

Proof. We show that there is an Adams resolution by $MGL_{(p)}$ -modules. The construction of the spectral sequence from the Adams resolution preserves such a module structure in every step and the claim will follow.

We use similar arguments as in Corollary 8.3.3, but starting with an Adams resolution $\{X_t, g_t, K_t, f_t\}$ for X. Then we take the smash product of such a resolution with Ak(s) and show that this yields an Adams resolution for $Y = Ak(s) \wedge X$. Since Ak(s)is an $MGL_{(p)}$ -module, this will be a resolution by $MGL_{(p)}$ -modules. By definition, f_t is surjective on cohomology, which is equivalent to g_t inducing zero in cohomology. It follows that $Ak(s) \wedge g_t$ induces zero in cohomology, hence, $Ak(s) \wedge f_t$ is surjective on cohomology by the long exact fiber sequence.

It remains to show that $Ak(s) \wedge K_t$ is a wedge of suspensions of H. Consider one wedge summand H of K_t . By Corollary 8.3.6, $\pi_{**}(Ak(s) \wedge H)$ is free over $\pi_{**}(H)$. Furthermore, $Ak(s) \wedge H$ is a cellular H-module. Therefore, [Hoy13, Lemma 5.3] implies that $Ak(s) \wedge H$ is split, that is, $Ak(s) \wedge H \cong \bigvee \Sigma^{**}H$, as we wanted to show. \Box

8.4 Vanishing criterion for motivic Morava K-theory

Now we can prove the following motivic version of the vanishing result [Mit85, Theorem 4.8].

Theorem 8.4.1. (Vanishing criterion)

Let p be any prime, s > 0 and k be as in Corollary 8.3.3. Let $X \in S\mathcal{H}(k)^{fin}$ be a finite motivic cell spectrum such that $H^{**}X$ is free over $\Lambda(Q_s)$ (the exterior algebra over H^{**}) as a module over the Steenrod algebra. Then

$$AK(s)_{**}X = 0.$$

Proof. With the preparations made so far, the rest of the proof is exactly as in [Mit85, Theorem 4.8]. Since $AK(s) = v_s^{-1}Ak(s)$, it suffices to show that $Ak(s)_{**}X$ is v_s -torsion. We apply the change of rings isomorphism, Corollary 8.2.3, to the Adams spectral sequence for $Ak(s) \wedge X$ and get:

$$E_2 \cong \operatorname{Ext}_{\Lambda(Q_s)}(\operatorname{H}^{**} X, \operatorname{H}^{**}) \Rightarrow Ak(s)_{**}X.$$

By the assumption on X, this collapses to $\operatorname{Hom}_{\Lambda(Q_s)}(\operatorname{H}^{**}X, \operatorname{H}^{**}) \cong Ak(s)_{**}X$, and, by the previous proposition, this is an isomorphism of $(MGL_{(p)})_{**}$ -modules. Since $(MGL_{(p)})_{**}$ acts trivially on the left hand side of this isomorphism, v_s acts trivially on $Ak(s)_{**}X$, too. Hence, $AK(s)_{**}X = 0$.

8.5 Construction of an X such that $H^{**}X$ is free over $\Lambda(Q_s)$

In this section, we assume $k = \mathbb{C}$ because we will work explicitly with $\mathrm{H}^{**} \cong \mathbb{F}_p[\tau]$, deg $(\tau) = (0, 1)$ (see Lemma 6.3.6). The construction of a spectrum X with the properties required in the previous theorem can be done similarly as in [Rav92, Appendix C]. That is, one starts with a so-called weakly type n spectrum (n > s) and then uses a particular idempotent to split off a (strongly) type n spectrum of a certain smash power of it.

8.5.1 Idempotents for free H^{**}-modules

For V^{**} an Adams graded graded abelian group (i.e., V^{**} has a sign rule in the first grading but not in the second one, see e.g. [NSØ09, Section 3]) which is a free H^{**}module, let $V^+ = \bigoplus_{p \text{ even}} V^{p,q}$ and $V^- = \bigoplus_{p \text{ odd}} V^{p,q}$ be the even and odd dimensional parts of V. That is, commuting with an element of V^+ does not change the sign but commuting two elements of V^- does. For a vector space V^* over \mathbb{F}_p , Ravenel defines a number k_V and an idempotent $e_V \in \mathbb{Z}_{(p)}[\Sigma_{k_V}]$, which only depend on $\dim_{\mathbb{F}_p} V^+$ and $\dim_{\mathbb{F}_p} V^-$ [Rav92, Appendix C.2]. The analogous definition can be formulated using $\dim_{\mathbb{H}^{**}} V^+$ and $\dim_{\mathbb{H}^{**}} V^-$ for our bigraded \mathbb{H}^{**} -modules. The symmetric group Σ_{k_V} acts on $V^{\otimes k_V}$ by permuting the factors. As V is an \mathbb{F}_p -module, this induces an action of $\mathbb{Z}_{(p)}[\Sigma_{k_V}]$ on $V^{\otimes k_V}$. For our purposes, it will not be important to know the precise definitions of k_V and e_V . We just need to know that they are defined in such a way that the following analogue of [Rav92, Theorem C.2.1] holds.

Proposition 8.5.1. Let k_V and $e_V \in \mathbb{Z}_{(p)}[\Sigma_{k_V}]$ be the number and idempotent defined in [Rav92, Appendix C.2] and let $W = V^{\otimes k_V}$. Then $e_V W \neq 0$. If $U \subset V$ has dim $U^+ \leq$ dim $V^+ - 1$ or dim $U^- \leq \dim V^- - (p-1)$, then $e_V U^{\otimes k_V} = 0$. Here, dim denotes the H^{**}-dimension and \otimes denotes the tensor product over H^{**}.

Proof. The proof of [Rav92, Theorem C.2.1] applies to our setting without changes. \Box

We will need the following Lemma.

Lemma 8.5.2. Let M be a module over $\mathbb{F}_p[\tau, Q]/Q^2$ which is free as a module over $\mathbb{F}_p[\tau]$ and free as a module over $\mathbb{F}_p[Q]/Q^2$. Then M is a free $\mathbb{F}_p[\tau, Q]/Q^2$ -module.

Proof. Let $\{m_i\}_{i\in I}$ be a basis of M over $\mathbb{F}_p[\tau]$ and $\{n_j\}_{j\in J}$ a basis over $\mathbb{F}_p[Q]/Q^2$. Then M is a free \mathbb{F}_p -module with bases $\{\tau^k m_i\}_{i\in I, k\in \mathbb{N}}$ and $\{n_j, Qn_j\}_{j\in J}$. As an $\mathbb{F}_p[Q]/Q^2$ -module, M decomposes as $M \cong M' \oplus QM'$ with $M' \cong QM' \cong QM$ as \mathbb{F}_p -modules. Hence, the elements m_i can be written as $m_i = a_i + Qb_i$ with $a_i, b_i \in M'$. For any $i \in I$ such that both a_i and b_i are nonzero, we replace the basis element m_i by the two elements a_i and Qb_i . Then we still have a set of generators for M over $\mathbb{F}_p[\tau]$, which can be turned into a basis by removing elements. Hence, we can assume that all m_i are of the form $m_i = a_i$ or $m_i = Qb_i$. Let $I' = \{i \in I \mid m_i = a_i \in M'\}$. Then $M' \cong \mathbb{F}_p\{\tau^k a_i\}_{i\in I', \tau\in \mathbb{N}}$ as an \mathbb{F}_p -module. Hence, $QM' \cong \mathbb{F}_p\{Q\tau^k a_i\}_{i\in I', \tau\in \mathbb{N}}$ and $M \cong \mathbb{F}_p\{\tau^k a_i, Q\tau^k a_i\}$. It follows that $\{a_i\}_{i\in I'}$ is a basis of M as a free $\mathbb{F}_p[\tau, Q]/Q^2$ -module. \square

Theorem C.2.2 of [Rav92] explains how to split off a free module over the exterior \mathbb{F}_p -algebra generated by Q_s from a module with nontrivial Q_s -action. In our setting, this also works for the exterior H^{**}-algebra $\Lambda(Q_s)$.

Proposition 8.5.3. Let $V = U \oplus F$ be a splitting of $\Lambda(Q_s)$ -modules which are free over H^{**} , and $F \neq 0$ be free over $\Lambda(Q_s)$. Then $e_V V^{\otimes k_V}$ is a free $\Lambda(Q_s)$ -module.

Proof. We write the tensor product as $V^{\otimes k_V} = U^{\otimes k_V} \oplus F'$, where $F' = \bigoplus_{\substack{a+b=k_V\\b\geq 1}} U^{\otimes a} \otimes F^{\otimes b}$.

By the proof of [Rav92, Theorem C.2.2], F' is free over the Hopf algebra $\mathbb{F}_p[Q_s]/Q_s^2$, which we abbreviate by E. Let us give the reason for this statement. We show that if the E-module U has basis $\{u_i\}_I$ over \mathbb{F}_p and F has basis $\{f_j\}_J$ over E, then $U \otimes_{\mathbb{F}_p} F$ has \mathbb{F}_p -basis $\{u_i \otimes f_j, Q_s(u_i \otimes f_j)\}_{I,J}$ and hence is a free module over E. The module structure of $U \otimes F$ is defined by $Q_s(u \otimes f) = Q_s u \otimes f + u \otimes Q_s f$. Since $\{f_j, Q_s f_j\}_J$ defines an \mathbb{F}_p -basis of F, an \mathbb{F}_p -basis of $U \otimes F$ can be given by $\{u_i \otimes f_j, u_i \otimes Q_s f_j\}$. In the formula $Q_s(u_i \otimes f_j) = Q_s u_i \otimes f_j + u_i \otimes Q_s f_j, Q_s u_i = \Sigma r_k u_k$ can be expressed by the basis elements of U, hence $Q_s u_i \otimes f_j \in \mathbb{F}_p\{u_k \otimes f_j\}$ and the basis elements $u_i \otimes Q_s f_j$ of $U \otimes F$ can be replaced by $Q_s(u_i \otimes f_j)$. We obtain $U \otimes F = \mathbb{F}_p\{u_i \otimes f_j, Q_s(u_i \otimes f_j)\}$. Inductively, it follows that all mixed summands $U^{\otimes a} \otimes F^{\otimes b}$ in $V^{\otimes k_V}$ are free and hence F' is free over E.

The analogue holds if we consider U and F as free modules over $\mathbb{F}_p[\tau]$ instead of \mathbb{F}_p and use $\otimes_{\mathbb{F}_p[\tau]}$. It follows that F' is free over $\Lambda(Q_s)$. The direct summands are invariant under the Σ_{k_V} -action. Hence, we have a short exact sequence

$$0 \to e_V U^{\otimes k_V} \to e_V V^{\otimes k_V} \to e_V F' \to 0.$$

Since $\deg(Q_s) = (2p^i - 1, p^i - 1)$, multiplication by Q_s sends V^+ to V^- and vice versa. It follows that $\dim F^+ > 0$ (and $\dim F^- > 0$) and, hence, $\dim U^+ < \dim V^+$. By the previous proposition, this implies $e_V U^{\otimes k_V} = 0$. It follows that $e_V V^{\otimes k_V} = e_V F'$. We have to show that $e_V F'$ is a free $\Lambda(Q_s)$ -module. As a module over the exterior \mathbb{F}_{p^-} algebra over Q_s , this is a direct summand of a free module over a local ring. Hence, $e_V F'$ is free over $\mathbb{F}_p[Q_s]/(Q_s^2)$. Since $e_V F'$ is also a free H^{**}-module, it is free over $\Lambda(Q_s) = \mathbb{F}_p[\tau, Q_s]/Q_s^2$ by Lemma 8.5.2.

We can apply this to motivic cohomology in the following way:

Theorem 8.5.4. (Splitting off free $\Lambda(Q_s)$ -modules)

Let $X \in S\mathcal{H}(\mathbb{C})_{(p)}^{fin}$ be a p-local finite cell spectrum such that Q_s acts nontrivially on $\mathrm{H}^{**}(X)$ as an element of the Steenrod algebra. Assume that $V = \mathrm{H}^{**}(X)$ is a free H^{**} -module and let $Y = e_V(X^{\wedge k_V})$. Then $\mathrm{H}^{**}(Y)$ is free over $\Lambda(Q_s)$.

Proof. This is analogous to a statement in [Rav92, Theorem C.3.2]. Since Q_s acts nontrivially, $H^{**}(X)$ contains a nontrivial summand which is free over $\Lambda(Q_s)$. The previous proposition yields the claim. Note that $H^{**}(e_V X^{\wedge k_V}) = e_V H^{**}(X)^{\otimes k_V}$ by the Künneth theorem (Proposition 8.1.3) and by the way $\mathbb{Z}_{(p)}[\Sigma_{k_V}]$ acts. The Künneth theorem also holds for *p*-local finite spectra because *p*-localisation commutes with $H^{**}(-)$.

Y is a retract of the p-local finite cell spectrum $X^{\wedge k_V}$, but maybe it is not finite itself. Therefore, we need an additional argument which shows that Theorem 8.4.1 holds for Y.

Corollary 8.5.5. For s > 0 and Y as in Theorem 8.5.4, $AK(s)_{**}(Y) = 0$.

Proof. We have to show $\mathrm{H}^{**}(Ak(s) \wedge Y) \cong \mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} \mathrm{H}^{**}(Y)$. Since $\mathrm{H}^{**}Y \cong e_V \mathrm{H}^{**}(X)^{\otimes k_V}$, Y is of motivically finite type. Remark 8.3.4 applies and the claim follows as in the proof of Theorem 8.4.1. The left hand side of the claimed isomorphism can be rewritten as

$$\mathrm{H}^{**}(Ak(s) \wedge e_V X^{\wedge k_V}) \cong \mathrm{H}^{**}((1 \wedge e_V)(Ak(s) \wedge X^{\wedge k_V})) \cong (1 \otimes e_V) \mathrm{H}^{**}(Ak(s) \wedge X^{\wedge k_V}).$$

Now we can apply the Künneth isomorphism and get $(1 \otimes e_V)(\mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} \mathrm{H}^{**}(X)^{\wedge k_V})$. This is isomorphic to $\mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} e_V \mathrm{H}^{**}(X)^{\wedge k_V} \cong \mathcal{A}/\mathcal{A}Q_s \otimes_{\mathrm{H}^{**}} \mathrm{H}^{**}(Y)$, which is the right hand side.

This result tells us that, given a nontrivial Q_s -action on $\mathrm{H}^{**}(X), X \in \mathcal{SH}(\mathbb{C})^{fin}_{(p)}$, we can construct a spectrum Y for which $AK(s)_{**}(Y) = 0$. So, let's construct such an X.

8.5.2 A finite cell spectrum with nontrivial Q_s -action and trivial Q_n action

Let $k = \mathbb{C}$. We combine ideas of Ravenel [Rav92] with computations by Voevodsky [Voe03b]. In [Rav92, Lemma 6.2.6], the given example of a spectrum with nontrivial Q_s^{top} -action, s < n, and trivial Q_n^{top} -action on $H^*(X)$ is $X = (B\mathbb{Z}/p)_2^{2p^n}$, that is, the suspension spectrum of the $2p^n$ -skeleton of the classifying space $B\mathbb{Z}/p$ modulo its 1skeleton. Cutting off higher dimensional cells leads to a trivial Q_n^{top} -action, which is needed for nontrivial *n*-th Morava K-theory. In [Voe03b, Section 6], the algebraic analogue to $B\mathbb{Z}/p$ is defined as $B\mu_p = \operatorname{colim}_n \tilde{V}_n/\mu_p$, where $\tilde{V}_n = \mathbb{A}^n \setminus \{0\}$ (see the proof of [Voe03b, Lemma 6.3]) and μ_p acts by multiplication with a *p*-th root of unity in each of the *n* coordinates. Under $R = R_{\mathbb{C}}$, this action realises to the \mathbb{Z}/p -action on $S^{2n-1} \subset \mathbb{C}^n$ rotating each \mathbb{C} factor by a *p*-th root of unity.

Lemma 8.5.6.

$$R(\tilde{V}_n/\mu_p) \cong S^{2n-1}/(\mathbb{Z}/p)$$

is the (2n-1)-skeleton of $B\mathbb{Z}/p$ in the CW-structure having one cell in each dimension.

Proof. \mathbb{BZ}/p is the infinite dimensional lens space, as studied for example in [Hat01, Example 2.43]. There, it is explained that the (2n-1)-skeleton is precisely the (2n-1)-dimensional lens space, which is defined as the orbit space $S^{2n-1}/(\mathbb{Z}/p)$.

By [Hat01, Example 2.43, page 146], the attaching map of the 2k-cell of $B\mathbb{Z}/p$ is the quotient map $S^{2k-1} \to S^{2k-1}/(\mathbb{Z}/p)$. We define $V_n \in S\mathcal{H}(\mathbb{C})$ to be the cofiber of the quotient map of suspension spectra $\tilde{V}_{p^n} \to \tilde{V}_{p^n}/\mu_p$, so that the following lemma holds.

Lemma 8.5.7.

$$R(V_n) = (B\mathbb{Z}/p)^{2p^n}.$$

Let \mathbb{B} be the cofiber of the composite map $\tilde{V}_1/\mu_p \to \tilde{V}_{p^n}/\mu_p \to V_n$. Then \mathbb{B} is a finite cell spectrum and satisfies the following corollary.
Corollary 8.5.8.

$$R(\mathbb{B}) = (B\mathbb{Z}/p)_2^{2p^n}.$$

The following is a special case of [Voe03b, Proposition 6.10] (as explained on [Voe03b, page 20]).

Proposition 8.5.9. $\mathrm{H}^{**}(B\mu_p)$ is a free H^{**} -module with basis $\{v^i, uv^i \mid i \geq 0\}$, where $v \in \mathrm{H}^{2,1}(B\mu_p)$ and $u \in \mathrm{H}^{1,1}(B\mu_p)$.

From this, it follows for dimensional reasons:

Proposition 8.5.10. The cohomology $H^{**} \mathbb{B}$ is the free H^{**} -module with basis $\{v^i \mid 1 \leq i \leq p^n\} \cup \{uv^i \mid 1 \leq i \leq p^n - 1\}.$

Furthermore, Voevodsky shows [Voe03b, Lemmas 11.2 and 11.3]:

Lemma 8.5.11. On

$$\mathrm{H}^{**}(B\mu_p)/\mathrm{H}^{*,>0}\,\mathrm{H}^{**}(B\mu_p)\cong \mathbb{F}_p[u,v]/(u^2=0),$$

 $\mathcal{A}/(\mathbf{H}^{*,>0}\,\mathcal{A}) \text{ acts by } \beta(u) = v, P^{i}(u) = 0 \text{ for } i > 0, \ \beta(v^{k}) = 0 \text{ and } P^{i}(v^{k}) = {k \choose i} v^{k+i(p-1)}.$

Over \mathbb{C} , the action of Q_i can be defined by $Q_0 = \beta$ and $Q_{i+1} = P^{p^i}Q_i - Q_iP^{p^i}$ [Voe96, Proposition 3.1]. From this, we can inductively compute the action of Q_s on $uv^k \in \mathrm{H}^{**}(\mathbb{B})/\mathrm{H}^{*,>0}\mathrm{H}^{**}(\mathbb{B})$. We get $Q_s(uv^k) = cv^{k+p^s}$ with $c \equiv 1 \mod p$, which is nontrivial for s < n and $k < p^n - p^s$. It follows that Q_s acts nontrivially on $\mathrm{H}^{**}(\mathbb{B})$ for s < n.

Now we have all ingredients for the motivic type n spectrum in $\mathcal{SH}(\mathbb{C})$.

Theorem 8.5.12. (A spectrum of motivic type n)

For a fixed n > 0, let $V = \mathrm{H}^{**}(\mathbb{B}_{(p)})$ and $X = e_V(\mathbb{B}_{(p)})^{\wedge k_V}$, then $AK(s)_{**}(X) = 0$ for all s < n and $AK(n)_{**}(X) \neq 0$.

Proof. For s > 0, $AK(s)_{**}(X) = 0$ follows from Corollary 8.5.5, whose assumptions are satisfied by the above considerations.

For s = 0, note that $AK(0) = p^{-1}MGL_{(p)}/(a_1, a_2, \cdots)$ by Definition 6.3.1. The main result of [Hoy13] implies $AK(0) \cong p^{-1}(H\mathbb{Z})_{(p)}$. It follows that

$$AK(0)_{**}X \cong \pi_{**}(p^{-1}(H\mathbb{Z} \wedge X)_{(p)})$$

But p acts trivially on \tilde{V}_m/μ_p , which implies that X is p-torsion. Therefore, $p^{-1}X \cong 0$ and $AK(0)_{**}X = 0$. It remains to show that $AK(n)_{**}(X) \neq 0$. This can either be done analogously to [Mit85, Theorem 4.8], using the motivic Atiyah Hirzebruch spectral sequence from Proposition 6.3.4, or by considering the topological realisation $R_{\mathbb{C}}(X)$. In Section 8.6, we show that R(X) is of type n. It follows that $X \in R^{-1}(\mathcal{C}_n \setminus \mathcal{C}_{n+1})$. In particular, $X \notin R^{-1}(\mathcal{C}_{n+1})$. By Proposition 6.3.8, $\mathcal{C}_{AK(n)} \subseteq R^{-1}(\mathcal{C}_{n+1})$. This proves $X \notin \mathcal{C}_{AK(n)}$. **Remark 8.5.13.** The spectrum X is a retract of the *p*-local finite cell spectrum $\mathbb{B}_{(p)}^{\wedge k_V}$ and it follows by Remark 5.2.4(1) that X is compact, i.e., $X \in (\mathcal{SH}(\mathbb{C})_f)_{(p)}$.

Remark 8.5.14. $X = e_V(\mathbb{B}_{(p)})^{\wedge k_V}$ is an example of a motivic spectrum with vanishing Margolis homology groups $MH_s^{p,q}(X)$ for all $s < n, p, q \in \mathbb{Z}$, as defined in [Voe03a, Section 3]: we have shown that $\mathrm{H}^{**}(X)$ is free over $\Lambda(Q_s)$, which implies that ker $Q_s = \mathrm{im} Q_s$ for $Q_s : \mathrm{H}^{**}(X) \to \mathrm{H}^{**}(X)$.

8.6 The type of the realisation R(X)

Let $k = \mathbb{C}$ and $B = (B\mathbb{Z}/p)_2^{2p^n}$. We already know that for $R : \mathcal{SH}(\mathbb{C}) \to \mathcal{SH}, R(\mathbb{B}) = B$. Since R preserves \wedge -products, it follows

$$R(X) = R(e_V(\mathbb{B}_{(p)})^{\wedge k_V}) = e_V R((\mathbb{B}_{(p)}))^{\wedge k_V} = e_V(B_{(p)})^{\wedge k_V}$$

The \mathbb{F}_p -vector space $\mathrm{H}^*(B_{(p)})$ is generated by similar elements as the $\mathbb{F}_p[\tau]$ -vector space $V = \mathrm{H}^{**}(\mathbb{B}_{(p)})$ (compare Proposition 8.5.10 with [Rav92, Lemma 6.2.6]). In particular, $\dim_{\mathbb{F}_p}(\mathrm{H}^*(B_{(p)}))^+ = \dim_{\mathrm{H}^{**}} V^+$ and $\dim_{\mathbb{F}_p}(\mathrm{H}^*(B_{(p)}))^- = \dim_{\mathrm{H}^{**}} V^-$. It follows that R(X) is the type-*n* spectrum defined in [Rav92, Theorem C.3.2].

8.7 The constant type-n spectrum

In this section, let $k = \mathbb{C}$ and let $X_n = e_V B^{\wedge k_V}$ be the type-n spectrum defined by Ravenel. We want to determine the motivic type of cX_n , where $c : S\mathcal{H} \to S\mathcal{H}(\mathbb{C})$ as in Chapter 4. X_n is constructed via an idempotent from the finite cell spectrum $B = B\mathbb{Z}/p_2^{2p^n}$ [Rav92, Lemma 6.2.6]. First, we calculate $\mathrm{H}^{**}(cB)$.

Proposition 8.7.1. Let $X = \Sigma^{\infty} Y$ be the suspension spectrum of a finite CW complex. Then $H^{**}(cX) \cong H^*(X)[\tau]$ as \mathbb{F}_p -modules, where a generator in degree *i* from the right hand side maps to bidegree (i, 0) on the left hand side.

Proof. For any $F \in \mathcal{SH}(\mathbb{C})$, we have $\mathrm{H}^{**}(F) \cong \mathrm{H}^{*+n,*}(F \wedge S_s^n)$. For $F = S^0$, we get $\mathrm{H}^{**}(S_s^n) \cong \mathrm{H}^{*-n,*}(S^0) \cong \mathrm{H}^{*-n,*} \cong \mathrm{H}^*(S^n)[\tau]$. Now let $(Y^k)_k$ be a CW decomposition of Y, that is, $\Sigma^{\infty}Y^k$ is the cofiber of some $\Sigma^{\infty}\alpha_k : S^{n_k} \to \Sigma^{\infty}Y^{k-1}$. We write X^k for $\Sigma^{\infty}Y^k$. Since c preserves cofiber sequences, we get a cofiber sequence of suspension spectra $S_s^{n_k} \xrightarrow{c\alpha} cX^{k-1} \to cX^k$. It induces a long exact sequence

$$\cdots \to \mathrm{H}^{*-1,*}(cX^{k-1}) \stackrel{(c\alpha)^*}{\to} \mathrm{H}^{*-1,*}(S^{n_k}_s) \to \mathrm{H}^{**}(cX^k) \to \mathrm{H}^{**}(cX^{k-1}) \stackrel{(c\alpha)^*}{\to} \mathrm{H}^{**}(S^{n_k}_s) \to \cdots$$

We assume inductively that $\mathrm{H}^{i,j}(cX^{k-1}) \cong \mathrm{H}^i(X^{k-1})\{\tau^j\}$. let $x\tau^j \in \mathrm{H}^{i,j}(cX^{k-1})$. Since $R(\tau) = 1$ and $R(c\alpha) = \alpha$, we have

$$R((c\alpha)^*(x\tau^j)) = \alpha^*(R(x\tau^j)) = \alpha^*(x).$$

The only element in $\mathrm{H}^{i,j}(S_s^{n_k})$ which is mapped to $\alpha^*(x)$ by R is $\alpha^*(x)\tau^j$. This proves $(c\alpha)^* \cong \alpha^*[\tau]$. By the five lemma, it follows that the map $\mathrm{H}^{**}(cX^k) \to \mathrm{H}^*(X^k)[\tau]$, given by sending $x \in \mathrm{H}^{i,j}(cX^k)$ to $R(x)\tau^j$, is an isomorphism and, inductively, $\mathrm{H}^{**}(cX) \cong \mathrm{H}^*(X)[\tau]$.

Corollary 8.7.2. As \mathbb{F}_p -modules,

$$\mathbf{H}^{**}(cB) \cong \mathbf{H}^{*}(B)[\tau] \cong \mathbf{H}^{**}\{x^{k} \mid 1 \le k \le p^{n}\} \cup \{yx^{k} \mid 1 \le k \le p^{n} - 1\},\$$

with $\deg(x) = (2,0)$ and $\deg(y) = (1,0)$.

Proof. The cohomology of B is described in the proof of [Rav92, Lemma 6.2.6]. We only have to add the polynomial generator $\tau \in H^{**}$.

For X = cY, Y a finite CW spectrum, Proposition 6.3.4 implies the convergence of the motivic Atiyah Hirzebruch spectral sequence,

$$E_2^{p,q,t} = \mathbf{H}^{p+2t,q+t}(X) \otimes K(n)_t \Rightarrow AK(n)^{p,q}(X)$$

By Remark 6.3.5(1), the realisation functor R maps this spectral sequence to the topological Atiyah Hirzebruch spectral sequence:



where $(E_r^{p,q,t}, d_r)$ is the spectral sequence mentioned above and $(F_r^{p,t}, d_r')$ is the Atiyah Hirzebruch spectral sequence (see, e.g., [Rav92, Theorem A.3.7])

$$F_2^{p,t} = \mathrm{H}^{p+2t}(R(X)) \otimes K(n)_t \Rightarrow K(n)^p(R(X)).$$

Note that there is no differential in between d'_r and d'_{r+1} , since $K(n)_*$ is concentrated in even degrees.

Now let r be small enough, so that E_r and F_r are still equal to the 2-pages. Using Proposition 8.7.1, the above square is

$$\begin{array}{ccc} \mathrm{H}^{p+2t}(R(X)) \cdot \tau^{q+t} \otimes K(n)_t & \longrightarrow \mathrm{H}^{p+2t+2r+1}(R(X)) \cdot \tau^{q+t+r} \otimes K(n)_{t+2r} \\ & & & & \downarrow R \\ & & & \downarrow R \\ \mathrm{H}^{p+2t}(R(X)) \otimes K(n)_t & \longrightarrow \mathrm{H}^{p+2t+2r+1}(R(X)) \otimes K(n)_{t+2r}. \end{array}$$

Since $R(\tau) = 1$, the vertical maps are isomorphisms. Hence, for this X, d_r is completely determined by d'_r through $d_r = d'_r \cdot \tau^r$. Taking homology, it follows by induction that $E_r^{p,q,t} = F_r^{p,t} \cdot \tau^q$ for all $r < \infty$. Hence, this also holds for $r = \infty$. Since both spectral sequences are strongly convergent, we get:

Proposition 8.7.3. Let Y be a finite CW spectrum. Then

$$AK(n)^{**}(c(Y)) = 0 \Leftrightarrow K(n)^{*}(Y) = 0.$$

We have proven:

Theorem 8.7.4. (Constant type-n spectra have motivic type n)

The spectrum cX_n , where X_n denotes Ravenel's type-n spectrum (or any other p-local finite type-n spectrum), has motivic type n.

This and the previous sections can be summarised by:

Corollary 8.7.5. Let $X = cX_n$ or $X = \mathbb{X}_n$, where $\mathbb{X}_n = e_V(\mathbb{B}_{(p)})^{\wedge k_V}$. Then

thickid(X)
$$\subseteq \mathcal{C}_{AK(n-1)} \subseteq R^{-1}(\mathcal{C}_n)$$

is a chain of thick ideals in $(\mathcal{SH}(\mathbb{C})_f)_{(p)}$, and thickid $(X) \nsubseteq R^{-1}(\mathcal{C}_{n+1})$.

Recall that we derived the second of the above inclusions in Proposition 6.3.8.

As cK(n) is another motivic model for Morava K-theory (see Section 6.3.2), we can likewise ask the question, whether also thickid $(X) \subseteq C_{cK(n-1)}$. Let us first consider cX_n .

Proposition 8.7.6. $cK(s) \wedge c(X_n) \cong 0$ if and only if s < n. Hence,

thickid
$$(cX_n) \subseteq \mathcal{C}_{cK(n-1)} \subseteq R^{-1}(\mathcal{C}_n)$$

in $(\mathcal{SH}(\mathbb{C})_f)_{(p)}$. Furthermore, thickid $(cX_n) \not\subseteq R^{-1}(\mathcal{C}_{n+1})$.

Proof. We have $c(K(s) \wedge X_n) \cong 0$ if and only if X_n is in \mathcal{C}_{s+1} , since c is fully faithful by [Lev14, Theorem 1]. Since $X_n \in \mathcal{C}_n \setminus \mathcal{C}_{n+1}$, the first claim follows. The second claim holds because $R(cX_n) = X_n$.

For \mathbb{X}_n , we know so far:

Lemma 8.7.7. If $s \ge n$, then $cK(s) \land \mathbb{X}_n \not\cong 0$.

Proof. Assume $cK(s) \wedge \mathbb{X}_n \cong 0$. Then $R(cK(s) \wedge \mathbb{X}_n) \cong K(s) \wedge X_n \cong 0$ and it follows s < n.

For now, the question whether $cK(s) \wedge \mathbb{X}_n \cong 0$ for s < n remains open. If it is true, then thickid $(\mathbb{X}_n) \subseteq \mathcal{C}_{cK(n-1)} \subseteq R^{-1}(\mathcal{C}_n)$. If it is not true, then thickid $(\mathbb{X}_n) \notin \mathcal{C}_{cK(n-1)}$ and in particular $\mathbb{X}_n \notin$ thickid (cX_n) and thickid $(\mathbb{X}_n) \neq$ thickid (cX_n) .

Summary.

We have constructed two different lifts of a topological type-*n* spectrum to the motivic category $(\mathcal{SH}(\mathbb{C})_f)_{(p)}$, one of them is in $c(\mathcal{SH}_{(p)}^{fin})$ and the other one is not. These are candidates for generators of different thick sub-ideals of $R^{-1}(\mathcal{C}_n)$ inside $(\mathcal{SH}(\mathbb{C})_f)_{(p)}$. We proved that both spectra have motivic type *n*, meaning that the thick ideals generated by them cannot be distinguished using motivic Morava K-theories AK(s). It is an open question, whether these ideals are equal or whether they can be distinguished by some other motivic model for Morava K-theory or any other method.

Chapter 9

Bousfield classes

So far, we have seen that the thick ideals $R_k^{-1}C_n$ form a descending chain and that $C_{AK(n-1)}$ is a thick ideal contained in $R_k^{-1}C_n$. However, we have not seen that the thick ideals $C_{AK(n)}$ form a descending chain themselves. The aim of this chapter is to prove this, at least for $k = \mathbb{C}$ and finite cell spectra. That is, we prove that $AK(n)_{**}X = 0$ implies $AK(n-1)_{**}X = 0$ for $X \in \mathcal{SH}(\mathbb{C})^{fin}$, where $n \geq 1$ and p > 2 (Theorem 9.6.4). As is done in topology [Rav84, Theorem 2.11], we will work in terms of Bousfield classes (Definition 9.0.1).

We proceed as follows. In the first two sections, k can be any subfield of \mathbb{C} . In Section 9.1, we show that v_n -torsion is also v_{n-1} -torsion (Theorem 9.1.1). Then we show that some basic results on Bousfield classes also apply to the motivic setting (Lemma 9.2.2). In Section 9.3, we construct a product on AP(n). Here, we need to know that $AP(n)_{**}$ vanishes in certain degrees, which we do if we assume $k = \mathbb{C}$, p > 2 and n > 0. We continue by showing that, for p > 2 and n > 0, $\langle AK(n) \rangle = \langle AB(n) \rangle$ in $\mathcal{SH}(\mathbb{C})$ (Corollary 9.4.18), passing from AK(n) to the cohomology theory $AB(n) = v_n^{-1}AP(n)$, which is slightly easier to understand. On the way, we need to compute a couple of things like $AP(m)^{**}AP(n)$, to construct stable operations $AP(n)^{**}(-) \to AP(n)^{**}(-)$ (Theorem 9.4.12). Here, the assumption $k = \mathbb{C}$ is also helpful, as we make explicit use of the formula $AP(n)_{**} \cong P(n)_*[\tau]$ (Lemma 6.3.7). An application of all these results is Theorem 9.5.1, where, for p > 2 and $k = \mathbb{C}$, we prove

$$\langle AE(n) \rangle = \bigvee_{0 \le i \le n} \langle AK(i) \rangle.$$

For the definitions of AK(n), AB(n), AP(n) and AE(n), see Definition 6.3.1.

Let's start with a definition of Bousfield classes.

Definition 9.0.1. Let k be a field and let $\mathcal{T} = S\mathcal{H}(k)$, $S\mathcal{H}(k)^{cell}$ or $S\mathcal{H}(k)^{fin}$. For any $E \in S\mathcal{H}(k)$, the class of all spectra $X \in \mathcal{T}$ satisfying $E_{**}X \neq 0$ is denoted by $\langle E \rangle$. We write $\langle E \rangle \leq \langle F \rangle$ if $\langle E \rangle$ is a subclass of $\langle F \rangle$. Meet and join of Bousfield classes are given by $\langle E \rangle \wedge \langle F \rangle = \langle E \wedge F \rangle$ and $\langle E \rangle \vee \langle F \rangle = \langle E \vee F \rangle$ [Rav84, Definition 1.20].

Remark 9.0.2. In [Rav84, Definition 1.19], $\langle E \rangle$ is defined to be the equivalence class of all F such that, for all X, $E_*X = 0$ if and only if $F_*X = 0$. Thus, $\langle E \rangle$ is determined by

the collection of all X such that $E_*X \neq 0$, as in the definition above. For $\mathcal{T} = \mathcal{SH}(k)^{cell}$ or $\mathcal{SH}(k)^{fin}$ and $E \in \mathcal{SH}(k)^{cell}$, $\langle E \rangle = \{X \in \mathcal{T} \mid E \land X \not\cong 0\} = \mathcal{C}_E$ by Proposition 6.2.2.

9.1 v_n -Torsion

In this section, we work in the category $\mathcal{SH}(k)$, $k \subseteq \mathbb{C}$. We prove the following theorem, refining [JY80, Theorem 0.1].

Theorem 9.1.1. Any v_n -torsion element in an $ABP_{**}ABP$ -comodule is also a v_{n-1} -torsion element.

Recall that $BP_*BP \cong BP_*\{t_{top}^E\}$, where $E = (e_1, e_2, \cdots)$ runs over all finite sequences of non-negative integers and $\deg(t_{top}^E) = \sum e_i(2p^i - 2)$, as is shown in [Ada74, Theorem II.16.1(ii)].

From [Vez01, Definition 5.3], we know that ABP is a commutative ring spectrum. It follows that $ABP_{**}ABP$ is an ABP_{**} -module. We describe its structure:

Lemma 9.1.2. (1) As a left ABP_{**} -module,

$$ABP_{**}ABP \cong ABP_{**}\{t^E\},\$$

where E runs over all finite sequences of non-negative integers,

$$\deg\left(t^{(e_1,e_2,\cdots)}\right) = \left(\sum_i e_i(2p^i - 2), \sum_i e_i(p^i - 1)\right),\$$

and $R_k(t^E) = t^E_{top}$ Consequently, as a right ABP_{**}-module,

$$ABP_{**}ABP \cong ABP_{**}\{c(t^E)\},\$$

where $c: ABP_{**}ABP \rightarrow ABP_{**}ABP$ is the conjugation, induced by the twist map $ABP \wedge ABP \rightarrow ABP \wedge ABP$.

(2) As a left ABP^{**} -module,

$$ABP^{**}ABP \cong ABP^{**}[[s^E]],$$

which is the completion of $ABP^{**}\{s^E\}$ under infinite sums. Here, $\deg(s^E) = \deg(t^E)$ and the s^E are the dual basis elements to t^E .

In particular, $ABP_{**}ABP$ is a flat ABP_{**} -module.

Proof. By Remark 6.3.3(4), [NSØ09, Proposition 9.1.(i)] applies to ABP, so we have $ABP_{**}ABP \cong ABP_{**} \otimes_{BP_*} BP_*BP$. Since $BP_*BP \cong BP_*\{t_{top}^E\}$, the first claim follows. As BP_*BP is projective over BP_* , [NSØ09, Proposition 9.7(i)] implies $ABP^{**}ABP \cong Hom_{BP_*}(BP_*BP, ABP_{**})$. Since the analogue holds for BP^*BP , this is the same as $ABP^{**} \otimes_{BP^*} BP^*BP$, which is $ABP^{**}[[s^E]]$ by [JY80, Lemma 5.12]. □

For any finite motivic cell spectrum X, the morphism

$$m_*: ABP_{**}(ABP) \otimes_{ABP_{**}} ABP_{**}(X) \to ABP_{**}(ABP \land X)$$

induced by the ABP-module structure of $ABP \wedge X$ is an isomorphism: this holds for $X = S^0$, since $ABP_{**}ABP$ is free over ABP_{**} and, hence, for any finite X by cellular induction via the five lemma, see also [Ada69, Lecture 3, Lemma 1]. More precisely, one has to check that a cofiber sequence $X \to Y \to Z$ induces a long exact sequence on both ends of m_* . On the right hand side, this is the long exact ABP_{**} -sequence induced by $ABP \wedge X \to ABP \wedge Y \to ABP \wedge Z$ and on the left hand side, we get a long exact ABP_{**} -sequence tensored over the field \mathbb{F}_p with $\mathbb{F}_p\{t^E\}$, which is still exact. Actually, the above map is an isomorphism for any motivic spectrum X by [NSØ09, Lemma 5.1(i)].

This can be used to define elementary ABP-operations as in [JY80, Section 1]:

Definition 9.1.3. Let

$$\psi_X : ABP_{**}X \to ABP_{**}ABP \otimes_{ABP_{**}} ABP_{**}X$$

be the map induced by $1 \wedge i \wedge 1 : ABP \wedge S^0 \wedge X \to ABP \wedge ABP \wedge X$ (where *i* is the unit of the ring spectrum ABP) followed by $(m_*)^{-1}$. Then the elementary ABP-operation $s_E : ABP_{**}X \to ABP_{**}X$ is defined by

$$\psi_X(x) = \sum_E c(t^E) \otimes s_E(x).$$

The $s^E \in ABP^{**}ABP$ from the above lemma are special cases of these operations (see [JY80, Lemma 5.12]). The ABP-operations satisfy a Cartan formula similar to [JY80, Formula (1.7)]:

Lemma 9.1.4. If $y \in ABP_{**}$ and $x \in ABP_{**}X$, then

$$s_E(yx) = \sum_{F+G=E} s_F(y) s_G(x).$$

Proof. We have to show that

$$\psi_X(yx) = \sum_E \left(c(t^E) \otimes \sum_{F+G=E} s_F(y) s_G(x) \right).$$

Since ψ_X is a map of ABP_{**} -modules, $\psi_X(yx) = \psi_{S^0}(y)\psi_X(x)$, which is equal to $\sum_E \sum_{F+G=E} (c(t^F)c(t^G) \otimes s_F(y)s_G(x))$. As F and G are exponent sequences, $t^F t^G = t^E$ and it follows that $\psi_X(yx) = \sum_E (c(t^E) \otimes \sum_{F+G=E} s_F(y)s_G(x))$.

The next lemma compares the Hopf algebroid structures of (BP_*, BP_*BP) and $(ABP_{**}, ABP_{**}ABP)$ and is closely related to [NSØ09, Section 5].

Lemma 9.1.5. $(ABP_{**}, ABP_{**}ABP)$ is a flat Hopf algebroid, and there is a map of Hopf algebroids $(BP_*, BP_*BP) \rightarrow (ABP_{**}, ABP_{**}ABP)$ such that the following hold: (1) $BP_* \rightarrow ABP_{**}$ is the inclusion into $\bigoplus ABP_{(2i,i)}$, mapping v_i^{top} to v_i .

- (2) $BP_*BP \to ABP_{**}ABP$ is the map $\stackrel{i}{BP_*}\{t_{top}^E\} \to ABP_{**}\{t^E\}$ given by (1) on BP_* and mapping t_{top}^E to t^E .
- (3) The map $\psi = \psi_{S^0}$ from Definition 9.1.3 is the coaction map of ABP_{**} as a left (ABP_{**}, ABP_{**}ABP)-comodule and, similarly, for the map ψ^{top} from [JY80]. Furthermore, the map of Hopf algebroids preserves the comodule structure in the sense that

commutes.

Proof. In [NSØ09, Corollary 5.2(i)], it is shown that $(E_{**}, E_{**}E)$ is a flat Hopf algebroid whenever E is a cellular ring spectrum and $E_{**}E$ is a flat E_{**} -module. This is the case for E = ABP by Remark 6.3.3 and Lemma 9.1.2(1). Furthermore, an orientation on E induces a map of Hopf algebriods $(MU_*, MU_*MU) \rightarrow (E_{**}, E_{**}E)$ by [NSØ09, Corollary 6.7], where $MU_* \rightarrow E_{**}$ is the map classifying the formal group law (FGL) given by the orientation on E and $MU_*MU \rightarrow E_{**}E$ classifies the strict isomorphism of formal group laws induced on $E \wedge E$ by the left and right units $E \rightarrow E \wedge E$.

If the FGL associated with the orientation of E is p-typical, the map of Hopf algebroids factors through (BP_*, BP_*BP) because BP_* and BP_*BP classify p-typical group laws and strict isomorphisms of p-typical group laws, respectively (see [Rav86, Appendix 2]). Recall that ABP is oriented by $MU_* \to MGL_{**} \to ABP_{**}$ and its FGL is p-typical because this factors as $MU_* \to BP_* \to ABP_{**}$ by the construction of ABP, where the latter map is as claimed in (1). Hence, we get a map of Hopf algebroids $(BP_*, BP_*BP) \to (ABP_{**}, ABP_{**}ABP)$ satisfying (1).

Before we pove (2), we will show the analogous statement for MGL. By [NSØ09, Corollary 6.7], as above, there is a map of Hopf algebroids

$$(MU_*, MU_*MU) \rightarrow (MGL_{**}, MGL_{**}MGL),$$

determined by the complex orientation on MGL. Let x be the orientation on $MGL \wedge MGL$ induced by the left unit $MGL \to MGL \wedge MGL$ and x' be the orientation induced by the right unit. By [NSØ09, Lemma 6.4.(ii)], $x' = \sum_{i\geq 0} b_i x^{i+1}$. The orientations x and x' correspond to formal group laws F_L and F_R and the formula implies that the b_i are the coefficients of the power series of the strict isomorphism φ between F_L and F_R (as in the proof of [Rav86, Theorem 4.1.11]). The same formula holds for the orientations on $MU \wedge MU$ by [Rav86, Lemma 4.1.8] and the strict isomorphism between the FGLs F_L^{top} and F_R^{top} therefore has coefficients b_i^{top} . By definition, b_i is the image of b_i^{top} under $MU_*[b_i^{\text{top}}] \cong MU_*MU \to MGL_{**} \otimes_{MU_*} MU_*MU \cong MGL_{**}MGL$, as in [NSØ09, Lemma 6.4.(i)].

 MU_*MU classifies strict isomorphisms $F \xrightarrow{f} G$ of FGLs in the following way: $MU_*MU \cong MU_*[b_i^{\text{top}}]$, where MU_* classifies F, f is given by a power series in b_i^{top} and G is determined by F and f.

In our setting, the map $MU_*MU \to MGL_{**}MGL$ is the map corresponding to the strict isomorphism $F_L \xrightarrow{\varphi} F_R$. Furthermore, F_L is the FGL associated with the orientation of $MGL_{**}MGL$ given by $MU_* \to MGL_{**} \to MGL_{**}[b_i] \cong MGL_{**}MGL$ (because the isomorphism herein is an isomorphism of left MGL_{**} -modules), and, similarly, F_L^{top} is the FGL associated with the orientation $MU_* \to MU_*[b_i^{\text{top}}] \cong MU_*MU$. This implies that the following square commutes, where the left horizontal maps are the obvious inclusions, and the vertical maps are the maps from [NSØ09, Corollary 6.7].

$$\begin{array}{ccc} MU_{*} \longrightarrow MU_{*}[b_{i}^{\mathrm{top}}] & \xrightarrow{\cong} & MU_{*}MU \\ & & & & & \\ & & & & & \\ MGL_{**} \longrightarrow MGL_{**}[b_{i}] & \xrightarrow{\cong} & MGL_{**}MGL. \end{array}$$

In terms of group laws, the right hand map sends φ^{top} to φ . Since these are the power series described above, b_i^{top} is sent to b_i . This proves the *MGL*-version of (2).

For the ABP-version, one has to show that the following diagram commutes, where the right map is the Hopf algebroid morphism and the left map is as described in (2).

The proof is exactly the same as in the case of MGL. One simply has to replace each FGL by the corresponding p-typical FGL.

For (3), note that, by its definition, ψ is the coaction map that comes naturally with any flat Hopf algebroid $(E_{**}, E_{**}E)$ (as in [NSØ09, Corollary 5.2(i)]), meaning in particular that the diagram in (3) commutes.

Definition 9.1.6. For an exponent sequence $E = (e_1, e_2, \cdots)$ as in Lemma 9.1.2(1), we set $|E| = \sum_i e_i (2p^i - 2)$. Let $I_m = (p, v_1, \cdots, v_{m-1}) \subset ABP_{**}$ be the usual prime ideal.

Corollary 9.1.7. Consider $s_E : ABP_{**} \to ABP_{**}$ as in Definition 9.1.3 and assume that $|E| \ge 2kp^s(p^n - p^m)$ for $n \ge m$, $s \ge 0$ and $k \ge 1$. Then

$$s_E(v_n^{kp^s}) = \begin{cases} v_m^{kp^s} & \mod I_m^{s+1} \text{ if } e_{n-m} = kp^{s+m} \text{ and } e_i = 0 \text{ for } i \neq n-m \\ 0 & \mod I_m^{s+1} \text{ otherwise.} \end{cases}$$

Proof. By the above Lemma, the following diagram commutes:

where the vertical arrows send v_n^{top} to v_n and t_{top}^E to t^E . We consider the element $(v_n^{\text{top}})^{kp^s} \in BP_*$. It is mapped horizontally to

$$\psi^{\mathrm{top}}((v_n^{\mathrm{top}})^{kp^s}) = \sum_E c^{\mathrm{top}}(t_{\mathrm{top}}^E) \otimes s_E^{\mathrm{top}}((v_n^{\mathrm{top}})^{kp^s}).$$

By [JY80, Lemma 2.1], $s_E^{\text{top}}((v_n^{\text{top}})^{kp^s})$ satisfies the formula we want to prove. Since all the elements from the topological case map to the corresponding elements in the lower row, the formula has to hold there, too.

The rest of the proof of Theorem 9.1.1 is exactly the same as [JY80, Lemmas 2.2 and 2.3], relying mainly on the above lemma and the Cartan formula. Theorem 9.1.1 implies the following corollary. The analogous topological statement can be found in the proof of [Rav84, Theorem 2.1(d)].

Corollary 9.1.8. Let $k \subseteq \mathbb{C}$. If $AE(n)_{**}X = 0$, then also $AE(i)_{**}X = 0$ for all $i \leq n$. In terms of Bousfield classes in SH(k):

$$\langle AE(n) \rangle \ge \langle AE(i) \rangle$$
 for all $n \ge i$.

Proof. Since E(n) is Landweber exact (see [Lan76] or [Rav86, Section 4.2]), the ABPversion of [NSØ09, Theorem 8.7] applies to $AE(n)_{**}(X)$, yielding

$$AE(n)_{**}(X) \cong ABP_{**}(X) \otimes_{BP_*} E(n)_{*},$$

which is an $ABP_{**}ABP$ -comodule via the map ψ_X from Definition 9.1.3.

which is an $ABP_{**}ABP$ -comodule via the map ψ_X from Definition 9.1.3. As $E(n) = (v_{n}^{\text{top}})^{-1}BP/(v_{n+1}^{\text{top}}, v_{n+2}^{\text{top}}, \cdots)$, the condition $AE(n)_{**}(X) = 0$ is equivalent to $ABP_{**}(X) \otimes_{BP_*} BP/(v_{n+1}^{\text{top}}, v_{n+2}^{\text{top}}, \cdots)$ being v_n -torsion. By Theorem 9.1.1, it follows that $ABP_{**}(X) \otimes_{BP_*} BP/(v_{n+1}^{\text{top}}, v_{n+2}^{\text{top}}, \cdots)$ is v_i -torsion for any $i \leq n$. This implies that also $ABP_{**}(X) \otimes_{BP_*} BP/(v_{i+1}^{\text{top}}, v_{i+2}^{\text{top}}, \cdots)$ is v_i -torsion, which is equivalent to AE(i) = 0 $AE(i)_{**}(X) = 0.$

9.2 **Properties of Bousfield classes**

Ravenel has shown the following properties of Bousfield classes [Rav84, Section 1]. They hold in any tensor triangulated category (\mathcal{T}, \wedge) .

Lemma 9.2.1. (1) In an exact triangle, each Bousfield class is less or equal to the wedge of the other two [Rav84, Proposition 1.23].

- (2) If M is a module spectrum over the ring spectrum E, then $\langle M \rangle \leq \langle E \rangle$ [Rav84, Proposition 1.24].
- (3) Let Σ be an auto-equivalence in \mathcal{T} . If Y is the homotopy cofiber of $\Sigma^d X \xrightarrow{f} X$ and $\hat{X} = \operatorname{colim}(\Sigma^{-kd}X)$, then $\langle X \rangle = \langle \hat{X} \rangle \lor \langle Y \rangle$ [Rav84, Lemma 1.34].

Furthermore, the following relations from [Rav84, Section 2] also hold in $\mathcal{SH}(k)$:

Lemma 9.2.2. (1) $\langle AE(n) \rangle \ge \langle AK(n) \rangle$, (2) $\langle AE(n) \rangle \land \langle AP(n+1) \rangle = \langle v_n^{-1}ABP \rangle \land \langle AP(n+1) \rangle = \langle 0 \rangle$, (3) $\langle AP(n) \rangle = \langle AB(n) \rangle \lor \langle AP(n+1) \rangle$.

Proof. Constructing AK(n) from AE(n), (1) follows from Lemma 9.2.1(1). (3) is a direct application of Lemma 9.2.1(3) (see also [Rav84, Theorem 2.1(c)]). For the first part of (2), note that $\langle AE(n) \rangle \leq \langle v_n^{-1}ABP \rangle$ since $AE(n) = v_n^{-1}ABP/(v_{n+1}, v_{n+2}, \cdots)$. When we prove the second equation in (2), the first one will follow from this inequality because $\langle 0 \rangle$ is the empty set.

It remains to show that $AP(n+1) \wedge v_n^{-1}ABP \cong 0$, as proven in the topological setting in [Rav84, Lemma 2.3]. This spectrum is the homotopy cofiber of the map

$$v_n \wedge 1 : AP(n) \wedge v_n^{-1}ABP \to AP(n) \wedge v_n^{-1}ABP.$$

We claim that $(v_n \wedge 1)_* = (1 \wedge v_n)_*$ on $\pi_{**}(AP(n) \wedge ABP)$. Since $(1 \wedge v_n)_*$ is an isomorphism on $\pi_{**}(AP(n) \wedge v_n^{-1}ABP)$, this will imply that the homotopy cofiber is contractible.

To prove this claim, note that $(v_n \wedge 1)_*$ and $(1 \wedge v_n)_*$ are induced by the respective maps on $\pi_{**}(ABP \wedge ABP)$, where they are given by applying the left respectively right unit $ABP_{**} \rightarrow ABP_{**}ABP$ to v_n . In the topological case, the left and right units applied to v_n are the same modulo I_n by [Ada74, II.16.1 (ii)]. By Lemma 9.1.2(1) and the inclusion of BP_* in ABP_{**} , this also holds motivically. Hence, $(v_n \wedge 1)_*$ and $(1 \wedge v_n)_*$ are the same modulo I_n . It remains to show that $I_n \subseteq ABP_{**}ABP$ maps to 0 under $ABP_{**}ABP \rightarrow AP(n)_{**}ABP$. For n = 0, there is nothing to show. Assume that I_n is mapped to zero in $AP(n)_{**}ABP$ for some n. Consider the map $AP(n)_{**}ABP \xrightarrow{i_*}$ $AP(n+1)_{**}ABP$ induced by the map to the cofiber in $AP(n) \xrightarrow{v_n} AP(n) \xrightarrow{i} AP(n+1)$. The inductive assumption implies that I_n is still zero in $AP(n+1)_{**}ABP$. Recall that $I_{n+1} \subseteq ABP_{**}ABP$ is the ideal generated by I_n and v_n . Since $i_* \circ (v_n)_* = 0$ in the long exact sequence

$$\cdots \to AP(n)_{**}ABP \xrightarrow{(v_n)_*} AP(n)_{**}ABP \xrightarrow{i_*} AP(n+1)_{**}ABP \to \cdots,$$

 i_* maps $v_n = (v_n)_*(1)$ to 0. Hence, $I_{n+1} = 0$ in $AP(n+1)_{**}ABP$.

9.3 The action of v_i on AP(n)

In [JW75, Appendix], a geometric proof using the Baas-Sullivan construction of P(n) shows that the action of v_i^{top} on $P(n)_*(X)$ is zero for any $0 \le i < n$. A non-geometric

proof of this result is given by [Nas95, Satz 1.3.4], which was motivated by [Wür77]. Nassau's proof can be simplified using the language of triangulated categories of modules, which is basically done in [Elm+97, Lemma V.2.4], as well as in [Str99, Lemma 3.2]. These proofs rely on the fact that the v_i are non-zero divisors of MU_* and that BP_* vanishes in certain degrees, which is not known in the motivic case. In the following, we will use ideas from [Nas95] and [Elm+97] to give a proof which also works in $S\mathcal{H}(\mathbb{C})$. The main difference is that we only know coefficients after passing to $MGL_{(p)}/(p)$ (see Lemma 6.3.7), which is why we have to work with R/(x, y), while [Elm+97, Chapter V] only works with R/x for some ring spectrum R.

In this section, we will prove that v_i acts trivially on AP(n) if $k = \mathbb{C}$. Furthermore, we will show that if p is odd, then $AP(n)_{**}(X)$ and $AP(n)^{**}(X)$ are $AP(n)_{**}$ -modules for any $X \in S\mathcal{H}(\mathbb{C})$.

Recall that MGL can be constructed as an E_{∞} -ring spectrum [Hu03, Theorem 14.2], which is equivalent to a strictly commutative ring spectrum by the motivic version of [Elm+97, Corollary II.3.6].

Let $R \in S\mathcal{H}(k)$ be a strictly commutative ring spectrum with multiplication $m : R \wedge R \to R$ and unit $i : S^{0,0} \to R$. Let $x : S^{k,l} \to R$ for some $k, l \in \mathbb{Z}$. In our application, we will have $R = MGL_{(p)}$. Note that $MGL_{(p)}$ is the homotopy colimit of the diagram of maps $MGL \xrightarrow{n} MGL$ for all positive integers relatively prime to p (see [Hu03, end of Section 14]). As these are maps of strictly commutative ring spectra, $MGL_{(p)}$ is also a strictly commutative ring spectrum ([SS00, Theorem 4.1(3)] applied to $S^{0,0}$ -algebras implies that the category of strictly commutative ring spectra is cocomplete).

Let M be an R-module with action map $\nu_M : R \wedge M \to M$. Let

$$\phi = \nu_M \circ (x \wedge 1_M) : S^{k,l} \wedge M \to R \wedge M \to M.$$

The map ϕ is the action of x on M.

The *R*-module structure on $S^{k,l} \wedge M$ is given by

$$\nu_{S^{k,l} \wedge M} : R \wedge S^{k,l} \wedge M \xrightarrow{\tau \wedge 1_M} S^{k,l} \wedge R \wedge M \xrightarrow{1_{S^{k,l} \wedge \nu_M}} S^{k,l} \wedge M.$$

Lemma 9.3.1. The map ϕ is an *R*-module map.

Proof. We have to check the commutativity of the following diagram:

$$\begin{array}{c|c} R \wedge S^{k,l} \wedge M & \xrightarrow{1 \wedge x \wedge 1} & R \wedge R \wedge M \xrightarrow{1 \wedge \nu_M} R \wedge M \\ & & & \downarrow \\ & & & \downarrow \\ & & & \downarrow \\ S^{k,l} \wedge R \wedge M \xrightarrow{1 \wedge \nu_M} S^{k,l} \wedge M \xrightarrow{x \wedge 1} R \wedge M \xrightarrow{\nu_M} M. \end{array}$$

In this diagram, we can replace $(x \wedge 1) \circ (1 \wedge \nu_M)$ by

$$S^{k,l} \wedge R \wedge M \xrightarrow{x \wedge 1 \wedge 1} R \wedge R \wedge M \xrightarrow{1 \wedge \nu_M} R \wedge M$$

and we can fill in a diagonal across the upper left corner,

$$S^{k,l} \wedge R \wedge M \xrightarrow{(\tau \wedge 1) \circ (x \wedge 1 \wedge 1)} R \wedge R \wedge M.$$

It follows that the above diagram commutes if and only if the following diagram commutes:

$$\begin{array}{c|c} S^{k,l} \wedge R \wedge M \xrightarrow{x \wedge 1 \wedge 1} R \wedge R \wedge M \xrightarrow{\tau \wedge 1} R \wedge R \wedge M \xrightarrow{1 \wedge \nu_M} R \wedge M \\ & & \downarrow^{\nu_M} \\ R \wedge R \wedge M \xrightarrow{1 \wedge \nu_M} R \wedge M \xrightarrow{\nu_M} M. \end{array}$$

Since R is commutative, we have $m \circ \tau = m$ and, hence,

$$\nu_M \circ (m \wedge 1) \circ (\tau \wedge 1) = \nu_M \circ (m \wedge 1).$$

Since M is an R-module, this is the same as

$$\nu_M \circ (1 \wedge \nu_M) \circ (\tau \wedge 1) = \nu_M \circ (1 \wedge \nu_M),$$

proving the commutativity of the above diagram.

In the following, we denote the homotopy category of *R*-modules by *R*-Mod. A stable model structure on *R*-modules is given by [SS00, Theorem 4.1] applied to the motivic stable model structure from [Jar00], so that *R*-Mod is a triangulated category (compare [NSØ09, page 554]). Since ϕ from above is a map of *R*-modules (Lemma 9.3.1), there is an exact triangle in *R*-Mod,

$$S^{k,l} \wedge M \xrightarrow{\phi} M \xrightarrow{\eta} N \xrightarrow{\partial} S^{k+1,l} \wedge M.$$

The cofiber N is also denoted M/x. Application of $[-, N]_{R-Mod}$ to this exact triangle yields a long exact sequence

$$\cdots \to [S^{2k+1,2l} \land M, N]_{R-\operatorname{Mod}} \xrightarrow{\partial^*} [S^{k,l} \land N, N]_{R-\operatorname{Mod}} \xrightarrow{\eta^*} [S^{k,l} \land M, N]_{R-\operatorname{Mod}} \to \cdots$$

Let $\psi = \nu_N \circ (x \wedge 1_N) : S^{k,l} \wedge N \to R \wedge N \to N$. This map is the action of x on M/x, and a map of R-modules by Lemma 9.3.1. We want to show that, under certain assumptions, $\psi = 0$, meaning that x acts trivially on M/x.

First, we consider

$$\eta^* \psi = \psi \circ (1_{S^{k,l}} \land \eta) : S^{k,l} \land M \xrightarrow{1 \land \eta} S^{k,l} \land N \xrightarrow{\psi} N.$$

By the definition of ψ , this is the map $\nu_N \circ (x \wedge 1_N) \circ (1_{S^{k,l}} \wedge \eta) = \nu_N \circ (1_R \wedge \eta) \circ (x \wedge 1_M)$. Since η is a map of *R*-modules, this is the same as $\eta \circ \nu_M \circ (x \wedge 1_M)$, which, by definition of ϕ , is the map $\eta \circ \phi$. By the above exact triangle, it follows that $\eta^* \psi = \eta \circ \phi = 0$. The long exact sequence implies that there is a map in *R*-Mod,

$$\overline{\psi}: S^{2k+1,2l} \wedge M \to N,$$

such that $\psi = \partial^* \overline{\psi}$.

Now, we assume that either M = R or that M = R/y for some $y \in R_{**}$. Furthermore, we assume that $\pi_{2k+1,2l}N = 0$.

Note that Case 1 is a special instance of Case 2 (with y = 0), so the reader may skip the following paragraph and continue reading at Case 2. However, Case 1 is easier, for which reason it might still be a good idea to read it, anyway.

Case 1. M = R.

We have $\overline{\psi}: S^{2k+1,2l} \wedge R \to N$. Since R is a ring spectrum, the unit $i: S^{0,0} \to R$ satisfies $1_R = m \circ (1_R \wedge i): R \wedge S^{0,0} \to R \wedge R \to R$. Hence,

$$\overline{\psi} = \overline{\psi} \circ (1_{S^{2k+1,2l}} \wedge m) \circ (1_{S^{2k+1,2l} \wedge R} \wedge i).$$

By the definition of the *R*-module structure $\nu_{S^{2k+1,2l}\wedge R}$ on $S^{2k+1,2l}\wedge R$,

 $(1_{S^{2k+1,2l}} \wedge m) = \nu_{S^{2k+1,2l} \wedge R} \circ (\tau \wedge 1_R) : S^{2k+1,2l} \wedge R \wedge R \to R \wedge S^{2k+1,2l} \wedge R \to S^{2k+1,2l} \wedge R.$

Hence,

$$\begin{split} \psi &= \psi \circ \nu_{S^{2k+1,2l} \wedge R} (\tau \wedge 1_R) (1_{S^{2k+1,2l} \wedge R} \wedge i) \\ &= \overline{\psi} \circ \nu_{S^{2k+1,2l} \wedge R} (1_R \wedge 1_{S^{2k+1,2l} \wedge i}) (\tau \wedge 1_R). \end{split}$$

 $= \psi \circ \nu_{S^{2k+1,2l} \wedge R} (1_R \wedge 1_{S^{2k+1,2l}} \wedge i) (\tau \wedge 1_R).$ Since $\overline{\psi}$ is an *R*-module map, $\overline{\psi} \nu_{S^{2k+1,2l} \wedge R} = \nu_N (1_R \wedge \overline{\psi})$, and therefore

$$\overline{\psi} = \nu_N (1_R \wedge \overline{\psi}) (1_R \wedge 1_{S^{2k+1,2l}} \wedge i) (\tau \wedge 1_R)$$
$$= \nu_N (1_R \wedge (\overline{\psi}(1_{S^{2k+1,2l}} \wedge i))) (\tau \wedge 1_R).$$

Now, $\overline{\psi}(1_{S^{2k+1,2l}} \wedge i) : S^{2k+1,2l} \wedge S^{0,0} \to N$ is in $\pi_{2k+1,2l}N$, which we assumed to be zero. Thus, $\overline{\psi} = 0$ and it follows that also $\psi = \partial^* \overline{\psi}$, which is the action of x on M/x, is zero, as we wanted to show.

Thus, we have shown:

Proposition 9.3.2. Let $R \in S\mathcal{H}(k)$ be a strictly commutative ring spectrum and $x \in \pi_{k,l}R$. Assume that $\pi_{2k+1,2l}(R/x) = 0$. Then x acts trivially on R/x, i.e., the map ψ from above is zero. The same holds if $x \in \pi_k R$ for a strictly commutative ring spectrum $R \in S\mathcal{H}$ such that $\pi_{2k+1}(R/x) = 0$.

Note that part of the above argument can be formulated more generally:

Lemma 9.3.3. Let R be a (homotopy) ring spectrum, M a left R-module, and $\pi_{k,l}M = 0$. Then any R-module map $\psi : S^{k,l} \wedge R \to M$ is homotopically trivial.

Proof. Let $i: S^{0,0} \to R$ be the unit of R. It satisfies $1_R = m(1_R \wedge i)$. Thus,

$$\psi = \psi \circ m(1_R \wedge i) = \nu_M(1_R \wedge \psi)(1_R \wedge i) = \nu_M(1_R \wedge \psi i),$$

with $\psi i \in \pi_{k,l} M = 0$. It follows $\psi = \nu_M (1_R \wedge 0) = 0$.

Now we pass on to case 2.

Case 2. M = R/y.

Let $y: S^{k',l'} \to R$ and let $\phi' = m(y \wedge 1_R): S^{k',l'} \wedge R \to R \wedge R \to R$ be the action of y on R. We have an exact triangle in R-Mod,

$$S^{k',l'} \wedge R \xrightarrow{\phi'} R \xrightarrow{\eta'} M \xrightarrow{\partial'} S^{k'+1,l'} \wedge R,$$

and, again, an exact sequence

$$\rightarrow [S^{2k+k'+2,2l+l'} \wedge R, N]_{R-\operatorname{Mod}} \xrightarrow{\partial^{*}} [S^{2k+1,2l} \wedge M, N]_{R-\operatorname{Mod}} \xrightarrow{\eta^{*}} [S^{2k+1,2l} \wedge R, N]_{R-\operatorname{Mod}} \rightarrow \cdots$$

We consider

$$\eta'^*\overline{\psi} = \overline{\psi} \circ (1_{S^{2k+1,2l}} \wedge \eta') : S^{2k+1,2l} \wedge R \to S^{2k+1,2l} \wedge M \to N.$$

Let $i: S^{0,0} \to R$ be the unit of R, as before. Since $\eta': R \to M$ is a map of R-modules (using Lemma 9.3.1), $\eta' \circ m = \nu_M(1_R \land \eta')$, and, hence,

$$\nu_M(1_R \wedge \eta')(1_R \wedge i) = \eta' \circ m(1_R \wedge i) = \eta'.$$

Thus,

$$\overline{\psi}(1_{S^{2k+1,2l}} \wedge \eta') = \overline{\psi}(1_{S^{2k+1,2l}} \wedge \nu_M(1_R \wedge \eta')(1_R \wedge i))$$
$$= \overline{\psi}(1_{S^{2k+1,2l}} \wedge \nu_M(1_R \wedge \eta'i)).$$

Since $\overline{\psi}$ is a map of *R*-modules, $\overline{\psi} \circ \nu_{S^{2k+1,2l} \wedge M} = \nu_N \circ (1_R \wedge \overline{\psi})$, where, by definition, $\nu_{S^{2k+1,2l} \wedge M} = (1_{S^{2k+1,2l}} \wedge \nu_M)(\tau \wedge 1_M)$. Hence, $\overline{\psi}(1_{S^{2k+1,2l}} \wedge \nu_M) = \nu_N(1_R \wedge \overline{\psi})(\tau \wedge 1_M)$, and, therefore,

$$egin{aligned} &\eta'^*\overline{\psi} =
u_N(1_R\wedge\overline{\psi})(au\wedge 1_M)(1_{S^{2k+1,2l}}\wedge 1_R\wedge\eta'i)\ &=
u_N(1_R\wedge\overline{\psi})(1_R\wedge 1_{S^{2k+1,2l}}\wedge\eta'i)(au\wedge 1_{S^{0,0}})\ &=
u_N(1_R\wedge(\overline{\psi}(1_{S^{2k+1,2l}}\wedge\eta'i)))(au\wedge 1_{S^{0,0}}). \end{aligned}$$

Now, $\overline{\psi}(1_{S^{2k+1,2l}} \wedge \eta' i) : S^{2k+1,2l} \wedge S^{0,0} \to N$ lies in $\pi_{2k+1,2l}N$, which we assumed to be zero. Hence, $\eta'^*\overline{\psi} = 0$. By the long exact sequence from above, it follows that $\overline{\psi} = \partial'^*\overline{\overline{\psi}}$ for some *R*-module map $\overline{\overline{\psi}} : S^{2k+k'+2,2l+l'} \wedge R \to N$. Thus, $\psi = \partial^*\overline{\psi} = \partial^*\partial'^*\overline{\overline{\psi}}$.

Consider the following commutative diagram. The map ψ is the precomposition of $\overline{\overline{\psi}}$ with the diagonal of the righthand square.

$$S^{2k,2l} \wedge M \xrightarrow{\phi} S^{k,l} \wedge M \xrightarrow{\eta} S^{k,l} \wedge N \xrightarrow{\partial} S^{2k+1,2l} \wedge M$$

$$\downarrow^{j} \qquad \downarrow^{\xi} \qquad \downarrow^{j} \qquad \downarrow^{j}$$

Since both rows are exact triangles, we can fill in a map $\zeta : S^{k,l} \wedge M \to S^{2k+1,2l} \wedge R$. We have $\phi' \circ \partial' = 0$, as both of these are maps in the lower triangle. Thus, the diagonal in the first square is zero and the map ζ lifts to a map $\xi : S^{k,l} \wedge N \to S^{2k+1,2l} \wedge R$. It follows that $\partial' \circ \partial = \partial' \circ \eta' \circ \xi = 0$, and, hence, $\psi = \overline{\psi} \circ \partial' \circ \partial = 0$.

We have proven:

Proposition 9.3.4. Let $R \in S\mathcal{H}(k)$ be a commutative ring spectrum, $x \in \pi_{k,l}R$ and $y \in \pi_{k',l'}R$. Assume that $\pi_{2k+1,2l}(R/(y,x)) = 0$. Then x acts trivially on R/(y,x).

The same holds if $x \in \pi_k R$ and $y \in \pi_{k'} R$ for a commutative ring spectrum $R \in SH$ such that $\pi_{2k+1}(R/(y,x)) = 0$.

This result can be applied to the action of v_i on AP(n) for $0 \le i < n$, at least for $k = \mathbb{C}$.

Corollary 9.3.5. Let $k = \mathbb{C}$ and $n \ge 1$. Then v_i acts trivially on AP(n) for any $0 \le i < n$ and v_i acts trivially on Ak(n) for any $i \ne n$.

Proof. First, we consider $MGL_{(p)}/(p, v_i)$ for some 0 < i < n (thus, $n \ge 2$). Since MU_* is concentrated in even degrees and Lemma 6.3.7 holds for $k = \mathbb{C}$ and quotients of $MGL_{(p)}/p$, we get $\pi_{2k+1,2l}(MGL_{(p)}/(p, v_i)) = 0$ for any k, l. By Proposition 9.3.4 it follows that $v_0 = p$ and v_i act trivially on $MGL_{(p)}/(p, v_i)$.

By [Elm+97, Lemma V.1.10],

$$MGL_{(p)}/(p, v_i) \cong MGL_{(p)}/p \wedge_{MGL_{(p)}} MGL_{(p)}/v_i$$

and, by [Hoy13, Remark 6.20], $AP(n) \cong MGL_{(p)}/J$, where J contains $a_i \in MGL_{**}$, $i \neq 2p^i - 2$, as well as $v_i, 0 \leq i \leq n - 1$. From [Elm+97, Lemma V.1.10], it follows that

$$AP(n) \cong MGL_{(p)}/(p, v_i) \wedge_{MGL_{(p)}} MGL_{(p)}/(J \setminus \{p, v_i\}).$$

Now v_i acts trivially on $MGL_{(p)}/(p, v_i)$, i.e., the respective map ϕ_i on $MGL_{(p)}/(p, v_i)$ is zero. It follows that also the map $\phi_i \wedge_{MGL_{(p)}} 1_{MGL_{(p)}/(J \setminus \{p, v_i\})}$ is zero, meaning that v_i acts trivially on AP(n). Similarly, p acts trivially on AP(n). This proves that all v_i , $0 \leq i < n$, act trivially on AP(n) if $n \geq 2$.

If n = 1, one has to replace $MGL_{(p)}/(p, v_i)$ by $MGL_{(p)}/(p)$ in the above argument. Furthermore,

$$Ak(n) \cong AP(n) \wedge_{MGL_{(p)}} MGL_{(p)}/(p, v_{n+1}, v_{n+2}, \cdots),$$

so v_i acts trivially on Ak(n), too, for $0 \le i < n$. For i > n, the claim follows analogously to the above argument.

Remark 9.3.6. Working with modules over $MGL_{(p)}$ has the advantage that the E_{∞} -structure allows us to use the isomorphism from [Elm+97, Lemma V.1.10], as well as the results from [SS00] (below Lemma 9.3.1). For ABP, we only know of a commutative ring structure in the weak sense (see [Vez01, Definition 5.3]). Note that for BP, an E_4 -structure is constructed in [BM13].

The *ABP*-module structure on AP(n) is the action of *ABP* on itself in $ABP \wedge_{MGL_{(p)}}$ $MGL_{(p)}/(v_0, \dots, v_{n-1}) \cong AP(n)$ and it is (by its construction) compatible with the $MGL_{(p)}$ -action on AP(n). Recall that $\nu : BP \wedge P(n) \to P(n)$ induces a BP_* -module structure on $P(n)_*(X)$ and $P(n)^*(X)$ for any $X \in S\mathcal{H}$ and that $P(n)_* = BP_*/(v_0, \dots, v_{n-1})$. Therefore, the classical version of the above corollary immediately implies that the BP_* -module structure on $P(n)_*(X)$ and $P(n)^*(X)$ induces a $P(n)_*$ -module structure, as also concluded in [JW75, Remark 2.5(a)].

Our next aim is to show that also for $X \in S\mathcal{H}(\mathbb{C})$, the ABP_{**} -module structure on $AP(n)_{**}(X)$ and $AP(n)^{**}(X)$ induces a structure of $AP(n)_{**}$ -modules, where the ring structure on $AP(n)_{**}$ is defined via the isomorphism $AP(n)_{**} \cong H_{**} \otimes_{\mathbb{F}_p} P(n)_*$ (Lemma 6.3.7). We will show in Lemma 9.3.10 that this is the right choice of ring structure on $AP(n)_{**}$.

Let R be a strictly commutative ring spectrum, let M = R/y satisfy $\pi_{2k'+1,2l'}M = 0$ (where (k', l') is the degree of y, as in Case 2 above), and let N = M/x satisfy $\pi_{2k+1,2l}N = 0$.

In the commutative diagram,

$$\begin{array}{c|c} S^{k',l'} \wedge R \wedge M^{\underbrace{y \wedge 1 \wedge 1}} & R \wedge R \wedge M \xrightarrow{m \wedge 1} R \wedge M \\ 1 \wedge \nu_M & & \downarrow \\ S^{k',l'} \wedge M \xrightarrow{\eta \wedge 1} R \wedge M \xrightarrow{\nu_M} M, \end{array}$$

the composition $\nu_M(y \wedge 1_M)$ is zero by Proposition 9.3.2. Furthermore, $(m \wedge 1_M) \circ (y \wedge 1_R \wedge 1_M) = \phi' \wedge 1_M$, where ϕ' is, as before, the map whose cofiber is M. Thus, we have

$$S^{k',l'} \wedge R \wedge M \xrightarrow{\phi' \wedge 1} R \wedge M \xrightarrow{\eta' \wedge 1} M \wedge M$$

and there exists a map $\mu_M : M \wedge M \to M$ in the homotopy category *R*-Mod such that $\mu_M \circ (\eta' \wedge 1_M) = \nu_M$.

Next, we define a map $\nu_{M,N} : M \wedge N \to N$ by $\nu_{M,N} = \mu_M \wedge_R \mathbf{1}_{R/x}$, using $N \cong M \wedge_R R/x$ [Elm+97, Lemma V.1.10]. It satisfies $\nu_{M,N}(\eta' \wedge \mathbf{1}_N) = \nu_N$ (by applying $- \wedge_R R/x$ to the analogous equation for μ_M) and $\nu_{M,N}(1 \wedge \eta) = \eta \circ \mu_M$ (because $\eta : M \to N$ is the canonical map $M \wedge_R R \to M \wedge_R R/x$).

In the commutative diagram (where the right square commutes because $\nu_{M,N}$ is a map of *R*-modules)

$$\begin{array}{c|c} S^{k,l} \wedge M \wedge N \xrightarrow{x \wedge 1 \wedge 1} & R \wedge M \wedge N \xrightarrow{\nu_M \wedge 1} & M \wedge N \\ 1 \wedge \nu_{M,N} \downarrow & & 1 \wedge \nu_{M,N} \downarrow & & \downarrow \nu_{M,N} \\ S^{k,l} \wedge N \xrightarrow{x \wedge 1} & R \wedge N \xrightarrow{\nu_N} & N, \end{array}$$

the lower composition is the action of x on N, which is trivial by Proposition 9.3.4. Thus,

 $\nu_{M,N}(\phi \wedge 1_N) = \nu_{M,N}(\nu_M \wedge 1_N)(x \wedge 1_M \wedge 1_N) = 0.$

Hence, there exists a map $\mu_N : N \wedge N \to N$ in *R*-Mod making the following diagram commutative.

$$S^{k,l} \wedge M \wedge N \xrightarrow{\phi \wedge 1} M \wedge N \xrightarrow{\eta \wedge 1} N \wedge N$$

In particular, this applies to $N = MGL_{(p)}/(p, x)$ as in Corollary 9.3.5, yielding an $MGL_{(p)}$ -module map

$$\mu_x: MGL_{(p)}/(p,x) \land MGL_{(p)}/(p,x) \to MGL_{(p)}/(p,x).$$

Lemma 9.3.7. AP(n) is isomorphic as an $MGL_{(p)}$ -module to the $\wedge_{MGL_{(p)}}$ -product of all $MGL_{(p)}/(p,x)$, $x \in J$, where J is as in the proof of Corollary 9.3.5.

Proof. By Proposition 9.3.2, p acts trivially on $MGL_{(p)}/p$, proving $MGL_{(p)}/(p, p) = MGL_{(p)}/p$. By [Elm+97, Lemma V.1.10], it follows that

$$MGL_{(p)}/(p,x) \wedge_{MGL_{(p)}} MGL_{(p)}/(p,y) \cong MGL_{(p)}/(p,x) \wedge_{MGL_{(p)}} MGL_{(p)}/y$$

for any $x, y \in J$. With $p = v_0 \in J$, this implies that the $\wedge_{MGL_{(p)}}$ -product of all $MGL_{(p)}/(p,x)$ is isomorphic to the $\wedge_{MGL_{(p)}}$ -product of all $MGL_{(p)}/x$, which is the quotient $MGL_{(p)}/J \cong AP(n)$ (as in Corollary 9.3.5).

We can, therefore, define a map of $MGL_{(p)}$ -modules,

$$\mu_{AP(n)}: AP(n) \land AP(n) \to AP(n)$$

by applying the maps μ_x on each factor $MGL_{(p)}/(p, x)$.

Lemma 9.3.8. If, in the above setting, $\pi_{k'+1,l'}M = \pi_{2k'+2,2l'}M = \pi_{3k'+3,3l'}M = 0$, then μ_M is homotopy associative.

If, furthermore, $\pi_{k+1,l}N = \pi_{2k+2,2l}N = \pi_{3k+3,3l}N = 0$, then μ_N is also homotopy associative.

Proof. Let $\mu = \mu_M$ and $\nu = \nu_N$. We have to show that

$$\delta = \mu(\mu \wedge 1_M - 1_M \wedge \mu) : M \wedge M \wedge M \to M$$

is zero. Let $\delta' = \delta(\eta' \wedge 1_M \wedge 1_M) : R \wedge M \wedge M \to M, \, \delta'' = \delta'(1_R \wedge \eta' \wedge 1_M) : R \wedge R \wedge M \to M$ and $\delta''' = \delta''(1_R \wedge 1_R \wedge \eta') = \delta \circ (\eta')^{\wedge 3} : R \wedge R \wedge R \to M$. Consider



The top triangle obviously commutes, the right triangle commutes (up to homotopy) by the definition of μ , and the large triangle commutes because η' is an *R*-module map. Thus, $\mu \circ (\eta')^{\wedge 2} = \eta' \circ m$, and it follows

$$\mu(\mu \wedge 1 - 1 \wedge \mu)(\eta')^{\wedge 3} = \mu(\eta' m \wedge \eta' - \eta' \wedge \eta' m)$$
$$= \mu \circ (\eta')^{\wedge 2}(m \wedge 1 - 1 \wedge m) = \eta' m(m \wedge 1 - 1 \wedge m),$$

which vanishes by the associativity of m. Thus, $\delta''' = \delta''(1 \wedge 1 \wedge \eta') = 0$.

This implies that δ'' factors through an *R*-module map $\zeta : S^{k'+1,l'} \wedge R \wedge R \wedge R \to M$, as in the following diagram:

$$R \wedge R \wedge R \xrightarrow{1 \wedge 1 \wedge \eta'} R \wedge R \wedge M \xrightarrow{1 \wedge 1 \wedge \partial'} S^{k'+1,l'} \wedge R \wedge R \wedge R$$

Now, $R \wedge R \wedge R$ is a ring spectrum and ζ can be considered as a map of $R \wedge R \wedge R$ modules. By Lemma 9.3.3 and the assumption on $\pi_{k'+1,l'}M$, ζ must be trivial. Thus, $\delta'' = 0$. Again, this implies $\delta' = \zeta'(1_R \wedge \partial' \wedge 1_M)$ for some $\zeta' : S^{k'+1,l'} \wedge R \wedge R \wedge M \to M$. The $R \wedge R \wedge R$ -module map $\zeta'(1_{S^{k'+1,l'}} \wedge 1_R \wedge 1_R \wedge \eta')$ is a degree (k'+1,l')-map from the ring spectrum $R \wedge R \wedge R$ to M, and therefore vanishes by Lemma 9.3.3. It follows that $\zeta' = \zeta''(1_{S^{k'+1,l'}} \wedge 1_R \wedge 1_R \wedge \partial')$ for some map $\zeta'' : S^{2k'+2,2l'} \wedge R \wedge R \wedge R \to M$.

$$S^{k'+1,l'} \wedge R \wedge R \xrightarrow{1 \wedge \eta'} S^{k'+1,l'} \wedge R \wedge R \wedge M \xrightarrow{1 \wedge \partial'} S^{2k'+2,2l'} \wedge R \wedge R \wedge R$$

By the second assumption on $\pi_{**}M$, $\zeta'' = 0$. It follows that $\delta' = \zeta'(1 \wedge \partial' \wedge 1) = \zeta''(1 \wedge \partial' \wedge \partial') = 0$. That is, $\delta(\eta' \wedge 1_M \wedge 1_M) = \delta' = 0$, which implies $\delta = \zeta'''(\partial' \wedge 1_M \wedge 1_M)$ for some $\zeta''' : S^{k'+1,l'} \wedge R \wedge M \wedge M \to M$.

Now, $\zeta'''(1 \wedge \eta' \wedge 1)(1 \wedge 1 \wedge \eta')$ is a map from $R \wedge R \wedge R$ to M of degree (k'+1,l'), and therefore trivial. Thus, $\zeta'''(1 \wedge \eta' \wedge 1) = \zeta^{(4)}(1 \wedge 1 \wedge \partial')$ with $\zeta^{(4)} : S^{2k'+2,2l'} \wedge R \wedge R \wedge R \to M$, which is also trivial. It follows that $\zeta''' = \zeta^{(5)}(1 \wedge \partial' \wedge 1)$ for some $\zeta^{(5)} : S^{2k'+2,2l'} \wedge R \wedge R \wedge M \to M$, and $\delta = \zeta'''(\partial' \wedge 1 \wedge 1) = \zeta^{(5)}(\partial' \wedge \partial' \wedge 1)$. The map $\zeta^{(5)}(1 \wedge 1 \wedge \eta) : S^{2k'+2,2l'} \wedge R \wedge R \wedge R \to M$ is again zero, which implies $\zeta^{(5)} = \zeta^{(6)}(1 \wedge 1 \wedge \partial')$ for some $\zeta^{(6)} : S^{3k'+3,3l'} \wedge R \wedge R \wedge R \to M$. By the third condition on $\pi_{**}M$, this $\zeta^{(6)}$ is zero. Finally, we have

$$\delta = \zeta^{(6)}(\partial' \wedge \partial' \wedge \partial') = 0.$$

The same line of proof can be used to derive the associativity of μ_N from that of

 μ_M . It only needs to be checked that



is commutative, but this was part of the definitions of $\nu_{M,N}$ and μ_N .

Proposition 9.3.9. If p > 2, then $\mu_{AP(n)}$ is homotopy associative.

Proof. Note that this does not follow immediately from the above lemma, as $\pi_{2k+2,2l}N \neq 0$ for $N = MGL_{(p)}/(p,x)$. However, we can use a trick from [Elm+97, Theorem V.3.1], where the topological analogue of this statement is proven.

Let N = R/(x, y) and $A = R/(J - \{x, y\})$ for some set J of elements in $\pi_{**}R$, and assume that we already know that A is equipped with a (homotopy) associative map $\mu_A : A \wedge A \to A$ as above. The product on $R/J \cong A/(x, y) \cong N \wedge_R A$ is given by

$$(N \wedge_R A) \wedge (N \wedge_R A) \xrightarrow{\tau} (N \wedge N) \wedge_R (A \wedge A) \xrightarrow{1 \wedge_R \mu_A} (N \wedge N) \wedge_R A \xrightarrow{\mu_N \wedge_R 1} N \wedge_R A.$$

Therefore, to prove associativity for $\mu_{N \wedge_R A}$, it suffices to prove that the associativity diagram for μ_N commutes after applying $- \wedge_R 1_A$ to it. Applying $- \wedge_R 1_A$ to all diagrams appearing in the above lemma yields the following result: If $\pi_{i,j}(M \wedge_R A) = 0$ for $(i, j) \in \{(k' + 1, l'), (2k' + 2, 2l'), (3k' + 3, 3l')\}$ and $\pi_{i,j}(N \wedge_R A) = 0$ for $(i, j) \in \{(k + 1, l), (2k + 2, 2l), (3k + 3, 3l)\}$, then $\mu_{N \wedge_R A}$ is associative.

Now, let $R = MGL_{(p)}$, A = ABP, $M = MGL_{(p)}/p$ and $N = MGL_{(p)}/(p, x)$. From [Vez01, Definition 5.3], we know that μ_{ABP} is associative. The assumptions on the homotopy groups are satisfied by Lemma 6.3.7, by which $\pi_{**}(ABP/p) \cong H_{**} \otimes \pi_*(BP/p)$ and $\pi_{**}(ABP/(p, x)) \cong H_{**} \otimes \pi_*(BP/(p, x))$. Note that these homotopy groups vanish in degrees (k + 1, l) and (3k + 3, 3l) for any p because π_*BP is concentrated in even degrees and k is even. However, for $\pi_{(2k+2,2l)}$ to vanish, we need to assume that p is odd.

This proves that $\mu_{ABP/(p,x)}$ is associative. Inductively, we can apply this argument to A = ABP/J' for some $(p) \subset J' \subset J$, where J is as in the proof of Corollary 9.3.5, using $ABP/(J' \cup \{y\}) = MGL_{(p)}/(p,y) \wedge_{MGL_{(p)}} ABP/J'$ (compare Lemma 9.3.7).

Recall that, for $k = \mathbb{C}$ and n > 0, $AP(n)_{**} \cong H_{**} \otimes_{\mathbb{F}_p} P(n)_*$ (Lemma 6.3.7), which is a ring. Hence, we can speak of $AP(n)_{**}$ -modules. Note that $AP(0)_{**}$ is a ring, anyway, since AP(0) = ABP is a ring spectrum.

Lemma 9.3.10. Let $k = \mathbb{C}$. On coefficients, the map

 $\mu_{AP(n)}: AP(n) \wedge AP(n) \to AP(n)$

induces the multiplication on $AP(n)_{**}$ given by $AP(n)_{**} \cong H_{**} \otimes_{\mathbb{F}_n} P(n)_{*}$.

Proof. By [Pel11, Theorem 3.6.16], the motivic Atiyah Hirzebruch spectral sequence in Lemma 6.3.7 is multiplicative, yielding an isomorphism of rings between $AP(n)_{**}$ with multiplication induced from $\mu_{AP(n)}$ and $E_2 = H_{**} \otimes_{\mathbb{F}_p} P(n)_*$ with multiplication induced from the ring structure of H_{**} and from $R_{\mathbb{C}}(\mu_{AP(n)}) : P(n) \wedge P(n) \to P(n)$. It therefore suffices to show that $R_{\mathbb{C}}(\mu_{AP(n)})$ induces the ring structure on $P(n)_*$. Now, $R_{\mathbb{C}}$ carries all the above diagrams to the analogous topological diagrams and, therefore, $BP \wedge P(n) \to P(n)$ (inducing the action of BP_* on $P(n)_*$) factors through $R_{\mathbb{C}}(\mu_{AP(n)})$, which, hence, induces the induced action of $P(n)_*$ on $P(n)_*$.

Corollary 9.3.11. In $SH(\mathbb{C})$, the action $\nu_n : ABP \wedge AP(n) \to AP(n)$ factors through a map $\mu_{AP(n)} : AP(n) \wedge AP(n) \to AP(n)$. If p > 2, the induced action of ABP_{**} on $AP(n)_{**}(X)$ and $AP(n)^{**}(X)$ gives $AP(n)_{**}(X)$ and $AP(n)^{**}(X)$ the structure of left $AP(n)_{**}$ -modules for any $n \ge 0$ and $X \in SH(\mathbb{C})$.

Proof. We have to check that the action of $AP(n)_{**}$ on $AP(n)_{**}(X)$ and $AP(n)^{**}(X)$ is unital and associative. This is equivalent to $\mu_{AP(n)}$ being homotopy left unital and associative. Unitality follows from the maps $\nu_M : R \wedge M \to M$ being unital, since, by definition of μ_M , $\nu_M = \mu_M \circ (\eta' \wedge 1) : R \wedge M \to M \wedge M \to M$. Associativity is proven in Proposition 9.3.9 for p > 2.

9.4 Bousfield classes of AK(n) and AB(n)

Recall $AB(n) = v_n^{-1}AP(n)$.

We want to use methods of [JW75] and [Wür76] to prove that

$$AK(n)_{**}X = 0 \Leftrightarrow AB(n)_{**}X = 0$$

for $X \in S\mathcal{H}(\mathbb{C})$. For this, we need the following two results, which hold in $S\mathcal{H}(k)$ for any $k \subseteq \mathbb{C}$ and are analogous to [Wür76, Formula (2.8)]. The reader may skip the first proposition, as it is a special case of the second one. The proof of the first one is maybe more illustrative.

Let $\mu_n : ABP^{**}(-) \to AP(n)^{**}(-)$ be induced from

$$ABP \xrightarrow{1 \wedge i} ABP \wedge AP(n) \xrightarrow{\nu_n} AP(n),$$

where $i: S \to AP(n)$ is induced from the unit map $S \to ABP$ and ν_n is the structure map of the ABP-module AP(n). For $s^E \in ABP^{**}ABP$ as in Lemma 9.1.2(2), we have $\mu_n(s^E) \in AP(n)^{**}(ABP)$.

Proposition 9.4.1. For any $n \ge 0$, the map

$$h_n: AP(n)^{**}[[\mu_n(s^E)]] \to AP(n)^{**}ABP,$$

given by $h_n(\sum_E x_E \cdot \mu_n(s^E)) = \sum_E \nu_n(s^E \wedge x_E)$, is an isomorphism of ABP^{**} -modules. If $k = \mathbb{C}$ and p > 2, it is an isomorphism of $AP(n)^{**}$ -modules by Corollary 9.3.11.

Proof. We proceed by induction. For AP(0) = ABP, μ_0 is the identity and the claim holds by Lemma 9.1.2(2). Now, assume h_n is an isomorphism for some $n \ge 0$. Consider the following diagram, consisting of two exact sequences induced by $AP(n) \xrightarrow{v_n} AP(n) \rightarrow P(n)$ AP(n+1).

$$AP(n)^{**}ABP \xrightarrow{v_n^*} AP(n)^{**}ABP \xrightarrow{\lambda_n^*} AP(n+1)^{**}ABP \xrightarrow{\delta^*} AP(n)^{**}ABP \xrightarrow{v_n^*} AP(n)^{**}ABP \xrightarrow{\lambda_n^*} AP(n)^{**}AP(n)^{**}ABP \xrightarrow{\lambda_n^*} AP(n)^{**}ABP \xrightarrow{\lambda_n^*} AP(n)^{**}AP \xrightarrow{\lambda_n^*} AP \xrightarrow{\lambda_n^*$$

The lower sequence is the exact sequence

$$\cdots \to AP(n)^{**} \to AP(n)^{**} \to AP(n+1)^{**} \to AP(n)^{**} \to AP(n)^{**} \cdots,$$

tensored over \mathbb{F}_p with $\mathbb{F}_p[[s^E]]$. We show that the diagram commutes. In the first square, the upper composition takes $x \cdot \mu_n(s^E) \in AP(n)^{**}[[\mu_n s^E]]$ to

$$v_n^*(h_n(x\mu_n s^E)) : ABP \xrightarrow{s^E \wedge x} ABP \wedge AP(n) \xrightarrow{\nu_n} AP(n) \xrightarrow{v_n} AP(n),$$

and the lower composition takes the same element to

$$h_n((v_n \cdot x) \cdot \mu_n(s^E)) : ABP \xrightarrow{s^E \wedge x} ABP \wedge AP(n) \xrightarrow{1 \wedge v_n} ABP \wedge AP(n) \xrightarrow{\nu_n} AP(n).$$

Therefore, the commutativity of the first square is equivalent to the commutativity of the square

This square commutes because v_n is a map of ABP-modules (compare Lemma 9.3.1).

In the second square, the upper composition takes $x \cdot \mu_n(s^E) \in AP(n)^{**}[[\mu_n s^E]]$ to

$$(\lambda^* \circ h_n)(x\mu_n s^E) : ABP \xrightarrow{s^E \wedge x} ABP \wedge AP(n) \xrightarrow{\nu_n} AP(n) \xrightarrow{\lambda_n} AP(n+1)$$

and the lower composition takes $x \cdot \mu_n(s^E)$ to

$$h_{n+1}(\lambda_n x \cdot \mu_n s^E) : ABP \xrightarrow{s^E \wedge x} ABP \wedge AP(n) \xrightarrow{1 \wedge \lambda_n} ABP \wedge AP(n+1) \xrightarrow{\nu_{n+1}} AP(n+1).$$

Thus, the commutativity of the second square is equivalent to the commutativity of

which holds because λ_n is, by definition, a map in the category of ABP-modules (see Definition 6.3.1).

In the third square, the upper composition takes $x \cdot \mu_{n+1}(s^E) \in AP(n+1)^{**}[[\mu_{n+1}s^E]]$ to $(\delta \circ h_{n+1})(x\mu_{n+1}s^E) = \delta(\nu_{n+1}(s^E \wedge x))$ and the lower composition takes $x \cdot \mu_{n+1}(s^E)$ to $\nu_n(s^E \wedge \delta(x))$. Thus, the commutativity of the third square follows from the commutativity of

$$\begin{array}{c} ABP \wedge AP(n+1) \xrightarrow{1 \wedge \delta} ABP \wedge AP(n) \\ \downarrow \nu_{n+1} & \qquad \qquad \downarrow \nu_n \\ AP(n+1) \xrightarrow{\delta} AP(n). \end{array}$$

Finally, the five lemma implies that h_{n+1} is an isomorphism.

The above proposition holds more generally:

Proposition 9.4.2. Let $h \in S\mathcal{H}(k)^{cell}$ be any cellular ABP-module spectrum. Then

$$h^{**}ABP \cong h^{**}[[s^E]]$$

as ABP^{**} -modules. In particular, this also holds for h = Ak(n).

Proof. We apply the universal coefficient spectral sequence from [DI05, Proposition 7.7] (see also Proposition 8.1.1) to E = ABP (which is a ring spectrum by [Vez01, Definition 5.3] and is cellular by Remark 6.3.3(1)), $M = ABP \wedge ABP$ and N = h.

$$\operatorname{Ext}_{ABP_{**}}^{***}(ABP_{**}ABP, h_{**}) \Rightarrow \pi_{**}F_{ABP}(ABP \land ABP, h),$$

converging conditionally to $\pi_{**}F_{ABP}(ABP \wedge ABP, h) \cong h^{**}(ABP)$. As $ABP_{**}ABP \cong$ $ABP_{**}{t^E}$ is free over ABP_{**} (Lemma 9.1.2(1)), the higher Ext-groups vanish and the sequence collapses to

$$h^{**}ABP \cong \operatorname{Hom}_{ABP_{**}}(ABP_{**}\{t^E\}, h_{**}),$$

which is isomorphic to $h^{**}[[s^E]]$, as in Lemma 9.1.2(2).

Würgler constructs operations

$$s_Q^E: MUQ^i(-) \to MUQ^{i+|E|}(-)$$

for regular sequences Q [Wür76, Theorem 5.1]. For MUQ = P(n), these operations specify a choice of the operations $(r_E)_n : P(n)^*(-) \to P(n)^{*+|E|}(-)$ considered in [JW75, Section 4]. They are needed in the proof of the isomorphism $B(n)_*(X) \cong$ $K(n)_*(X) \otimes \mathbb{F}_p[v_{n+1}, v_{n+2}, \cdots]$ in [JW75, Proposition 4.14]. We will now state motivic analogues of some of Würgler's lemmas for MUQ = P(n).

In $\mathcal{SH}(k)^{cell}$, let $AP(n)[\mathbf{t}]$ represent the cohomology theory

$$AP(n)^{**}(-)[\mathbf{t}] = ABP^{**}[t_1, t_2, \cdots] \otimes_{ABP^{**}} AP(n)^{**}(-)$$

with $\deg(t_i) = -(2(p^i - 1), p^i - 1).$

Remark 9.4.3. Würgler defines $h^*(-)[\mathbf{t}] = h^*(-) \otimes_{h^*} h^*[t_1, t_2, \cdots]$, where the t_i take any even degree, i.e., $\deg(t_i) = -2i$ in [Wür76, Section 5]. He also considers operations s^E of any degree $\sum_i 2ie_i$. However, Johnson and Wilson [JW75, Sections 1 and 4] are only interested in operations of degree $|E| = \sum_i 2(p^i - 1)e_i$. We will see that it suffices to restrict to those degrees. Most of Würgler's results we refer to in the following are formulated in a much greater generality and do not depend on any degrees of particular elements. The only point where the degrees of the t_i are important is in [Wür76, Theorem 5.1] and we will comment on that below Theorem 9.4.12.

Lemma 9.4.4. For $X \in S\mathcal{H}(k)^{fin}$,

$$(ABP \land AP(n))^{**}(X) \cong AP(n)^{**}(X)[\mathbf{t}].$$

As any cohomology theory on $\mathcal{SH}(k)^{fin}$ extends uniquely to $\mathcal{SH}(k)^{cell}$ [NSØ09, Lemma 4.10], it follows that we can take

$$AP(n)[\mathbf{t}] = ABP \wedge AP(n).$$

Proof. We consider the map

$$(ABP \land ABP)^{**} \otimes_{ABP^{**}} AP(n)^{**}(X) \to (ABP \land AP(n))^{**}(X),$$

induced by the ABP-module structure on AP(n). First, we see that

$$(ABP \land ABP)^{**} \cong (ABP \land ABP)_{-*,-*} \cong ABP_{-*,-*}[\mathbf{t}] \cong ABP^{**}[\mathbf{t}]$$

by Lemma 9.1.2(1). Note that the t_i appearing here are dual to the t_i in Lemma 9.1.2(1), hence $\deg(t_i) = -(2(p^i - 1), p^i - 1)$. To prove the claim, it suffices to show that the above map is an isomorphism.

For n = 0 and $X = S^0$, this is clear. Induction on n shows

$$(ABP \land ABP)^{**} \otimes_{ABP^{**}} AP(n)^{**} \cong (ABP \land AP(n))^{**}$$

for any n, since $AP(n) \to AP(n) \to AP(n+1)$ induces long exact sequences on both sides and the diagram commutes because $AP(n) \to AP(n) \to AP(n+1)$ are maps of ABP-modules. As also any cofiber sequence $X \to Y \to Z$ induces long exact sequences on both sides, cellular induction proves the claim for any finite spectrum X.

We state an analogue of [Wür76, Lemma 3.14].

Lemma 9.4.5. Let $h \in S\mathcal{H}(k)^{cell}$, $k \subseteq \mathbb{C}$, be an ABP-module. The multiplication $ABP \wedge h \rightarrow h$ induces an isomorphism of ABP^{**} -modules,

$$ABP^{**}ABP \otimes_{ABP^{**}} h^{**}AP(n) \cong h^{**}(ABP \wedge AP(n)).$$

Proof. We apply the spectral sequence from [DI05, Proposition 7.7] to E = ABP, $M = ABP \land ABP \land ABP$ and N = h:

$$\operatorname{Ext}_{ABP_{**}}^{***}(ABP_{**}(ABP \land ABP), h_{**}) \Rightarrow \pi_{**}F_{ABP}(ABP \land ABP \land ABP, h),$$

converging conditionally to $h^{**}(ABP \wedge ABP)$. By [NSØ09, Lemma 5.1(i)],

$$ABP_{**}(ABP \land ABP) \cong ABP_{**}ABP \otimes_{ABP_{**}} ABP_{**}ABP,$$

which is free over ABP_{**} . Thus, the spectral sequence collapses and

$$h^{**}(ABP \land ABP) \cong \operatorname{Hom}_{ABP_{**}}(ABP_{**}ABP \otimes_{ABP_{**}} ABP_{**}ABP, h_{**}).$$

Now, $ABP_{**}ABP \otimes_{ABP_{**}} ABP_{**}ABP \cong ABP_{**}\{t_1^E t_2^F\}$, and, therefore,

 $\operatorname{Hom}_{ABP_{**}}(ABP_{**}ABP \otimes_{ABP_{**}} ABP_{**}ABP, h_{**}) \cong h^{**}[[s_1^E s_2^F]]$

$$\cong ABP^{**}[[s_1^E]] \otimes_{ABP_{**}} h^{**}[[s_2^F]] \cong ABP^{**}ABP \otimes_{ABP_{**}} h^{**}ABP,$$

using Proposition 9.4.2. This proves the claim for n = 0.

Assume we have shown that $ABP \wedge h \rightarrow h$ induces an isomorphism

$$ABP^{**}ABP \otimes_{ABP^{**}} h^{**}AP(n) \cong h^{**}(ABP \wedge AP(n))$$

for some $n \ge 0$. The cofiber sequence $AP(n) \to AP(n) \to AP(n+1)$ induces long exact sequences on both sides of this isomorphism, which form a commutative diagram because the maps $AP(n) \to AP(n) \to AP(n+1)$ are maps of ABP-modules. Thus, the five lemma implies the claim for n+1.

Combining the above isomorphism with the one from Proposition 9.4.2, we get:

Lemma 9.4.6. Let $k = \mathbb{C}$ and p > 2. Let X = AP(n) or $X = AP(n)[\mathbf{t}]$. The action of ABP on AP(m), $ABP \land AP(m) \rightarrow AP(m)$, induces an isomorphism of $AP(m)^{**}$ -modules

$$AP(m)^{**}(ABP) \otimes_{AP(m)^{**}} AP(m)^{**}(X) \xrightarrow{\cong} AP(m)^{**}(ABP \wedge X).$$

Proof. From Proposition 9.4.1, we know that

$$AP(m)^{**}(ABP) \cong AP(m)^{**} \otimes_{ABP^{**}} ABP^{**}ABP$$

as $AP(m)^{**}$ -modules. With h = AP(m), the above lemma immediately implies

$$AP(m)^{**}(ABP) \otimes_{AP(m)^{**}} AP(m)^{**}(AP(n)) \xrightarrow{\cong} AP(m)^{**}(ABP \land AP(n))$$

as ABP^{**} -modules and, by Corollary 9.3.11, also as $AP(m)^{**}$ -modules.

For $X = AP(n)[\mathbf{t}] = ABP \wedge AP(n)$, we set $M = ABP^{\wedge 4}$ in the proof of the previous lemma. Using $\pi_{**}ABP^{\wedge 4} \cong ABP_{**}ABP \otimes_{ABP_{**}} \pi_{**}ABP^{\wedge 3}$ [NSØ09, Proposition 5.1(i)], the proof proceeds with exactly the same arguments as the proof of Lemma 9.4.5.

As a consequence of Corollary 9.3.5, we show the following (see [JW75, Lemma 2.8(b)]):

Corollary 9.4.7. Let $k = \mathbb{C}$ and p > 2. The cofibration $S^{2p^n-2,p^n-1} \wedge AP(n) \xrightarrow{v_n} AP(n) \to AP(n+1)$ induces short exact sequences

$$0 \to Ak(j)^{**}(S^{2p^n-1,p^n-1} \land AP(n)) \to Ak(j)^{**}(AP(n+1)) \to Ak(j)^{**}(AP(n)) \to 0$$

for every j > n.

Proof. We have to show that, in the $Ak(j)^{**}$ -long exact sequence, the map induced by v_n is zero. This map is defined as the composition of the two left arrows in the following commutative diagram:

$$Ak(j)^{**}(S^{2p^{n}-2,p^{n}-1} \wedge AP(n)) \xleftarrow{\cong} ABP^{**}(S^{2p^{n}-2,p^{n}-1}) \otimes_{ABP^{**}} Ak(j)^{**}(AP(n))$$

$$Ak(j)^{**}(v_{n} \wedge 1) \uparrow \qquad ABP^{**}(v_{n}) \otimes 1 \uparrow$$

$$Ak(j)^{**}(ABP \wedge AP(n)) \xleftarrow{\cong} ABP^{**}(ABP) \otimes_{ABP^{**}} Ak(j)^{**}(AP(n))$$

$$Ak(j)^{**}(v_{n}) \uparrow$$

$$Ak(j)^{**}(AP(n))$$

The horizontal maps are induced by the ABP-module structure on Ak(j). The lower horizontal map is an isomorphism by the previous lemma. The upper isomorphism is proven similarly, setting $M = ABP \wedge ABP$ in the above proof. We show that the right hand map is zero. Let $\sum s^E \otimes x_E$ be an element of $ABP^{**}(ABP) \otimes_{ABP^{**}} Ak(j)^{**}(AP(n))$. It maps to $\sum s^E(v_n) \otimes x_E$. In Corollary 9.1.7, we set k = 1, s = 0and m = n, to see that for all $|E| \geq 0$, either $s^E(v_n) \equiv v_n \mod I_n$ or $s^E(v_n) \equiv 0$ mod I_n . Thus, $s^E(v_n) \in I_{n+1} = (v_0, \dots, v_n)$. By Corollary 9.3.5, I_{n+1} acts trivially on $Ak(j)^{**}(AP(n))$ for j > n, hence, the right hand map is zero.

It follows that the left map factors through zero, proving the claim.

The isomorphism from Lemma 9.4.6 is now used to define an $AP(m)^{**}(ABP)$ comodule structure on $AP(m)^{**}(AP(n))$ by

$$AP(m)^{**}AP(n) \xrightarrow{\nu_n^*} AP(m)^{**}(ABP \land AP(n)) \stackrel{\cong}{\leftarrow} AP(m)^{**}ABP \otimes_{AP(m)^{**}} AP(m)^{**}AP(n),$$

where ν_n is the *ABP*-module structure map on *AP*(*n*). Note that, for $k \neq \mathbb{C}$ or p = 2, we do not know if these groups are *AP*(*m*)**-modules (see Corollary 9.3.11). In this case, we might only get an *ABP****ABP*-comodule structure on *AP*(*m*)***AP*(*n*).

Lemma 9.4.8. Let $k = \mathbb{C}$, $m > n \ge 0$ and E be an exponent sequence. Let $s^E : ABP \rightarrow ABP$ be as in Lemma 9.1.2 and $\mu_m : ABP \rightarrow AP(m)$ be as defined in the beginning of this section. Then

$$\mu_m \circ s^E \circ v_n : S \to ABP \to ABP \to AP(m)$$

is the zero map.

Proof. The realisation functor $R_{\mathbb{C}}$ takes the composition $\mu_m \circ s^E \circ v_n \in AP(m)_{**}$ to $\mu_m^{\text{top}} \circ s_{\text{top}}^E \circ v_n^{\text{top}} \in P(m)_*$. By the *BP*-version of [Wür76, Lemma 2.2], μ_m^{top} : $BP \to P(m)$ is the canonical projection. By the invariant prime ideal theorem, $I_{n+1}^{\text{top}} = (v_0^{\text{top}}, v_1^{\text{top}}, \cdots, v_n^{\text{top}}) \subset BP_*$ is invariant under the action of $s_{\text{top}}^E \in BP^*BP$, hence, $s_{\text{top}}^E(v_n^{\text{top}}) \in I_{n+1}^{\text{top}}$. Thus, $\mu_m^{\text{top}}(s_{\text{top}}^E \circ v_n^{\text{top}}) \in P(m)_*$ lies in the image of I_{n+1} under the projection $BP_* \to P(m)_* = BP_*/(v_0^{\text{top}}, \cdots, v_{m-1}^{\text{top}})$, which is zero, since m > n. This implies $R_{\mathbb{C}}(\mu_m \circ s^E \circ v_n) = 0 \in P(m)_*$.

For $k = \mathbb{C}$, $AP(m)_{**} \cong P(m)_*[\tau]$ by Lemma 6.3.7, and any homogeneous element in $AP(m)_{**}$ that realises to zero in $P(m)_*$ already has to be zero in $AP(m)_{**}$. Hence, $\mu_m \circ s^E \circ v_n = 0$ in $AP(m)_{**}$.

Now we present an analogue of a special case of [Wür76, Proposition 4.12].

Lemma 9.4.9. Let $k = \mathbb{C}$, p > 2 and $m \ge n$. As $AP(m)^{**}ABP$ -comodules,

$$AP(m)^{**}AP(n) \cong AP(m)^{**}ABP \otimes_{AP(n)^{**}} \Lambda_{AP(n)^{**}}[[\beta_0, \cdots, \beta_{n-1}]]$$

with $\deg(\beta_i) = (2p^i - 1, p^i - 1).$

Proof. For n = 0, the statement is trivial. Assume the proposition holds for some pair (m, n) with m > n. We show that it also holds for (m, n + 1).

Consider $AP(n) \xrightarrow{v_n} AP(n)$. We show that the induced map

$$v_n^* : AP(m)^{**}AP(n) \to AP(m)^{**}AP(n)$$

is trivial. This works similar to [Wür76, Lemma 3.15]. Let $\phi : S \to ABP$ represent $v_n \in ABP_{**}$ and let $\phi^* : AP(m)^{**}ABP \to AP(m)^{**}$ be the map induced on $AP(m)^{**}(-)$. For $x \in AP(m)^{**}ABP = AP(m)^{**}[[\mu_m s^E]]$ (see Proposition 9.4.1), we have $x = \sum_E \lambda_E \mu_m s^E$ and

$$\phi^*(x) = \phi^*(\sum_E \lambda_E \mu_m s^E) = \sum_E \lambda_E \mu_m(s^E(v_n)),$$

because ϕ^* is precomposition with $v_n \in ABP_{**}$. By Lemma 9.4.8, $\mu_m(s^E(v_n)) = 0$ in $AP(m)^{**}$, hence, $\phi^* = 0$. Now consider the following commutative square:

$$\begin{array}{ccc} AP(m)^{**}(ABP \wedge AP(n)) & \xrightarrow{(\phi \wedge 1)^{*}} & AP(m)^{**}(AP(n)) \\ & \cong & & & \uparrow \\ & & & \uparrow \\ AP(m)^{**}ABP \otimes_{AP(m)^{**}} AP(m)^{**}AP(n)_{\phi^{*} \otimes 1} & AP(m)^{**} \otimes_{AP(m)^{**}} AP(m)^{**}AP(n) \end{array}$$

The left map is an isomorphism by Lemma 9.4.6. Since $\phi^* = 0$, it follows that $(\phi \wedge 1)^* = 0$. Since v_n^* is, by definition, the precomposition of $(\phi \wedge 1)^*$ with the map $AP(m)^{**}AP(n) \rightarrow AP(m)^{**}(ABP \wedge AP(n)), v_n^* = 0$, as claimed.

It follows that the long exact $AP(m)^{**}(-)$ -sequence induced by

$$S^{2p^n-2,p^n-1} \wedge AP(n) \xrightarrow{v_n} AP(n) \to AP(n+1)$$

splits into short exact sequences of $AP(m)^{**}ABP$ -comodules

$$0 \to AP(m)^{**}(S^{2p^n-1,p^n-1} \land AP(n)) \to AP(m)^{**}AP(n+1) \to AP(m)^{**}AP(n) \to 0.$$

Analogously to [Wür76, Proposition 4.12], it follows inductively that

$$AP(m)^{**}AP(n+1) \cong AP(m)^{**}AP(n) \otimes_{AP(m)^{**}} \Lambda_{AP(m)^{**}}(\beta_n)$$

and the degree of β_n is determined by the degrees appearing in the exact sequence. \Box

If M is an $AP(n)^{**}(ABP)[\mathbf{t}]$ -comodule with structure map ψ , an element $a \in M$ is called primitive if $\psi(a) = 1 \otimes a$. Similarly to [Wür76, Lemma 4.13], the following holds:

Lemma 9.4.10. Let $k = \mathbb{C}$ and p > 2. Let $g : AP(n) \to AP(n)[\mathbf{t}]$ be a map of spectra. Then g is a map of ABP-module spectra if and only if it is a primitive element of the $AP(n)^{**}(ABP)[\mathbf{t}]$ -comodule $AP(n)^{**}(AP(n))[\mathbf{t}]$.

Proof. This follows from Lemma 9.4.6 in the same manner as [Wür76, Lemma 4.13] follows from [Wür76, Lemma 3.14] for X = P(n) and $X = P(n)[\mathbf{t}]$.

Now we state a result which is directly used for the construction of the operation we are aiming at. This corresponds to [Wür76, Theorem 4.17].

Proposition 9.4.11. Let $k = \mathbb{C}$ and p > 2. There is a degree-preserving group isomorphism

$$\operatorname{Hom}_{ABP}^{**}(AP(n), AP(n)[\mathbf{t}]) \cong \Lambda_{AP(n)^{**}[\mathbf{t}]}[[\beta_0, \beta_1, \cdots]],$$

where the left hand side is the bigraded abelian group of maps of ABP-module spectra and the right hand side is an exterior algebra over $AP(n)^{**}[t]$.

Proof. Würgler derives this from [Wür76, Proposition 4.12] and [Wür76, Lemma 4.13] using that inverse limits of primitive elements are primitive [Wür76, Lemma 4.16]. The same line of proof proves this proposition using Proposition 9.4.9 and Lemma 9.4.10. Denoting the set of primitive elements in M by $\Pr\{M\}$, the proof can be summarised by the following sequence of isomorphisms:

$$\operatorname{Hom}_{ABP}^{**}(AP(n), AP(n)[\mathbf{t}]) \cong \operatorname{Pr} \left\{ AP(n)^{**}(AP(n))[\mathbf{t}] \right\}$$
$$\cong \operatorname{Pr} \left\{ AP(n)^{**}(ABP)[\mathbf{t}] \otimes_{AP(n)^{**}[\mathbf{t}]} \Lambda_{AP(n)^{**}[\mathbf{t}]}[[\beta_0, \beta_1, \cdots]] \right\}$$
$$\cong \operatorname{Pr} \left\{ AP(n)^{**}(ABP)[\mathbf{t}] \right\} \otimes_{AP(n)^{**}[\mathbf{t}]} \Lambda_{AP(n)^{**}[\mathbf{t}]}[[\beta_0, \beta_1, \cdots]]$$
$$\cong AP(n)^{**}[\mathbf{t}] \otimes_{AP(n)^{**}[\mathbf{t}]} \Lambda_{AP(n)^{**}[\mathbf{t}]}[[\beta_0, \beta_1, \cdots]] \cong \Lambda_{AP(n)^{**}[\mathbf{t}]}[[\beta_0, \beta_1, \cdots]].$$

This completes the preparation Würgler needs for [Wür76, Theorem 5.1]. We state our version of this theorem, now using the notation from [JW75].

Theorem 9.4.12. Let p > 2. In $\mathcal{SH}(\mathbb{C})^{cell}$, there exists a family $(r_E)_n$, $n \ge 0$, of natural stable operations

$$(r_E)_n : AP(n)^{**}(-) \to AP(n)^{**}(-)$$

of degree (|E|, |E|/2), such that

$$(r_E)_n(ux) = \sum_{F+G=E} s_F(u)(r_G)_n(x)$$

for $u \in ABP^{**}(X)$ and $x \in AP(n)^{**}(X)$ (see Definition 9.1.3 for the definition of s_F) and such that

commutes.

Proof. Let $s_t : ABP^{**}(-) \to ABP^{**}(-)[\mathbf{t}]$ be given by $s_t(x) = \sum_E s_E(x) \otimes t^E$. As in [Wür76, Theorem 5.1], the square

can be completed with the help of Proposition 9.4.11 and we define $(r_E)_n$ by $s_{t,n}(x) = \sum_E (r_E)_n(x) \otimes t^E$.

Since Proposition 9.4.11 gives us a map of *ABP*-module spectra, the operation $s_{t,n}$ satisfies $s_{t,n}(ux) = s_t(u)s_{t,n}(x)$. It follows that

$$\sum_{E} (r_E)_n(ux) \otimes t^E = \sum_{F+G=E} s_F(u)(r_G)_n(x) \otimes t^F t^G$$

and, hence, $(r_E)_n(ux) = \sum_{F+G=E} s_F(u)(r_G)_n(x)$, which proves the first property claimed. The second property holds because the above commutative square has to commute on the level of each t^E .

Remark 9.4.13. Originally (see e.g. [Qui71, Formula (2.4)]), one first defines the operation $s_t : BP^*(-) \to BP^*(-)[t_1, t_2, \cdots]$, $\deg(t_i) = -2i$, and then uses it to construct operations s_E of degree $\sum_i 2ie_i$. The s_t used in the above proof contains only those summands $s_E \otimes t^E$ with $|E| = \sum_i 2(p^i - 1)$, because we work with ABP-modules instead of MGL-modules. Hence, the $s_{t,n}$ also consists of less summands than the corresponding operation in [Wür76, Theorem 5.1], but this suffices to define exactly those $(r_E)_n$ with the degrees needed in [JW75, Section 4].

In [JW75, Proposition 4.14], Johnson and Wilson show that, for any $X \in S\mathcal{H}^{fin}$, there is a natural isomorphism

$$B(n)_*(X) \cong K(n)_*(X) \otimes \mathbb{F}_p[v_{n+1}, v_{n+2}, \cdots]$$

for n < 2p - 2. Johnson and Wilson needed the condition on n because they were only able to canonically define the operations $(r_E)_n$ in this range. As stated in [Wür76, Remark 6.19], the condition becomes redundant if Würgler's operations are used.

With the above operations at hand, we can proceed exactly as in [JW75, Section 4]:

Definition 9.4.14. Let *n* be fixed and let \mathcal{E} be the set of all exponent sequences of the form $E = (0, \dots, 0, e_{n+1}, e_{n+2}, \dots)$. Recall the definition of |E| from Definition 9.1.6. For $E \in \mathcal{E}$, let $q = |E| = 2(p^n - 1)b + a$ with $0 \leq a < 2(p^n - 1)$. With $\sigma^n E = (p^n e_{n+1}, p^n e_{n+2}, \dots)$ and $c = b - (e_{n+1} + e_{n+2} + \dots)$, it follows $|\sigma^n E| = c2(p^n - 1) + a$, exactly as in [JW75]. Note that *q* and, hence, *a* are even. We define $\bar{s}_E \in Ak(n)^{a,a/2}(AP(n))$ by

$$\bar{s}_E: AP(n) \xrightarrow{(r_{\sigma^n E})_n} \Sigma^{c2(p^n-1)+a,c(p^n-1)+a/2} AP(n) \xrightarrow{v_n^c} \Sigma^{a,a/2} AP(n) \xrightarrow{\lambda_n} \Sigma^{a,a/2} Ak(n),$$

where λ_n is the quotient map from AP(n) to $Ak(n) = AP(n)/(v_{n+1}, \cdots)$.

Furthermore, we assume that the set $\{E \in \mathcal{E} \mid |E| = q\}$ is ordered as $\{E_1, \dots, E_v\}$, where v is the \mathbb{F}_p -dimension of $(\mathbb{F}_p[v_{n+1}, v_{n+2}, \dots])_q$, and we denote \bar{s}_{E_u} by s_u .

Lemma 9.4.15. Let $k = \mathbb{C}$, p > 2 and $\bar{s}_E : AP(n)_{**} \to Ak(n)_{**}$ be induced by the above map. Assume n > 0 and let $\tau \in AP(n)_{**}$ be as in Lemma 6.3.7. Then $\bar{s}_E(\tau) = 0$ for $E \neq 0$ and $\bar{s}_0(\tau) = \tau$. If, furthermore, $x \in AP(n)_{**}X$ with $X \in S\mathcal{H}(\mathbb{C})^{cell}$, then $\bar{s}_E(x\tau) = \bar{s}_E(x)\tau$.

Proof. Since $\bar{s}_E(\tau) = (\lambda_n \circ v_n^c \circ (r_{\sigma^n E})_n)(\tau)$ with $\lambda_n(\tau) = \tau$, and c = 0 for E = 0, the first claim is equivalent to $(r_F)_n(\tau) = 0$ for $F \neq 0$ and $(r_0)_n(\tau) = \tau$. The realisation functor $R_{\mathbb{C}}$ maps τ to $1 \in P(n)^0$ and it takes $(r_F)_n : AP(n)^{**} \to AP(n)^{**}$ to $(r_F^{\mathrm{top}})_n : P(n)^* \to P(n)^*$, which, by [Wür76, Theorem 5.1], is compatible with s_F^{top} in the sense that the following diagram commutes:

$$\begin{array}{c} BP^* \xrightarrow{s_F^{\mathrm{top}}} BP^* \\ \downarrow \\ P(n)^* \xrightarrow{(r_F^{\mathrm{top}})_n} P(n)^*. \end{array}$$

The map s_F^{top} is defined via the coaction map

$$\psi^{\mathrm{top}}: BP_* \stackrel{(1 \wedge i)_*}{\longrightarrow} BP_*BP \stackrel{m_*^{-1}}{\longrightarrow} BP_*BP \otimes_{BP_*} BP_*,$$

which clearly takes 1 to $1 \otimes 1$. Therefore, in the formula

$$\psi^{\text{top}}(1) = \sum_{F} c(t_{\text{top}}^{F}) \otimes s_{F}^{\text{top}}(1),$$

all $s_F^{\text{top}}(1)$ have to be 0, except for $s_0^{\text{top}}(1)$, which is 1. It follows that also $(r_F^{\text{top}})_n(1) = 0$ for $F \neq 0$ and $(r_0^{\text{top}})_n(1) = 1$. Hence, $R_{\mathbb{C}}((r_F)_n(\tau))$ is 0 for $F \neq 0$ and it is 1 for F = 0. The projection of $\pi \in P(n)^*$ under $P_{\pi} \in AP(n)^{**} \to P(n)^*$ is

The preimage of $x \in P(n)^*$ under $R_{\mathbb{C}} : AP(n)^{**} \to P(n)^*$ is

$$\left\{\sum_{k\geq 0} x_k \cdot \tau^k \mid \sum_{k\geq 0} x_k = x\right\}.$$

Since $(r_F)_n$ is a map of degree (|F|, |F|/2), the only possible preimage of $R_{\mathbb{C}}((r_F)_n(\tau))$ is $(r_F)_n(\tau) = 0$ for $F \neq 0$ and $(r_0)_n(\tau) = \tau$.

The claim on $\bar{s}_E(x\tau)$ now follows from the Cartan formula, Lemma 9.1.4.

The following is a motivic analogue to [JW75, Proposition 4.14].

Theorem 9.4.16. For any $X \in S\mathcal{H}(\mathbb{C})^{fin}$ and any n > 0, p > 2, there is a natural isomorphism

$$AB(n)_{**}(X) \to AK(n)_{**}(X) \otimes \mathbb{F}_p[v_{n+1}, v_{n+2}, \cdots].$$

Furthermore, for any $X \in \mathcal{SH}(\mathbb{C})$, $AK(n)_{**}(X) = 0$ if and only if $AB(n)_{**}(X) = 0$.

Proof. For a given exponent sequence E, consider the composition

$$AP(n) \xrightarrow{\overline{s}_E} \Sigma^{a,a/2} Ak(n) \to \Sigma^{a,a/2} AK(n) \xrightarrow{v_n^{-b}} \Sigma^{q,q/2} AK(n),$$

where $a = a_E$ and $b = b_E$ are as in Definition 9.4.14. These induce a natural homomorphism

$$\hat{\Lambda} : AP(n)_{**}(X) \to AK(n)_{**}(X) \otimes \mathbb{F}_p[v_{n+1}, v_{n+2}, \cdots]$$

by
$$\hat{\Lambda}(y) = \sum_{E \in \mathcal{E}} v_n^{-b_E} \bar{s}_E(y) \otimes v^E.$$

By Corollary 9.1.7 and the Cartan formula, $r_E(yv_n) \equiv r_E(y)v_n \mod I_n$. As $I_n = (v_0, \dots, v_{n-1})$ acts trivially on AP(n) and Ak(n) (Proposition 9.3.5), it follows that $\bar{s}_E : AP(n)_{**}(X) \to Ak(n)_{**}(X)$ is an $\mathbb{F}_p[v_n]$ -homomorphism. Therefore, $\hat{\Lambda}$ can be extended to $AB(n)_{**}(X) = v_n^{-1}AP(n)_{**}X$. This yields a map

$$\Lambda: AB(n)_{**}(X) \to AK(n)_{**}(X) \otimes \mathbb{F}_p[v_{n+1}, v_{n+2}, \cdots].$$

We show that Λ is an isomorphism for $X = S^0$, and, thus, for all $X = S^{p,q}$. By Lemma 6.3.7, we are considering a map $B(n)_*[\tau] \to K(n)_*[\tau] \otimes \mathbb{F}_p[v_{n+1}, v_{n+2}, \cdots]$ which, under $R_{\mathbb{C}}$, realises to the isomorphism $B(n)_* \xrightarrow{\cong} K(n)_* \otimes \mathbb{F}_p[v_{n+1}, v_{n+2}, \cdots]$ in [JW75, Proposition 4.14]. Thus, it suffices to calculate $\Lambda(\tau)$, or, equivalently, $\hat{\Lambda}(\tau) = \sum_{E \in \mathcal{E}} v_n^{-b_E} \bar{s}_E(\tau) \otimes v^E$. By the previous lemma, this is equal to $v_n^{-b_0} \bar{s}_0(\tau) \otimes v^0 = \tau$. Thus, Λ is an isomorphism if X is a sphere.

Cellular induction via the five lemma shows that Λ is an isomorphism for all $X \in S\mathcal{H}(\mathbb{C})^{fin}$.

By Definition 5.2.1(3), any cell spectrum is the colimit of a diagram of finite cell spectra. As in [Rav84, Theorem 2.1(a)], it follows that, for any $X \in S\mathcal{H}(\mathbb{C})^{cell}$,

 $AB(n)_{**}X = 0$ if and only if $AK(n)_{**}X = 0$. Here, we use that $\pi_{**}(-)$ commutes with filtered colimits by [DI05, Proposition 9.3].

Furthermore, by [DI05, Proposition 7.3], any $X \in S\mathcal{H}(\mathbb{C})$ has a cellular approximation $X' \in S\mathcal{H}(\mathbb{C})^{cell}$, $f: X' \to X^{fib}$ (X^{fib} a fibrant replacement of X), such that $\pi_{**}(f): \pi_{**}X' \xrightarrow{\cong} \pi_{**}X$. Hence, also $\pi_{**}(1_E \wedge f): E_{**}X' \to E_{**}X$ is an isomorphism for any $E \in S\mathcal{H}(\mathbb{C})$. In particular,

$$AB(n)_{**}X = 0 \Leftrightarrow AB(n)_{**}X' = 0 \Leftrightarrow AK(n)_{**}X' = 0 \Leftrightarrow AK(n)_{**}X = 0.$$

Remark 9.4.17. In addition to the above theorem, I also worked on a motivic version of [JW75, Theorem 4.8], which I was almost able to prove, except for one detail which is necessary for part (c), where [JW75, Lemma 4.10] enters the argument. Large parts of the proof of this lemma can be translated motivically, using motivic cellular approximation as introduced in [DI05, Proposition 7.3] and the motivic Hurewicz theorem from [Mor06, Theorem 5.57]. The only problematic point is part (e) in the proof of [JW75, Lemma 4.10], where a spectrum Y^k with finitely generated homology is approximated by a finite spectrum F. A possible construction of $F \to Y^k$ is described in [Hat01, Proposition 4C.1]. A similar construction can be performed in the motivic setting. However, it is not clear that the resulting spectrum F is finite. This would be the case if $(H\mathbb{Z}_{(P)})^{**}$ is a coherent (or Noetherian) ring, because then, $H_{**}(X, \mathbb{Z}_{(p)})$ is finitely generated over $(H\mathbb{Z}_{(P)})^{**}$ for finite cell spectra X, making a finite induction step possible.

Corollary 9.4.18. For p > 2 and n > 0,

$$\langle AK(n) \rangle = \langle AB(n) \rangle$$
 in $\mathcal{SH}(\mathbb{C})$.

This also holds for n = 0, in which case we do not need to assume $k = \mathbb{C}$ or p > 2.

Proposition 9.4.19. For any $k \subseteq \mathbb{C}$ and any prime p,

$$\langle AK(0) \rangle = \langle AB(0) \rangle$$
 in $\mathcal{SH}(k)$.

Proof. By definition, $AB(0) = p^{-1}ABP$ and

$$AK(0) = p^{-1}ABP/(v_1, v_2, \cdots) = p^{-1}MGL_{(p)}/(a_1, a_2, \cdots) = MGL_{\mathbb{Q}}/(a_1, a_2, \cdots).$$

By the main result in [Hoy13], $MGL/(a_1, a_2, \dots) \cong H\mathbb{Z}$, which implies

$$p^{-1}MGL_{(p)}/(a_1, a_2, \cdots) \cong p^{-1}H\mathbb{Z}_{(p)} = H\mathbb{Q}.$$

Hence, $AK(0) \cong H\mathbb{Q}$. From Lemma 9.2.1(1), we already know

 $\langle AK(0) \rangle \le \langle AB(0) \rangle.$

To prove

 $\langle AK(0) \rangle \ge \langle AB(0) \rangle,$

let $X \in S\mathcal{H}(k)^{fin}$ satisfy $AK(0)_{**}(X) = 0$. We have to show $AB(0)_{**}(X) = 0$. We use [Hoy13, Lemma 7.10], in which we rationalise $H\mathbb{Z}$ and MGL and set $X = MGL_{\mathbb{Q}}$. Here, rationalising a spectrum E means forming the homotopy colimit $E_{\mathbb{Q}} = p^{-1}E_{(p)}$. The lemma then states the following.

If $F \in \mathcal{SH}(k)$ satisfies $H\mathbb{Q} \wedge F = 0$, then $[F, MGL_{\mathbb{Q}}] = 0$.

When rationalising the proof of [Hoy13, Lemma 7.10], one uses that $\kappa_0(MGL_{\mathbb{Q}}) \cong (MGL_{\mathbb{Q}})_{\leq 0}$ and $\kappa_0(H\mathbb{Q}) \cong H\mathbb{Q}_{\leq 0}$, which hold because $MGL_{\mathbb{Q}}$ and $H\mathbb{Q}$ are connective (which follows from the connectivity of MGL [Hoy13, Corollary 3.9] and of $H\mathbb{Z}$ [Hoy13, Lemma 7.3]). Furthermore, one uses that

$$(MGL_{\mathbb{Q}})_{\leq 0} \cong (MGL_{\leq 0})_{\mathbb{Q}} \cong (H\mathbb{Z}_{\leq 0})_{\mathbb{Q}} \cong H\mathbb{Q}_{\leq 0}$$

where the first and third isomorphisms follow from the fact that $(-)_{\leq d}$ preserves filtered homotopy colimits [Hoy13, Lemma 2.1] and the middle isomorphism is by [Hoy13, Lemma 7.5].

We apply this version of [Hoy13, Lemma 7.10] to see that $(MGL_{\mathbb{Q}})^{**}X = 0$ for X as above, which is equivalent to $(MGL_{\mathbb{Q}})_{**}X = 0$, since X is finite (Proposition 6.2.2). Using Lemma 9.2.1(1) again, it follows that $(p^{-1}ABP)_{**}(X) = 0$, as we wanted to show.

As in the previous proof, the equivalence

$$AK(0)_{**}X = 0 \Leftrightarrow AB(0)_{**}X = 0$$

passes from finite spectra to cellular spectra and then to arbitrary spectra $X \in \mathcal{SH}(k)$.

As a corollary of the above results, we can prove the following analogue of [Rav84, Theorem 2.1(i)].

Corollary 9.4.20. In $SH(\mathbb{C})$,

$$\langle AK(n) \rangle \wedge \langle AK(m) \rangle = 0$$

for any $m \neq n$.

Proof. Assume m > n. By (1) and (2) of Lemma 9.2.2,

$$\langle AK(n) \rangle \land \langle AP(n+1) \rangle \le \langle AE(n) \rangle \land \langle AP(n+1) \rangle = \langle 0 \rangle.$$

Furthermore, by Lemma 9.2.2(3) and the above result,

$$\langle AP(m) \rangle = \langle AB(m) \rangle \lor \langle AP(m+1) \rangle = \langle AK(m) \rangle \lor \langle AP(m+1) \rangle.$$

This implies $\langle AK(m) \rangle \leq \langle AP(m) \rangle$. Since m > n, $\langle AP(m) \rangle \leq \langle AP(n+1) \rangle$ by Lemma 9.2.1(1). Hence,

$$\langle AK(n) \rangle \wedge \langle AK(m) \rangle \leq \langle AK(n) \rangle \wedge \langle AP(n+1) \rangle = \langle 0 \rangle.$$

9.5 Decomposition of $\langle AE(n) \rangle$

Recall from Definition 6.3.1 that

$$AE(n) = v_n^{-1}ABP/(v_{n+1}, v_{n+2}, \cdots).$$

With the above preparations, we are ready to prove an analogue of the decomposition of Bousfield classes given in [Rav84, Theorem 2.1(d)]. This answers a special case of [Hor06, Question 2.17].

Theorem 9.5.1. For p > 2,

$$\langle AE(n) \rangle = \bigvee_{0 \le i \le n} \langle AK(i) \rangle \text{ in } \mathcal{SH}(\mathbb{C}).$$

Proof. These are the same arguments as for [Rav84, Theorem 2.1(d)]: By Lemma 9.2.2(3) and Corollary 9.4.18, $\langle AP(n) \rangle = \langle AB(n) \rangle \lor \langle AP(n+1) \rangle = \langle AK(n) \rangle \lor \langle AP(n+1) \rangle$. Since AP(0) = ABP, it follows inductively:

$$\langle ABP \rangle = \langle AK(0) \rangle \lor \langle AK(1) \rangle \lor \cdots \lor \langle AK(n) \rangle \lor \langle AP(n+1) \rangle.$$

Since AE(n) is an ABP-module spectrum, $\langle AE(n) \rangle \leq \langle ABP \rangle$ by Lemma 9.2.1(2). By Lemma 9.2.2(2), $\langle AE(n) \rangle \wedge \langle AP(n+1) \rangle = \langle 0 \rangle$. It follows that

$$\langle AE(n) \rangle \leq \langle AK(0) \rangle \lor \langle AK(1) \rangle \lor \cdots \langle AK(n) \rangle.$$

By Corollary 9.1.8 and Lemma 9.2.2(1), $\langle AE(n) \rangle \ge \langle AE(i) \rangle \ge \langle AK(i) \rangle$ for $i \le n$ and hence also $\langle AE(n) \rangle \ge \bigvee_{i \le n} \langle AK(i) \rangle$.

Remark 9.5.2. The restriction p > 2 originates from Section 9.3, where it was needed to prove homotopy associativity for the map $\mu_{AP(n)} : AP(n) \wedge AP(n) \rightarrow AP(n)$. There might be a different way to show that the ABP_{**} -action on $AP(n)_{**}(X)$ induces an $AP(n)_{**}$ -action, in which case the condition p > 2 could be removed in the previous section and in the theorem.

9.6 AK(n) and AK(n+1)

Lemma 9.6.1. Let p be any fixed prime. For $n \ge 1$ and $X \in S\mathcal{H}(\mathbb{C})^{fin}$, $AP(n)_{**}(X)$ is an $AP(1)_{**}AP(1)$ -comodule and a coherent module over $AP(1)_{**}$.

Proof. Note that AP(1) = ABP/p is a ring spectrum because $p : ABP \to ABP$ is a map of ring spectra and the category of ring spectra is cocomplete. The coaction on $AP(n)_{**}(X)$ is defined via

$$AP(n)_{**}X \to AP(n)_{**}(AP(1) \land X) \leftarrow AP(1)_{**}AP(1) \otimes_{AP(1)_{**}} AP(n)_{**}X$$

where the left map is induced by the unit of AP(1) and we need to show that the right map (induced by $AP(1) \wedge AP(n) \rightarrow AP(n)$, compare Corollary 9.3.11) is an isomorphism. To prove this isomorphism, it suffices to show that, in the spectral sequence from [DI05, Proposition 7.7],

$$\operatorname{Tor}^{AP(1)_{**}}(AP(1)_{**}AP(1), AP(n)_{**}X) \Rightarrow AP(n)_{**}(AP(1) \land X),$$

 $AP(1)_{**}AP(1)$ is free over $AP(1)_{**}$, so that the spectral sequence collapses immediately. We want to apply Lemma 6.3.7 to $AP(1) \wedge AP(1)$. Recall that the slice spectral sequence considered in Lemma 6.3.7 converges for quotients of Landweber exact spectra by [Hoy13, Theorem 8.12 and Example 8.13]. Now, $ABP \wedge ABP$ is a product of Landweber exact spectra and is therefore Landweber exact, see e.g. [NSØ09, Remark 9.2], and $AP(1) \wedge AP(1)$ is a quotient of $ABP \wedge ABP$. Hence, the slice spectral sequence converges strongly, and it collapses for the same degree reasons as in Lemma 6.3.7. Thus, $AP(1)_{**}AP(1) \cong P(1)_*P(1)[\tau]$. By [JY80, Section 1], this is isomorphic to $P(1)_*[\tau, z^{E,A}]$ for certain $z^{E,A}$. In particular, it is free over $AP(1)_{**} \cong P(1)_*[\tau]$, as we wanted to show.

Now we show that $AP(n)_{**}(X)$ is coherent over $AP(1)_{**}$. Recall that $P(1)_* \cong \mathbb{F}_p[v_1^{\text{top}}, v_2^{\text{top}}, \cdots]$ is a coherent ring (see [CS69, Section 1]). The same holds for $AP(1)_{**} \cong \mathbb{F}_p[\tau, v_1, v_1, \cdots]$. By [CS69, Proposition 1.2], coherence of modules satisfies the two out of three property for exact triangles of graded modules (i.e., long exact sequences). It follows that $AP(n)_{**}$ is a coherent $AP(1)_{**}$ -module, and cellular induction implies that $AP(n)_{**}(X)$ is a coherent $AP(1)_{**}$ -module, too.

Setting n = 1 and $X = S^0$, the above lemma tells us that $AP(1)_{**}$ is a coherent $AP(1)_{**}AP(1)$ -comodule, where coherent means coherent as an $AP(1)_{**}$ -module.

Lemma 9.6.2. (Invariant prime ideals)

For $k = \mathbb{C}$ and p any prime, the invariant prime ideals of $AP(1)_{**}$ (that is, prime ideals which are also sub-comodules) are given by $I_m = (v_1, \dots, v_{m-1})$ and $\overline{I}_m = (\tau, v_1, \dots, v_{m-1})$.

Proof. We have $AP(1)_{**} \cong P(1)_*[\tau]$ and $AP(1)_{**}AP(1) \cong P(1)_*P(1)[\tau]$, as in the proof of the previous lemma. Under the functor $R_{\mathbb{C}}$, the coaction

$$AP(1)_{**} \to AP(1)_{**}AP(1) \otimes_{AP(1)_{**}} AP(1)_{**}$$

realises to

$$P(1)_* \to P(1)_* P(1) \otimes_{P(1)_*} P(1)_*$$

(similarly as in the proof of Corollary 9.1.7). By the classical invariant prime ideal theorem (see [Lan73a, Theorem 2.7] or compare [JY80, Theorem 1.16]), the invariant prime ideals of $P(1)_*$ are given by $I_m^{\text{top}} = (v_1^{\text{top}}, \cdots, v_{m-1}^{\text{top}})$. The isomorphism

$$AP(1)_{**}AP(1) \otimes_{AP(1)_{**}} AP(1)_{**} \cong P(1)_*P(1) \otimes_{P(1)_*} AP(1)_{**}$$

implies that I_m is an invariant prime ideal in $AP(1)_{**}$. By $AP(1)_{**} \cong P(1)_*[\tau]$, it follows also that \overline{I}_m is an invariant prime ideal of $AP(1)_{**}$, too.

It remains to show that there cannot be any further invariant prime ideals. As in [Lan73a], this follows from the fact that the only primitive elements in $AP(1)_{**}/\overline{I}_m \cong P(1)_*/I_m^{\text{top}}$ are multiples of powers of v_m (compare [Lan73a, Proposition 2.11]).

Corollary 9.6.3. (Motivic Landweber filtration theorem)

Let p be any prime and $n \ge 1$. For $X \in S\mathcal{H}(\mathbb{C})^{fin}$, $AP(n)_{**}(X)$ can be filtered by $AP(1)_{**}$ -modules

$$AP(n)_{**}(X) = M_0 \supset \cdots \supset M_k = 0$$

such that $M_i/M_{i+1} \cong AP(1)_{**}/I_m$ or $AP(1)_{**}/\overline{I}_m$ for some I_m and \overline{I}_m $(m \ge n)$ as above.

Proof. By Lemma 9.6.1, $AP(n)_{**}(X)$ is a coherent $AP(1)_{**}AP(1)$ -comodule. Landweber's filtration theorem [Lan73b, Theorem 3.3] (see also [JY80, Theorem 1.16]) implies that $AP(n)_{**}(X)$ has a filtration

$$AP(n)_{**}(X) = M_0 \supset \cdots \supset M_k = 0$$

such that $M_i/M_{i+1} \cong AP(1)_{**}/I$ for some I which is invariant under the comodule action. Thus, the claim follows from the previous lemma.

The following is a motivic version of one statement in [Rav84, Theorem 2.11]. In the proof, we use ideas from Ravenel's proof.

Theorem 9.6.4. Let p > 2. If $X \in S\mathcal{H}(\mathbb{C})^{fin}$ satisfies $AK(n+1)_{**}(X) = 0$, then also $AK(n)_{**}(X) = 0$. That is, $\langle AK(n+1) \rangle \geq \langle AK(n) \rangle$ in $S\mathcal{H}(\mathbb{C})^{fin}$.

Proof. Assume n > 0. The case n = 0 will be considered at the end of the proof. Let $E_{**}(-)$ be defined by

$$E_{**}X = AE(n+1)_{**} \otimes_{ABP_{**}} AP(n)_{**}(X).$$

As in [JY80, Lemma 3.5], the above Landweber filtration theorem and the fact that AE(n+1) is Landweber exact (see [NSØ09, Theorem 8.7]) yield

$$\operatorname{Tor}_{1}^{AP(n)_{**}}(AE(n+1)_{**} \otimes_{ABP_{**}} AP(n)_{**}, AP(n)_{**}/I) = 0$$

for all invariant prime ideals $I \subseteq AP(n)_{**}$ as above. (Alternatively, this can be derived from the topological analogue and Lemma 6.3.7, using [NSØ09, Theorem 8.7].) As in [Rav84, Theorem 2.11], this implies that $E_{**}(-)$ is an exact functor.

In analogy to [Rav84], we show that there is an injective pairing

$$AK(n+1)_{**} \otimes_{E_{**}} E_{**}X \hookrightarrow AK(n+1)_{**}X.$$

To construct this pairing, note that the ABP-action on AK(n+1) factors through a map $AP(n) \wedge AK(n+1) \rightarrow AK(n+1)$ by methods from Section 9.3, since v_i , i < n+1, acts trivially on AK(n+1) by Corollary 9.3.5. This induces a map

$$AK(n+1)_{**} \otimes_{ABP_{**}} AP(n)_{**}X \to AK(n+1)_{**}X$$
As in [Rav84, Theorem 2.11], this map factors through a pairing

$$AK(n+1)_{**} \otimes_{E_{**}} E_{**}X \to AK(n+1)_{**}X,$$

the reason being again that the relevant elements act trivially. Such a pairing induces a universal coefficient spectral sequence, whose motivic version is constructed in [DI05, Propositions 7.7 and 7.10],

$$\operatorname{Tor}_{i}^{E_{**}}(E_{**}(X), AK(n+1)_{**}) \Rightarrow AK(n+1)_{**}(X).$$

As in [Rav84], to prove the injectivity of the pairing, it suffices to prove the vanishing of

$$\operatorname{Tor}_{i}^{ABP_{**}}(AP(n)_{**}(X), AK(n+1)_{**})$$

for i > 1. Since $n \ge 1$, p acts trivially on both of these modules, and we can replace ABP by AP(1) = ABP/p. Hence, we have to show the vanishing of

$$\operatorname{Tor}_{i}^{AP(1)_{**}}(AP(n)_{**}(X), AK(n+1)_{**})$$

and, by Corollary 9.6.3, the question reduces to the vanishing of

1. D(1)

$$\operatorname{Tor}_{i}^{AP(1)_{**}}(AP(1)_{**}/I, AK(n+1)_{**})$$

for i > 1 and $I = I_m$ or \overline{I}_m , $m \ge n$. By Lemma 6.3.7, this equals

$$\operatorname{Tor}_{i}^{P(1)_{*}[\tau]}(P(1)_{*}/I_{m}[\tau], K(n+1)_{*}[\tau]) \text{ or}$$
$$\operatorname{Tor}_{i}^{P(1)_{*}[\tau]}(P(1)_{*}/I_{m}, K(n+1)_{*}[\tau]), \text{ respectively.}$$

A projective resolution of $K(n+1)_*$ over $P(1)_*$ yields a projective resolution of $K(n+1)_*[\tau]$ over $P(1)_*[\tau]$ by applying $-\otimes \mathbb{F}_p[\tau]$. It follows that both of the above torsion terms vanish if

$$\operatorname{Tor}_{i}^{P(1)_{*}}(P(1)_{*}/I_{m}, K(n+1)_{*}) = 0.$$

For i > 1, this follows from

$$\operatorname{Tor}_{i}^{BP_{*}}(BP_{*}/I_{m}, K(n+1)_{*}) = 0,$$

as in the proof of [Rav84, Theorem 2.11]. This proves the injectivity of the above pairing.

Now, assume that $AK(n+1)_{**}X = 0$. The injectivity of the pairing implies that $AK(n+1)_{**} \otimes_{E_{**}} E_{**}X = 0$. Recall $AK(n+1)_{**} \cong H_{**}[v_{n+1}^{\pm 1}]$ and note that

$$E_{**} \cong AE(n+1)_{**} \otimes_{ABP_{**}} AP(n)_{**} \cong E(n+1)_* \otimes_{BP_*} P(n)_*[\tau] \cong H_{**}[v_n, v_{n+1}^{\pm 1}]$$

by [NSØ09, Theorem 8.7] and Lemma 6.3.7. Therefore, $AK(n+1)_{**} \otimes_{E_{**}} E_{**}X = 0$ implies $v_n^{-1}E_{**}X = 0$. Now,

$$0 = v_n^{-1} E_{**} X = A E(n+1)_{**} \otimes_{ABP_{**}} A B(n)_{**} X,$$

by the definitions of $E_{**}(-)$ and of AB(n). By Theorem 9.4.16, since X is finite, this is equal to

$$AE(n+1)_{**} \otimes_{ABP_{**}} AK(n)_{**}(X) \otimes \mathbb{F}_p[v_{n+1}, v_{n+2}, \cdots].$$

By [NSØ09, Theorem 8.7], $AE(n+1)_{**} \cong E(n+1)_* \otimes_{BP_*} ABP_{**}$. It follows

$$0 = \mathbb{Z}_{(p)}[v_1, \cdots, v_n, v_{n+1}^{\pm 1}] \otimes_{\mathbb{Z}_{(p)}[v_1, \cdots]} AK(n)_{**}X \otimes \mathbb{F}_p[v_{n+1}, \cdots]$$

As v_m acts trivially on $AK(n)_{**}X$ for all $m \neq n$ by Corollary 9.3.5, this implies that $AK(n)_{**}X = 0$.

It remains to show that $AK(1)_{**}X = 0$ implies $AK(0)_{**}X = 0$. Recall the definitions $AE(1) = v_1^{-1}ABP/(v_2, \cdots)$ and $AK(1) = v_1^{-1}ABP/(p, v_2, \cdots) = AE(1)/p$. Since multiplication by p commutes with $\pi_{**}(-)$, it follows that $AK(1)_{**}X = 0$ implies $p^{-1}AE(1)_{**}X = 0$. (Note that $p^{-1}(AE(1)_{**}X) \cong (p^{-1}AE(1))_{**}X$ since $\pi_{**}(-)$ commutes with filtered colimits by [DI05, Proposition 9.3].) The following argument is closely related to Corollary 9.1.8. Since $p^{-1}AE(1)_{**}X = 0$, $p^{-1}(ABP/(v_2, \cdots))_{**}X$ is v_1 -torsion. By Theorem 9.1.1, this implies that $p^{-1}(ABP/(v_2, \cdots))_{**}X$ is p-torsion. But this can only be the case if $p^{-1}(ABP/(v_2, \cdots))_{**}X = 0$. It follows that

$$AE(0)_{**}X = p^{-1}(ABP/(v_1, v_2, \cdots))_{**}X = 0.$$

Now, by Theorem 9.5.1, $\langle AE(0) \rangle = \langle AK(0) \rangle$. Hence, $AK(0)_{**}X = 0$.

In terms of thick ideals in $\mathcal{SH}(\mathbb{C})_{(p)}^{fin}$, p > 2, we have proven that the motivic Morava K-theory spectra AK(n) indeed describe a descending chain of thick ideals, similarly to the chain of thick subcategories in $\mathcal{SH}_{(p)}^{fin}$. The inclusions are proper by Chapter 8.

Corollary 9.6.5. *For* p > 2*,*

$$\mathcal{SH}(\mathbb{C})^{fin}_{(p)} \supseteq \mathcal{C}_{AK(0)} \supseteq \mathcal{C}_{AK(1)} \supseteq \cdots$$

To sum up, we have identified three sequences of thick ideals which may or may not be equal:

$$\begin{split} \mathcal{SH}(\mathbb{C})_{(p)}^{fin} &\supset R^{-1}(\mathcal{C}_1) &\supset R^{-1}(\mathcal{C}_2) \supset \cdots \\ & || & \cup | & \cup | \\ \mathcal{SH}(\mathbb{C})_{(p)}^{fin} \supset \mathcal{C}_{AK(0)} \supset \mathcal{C}_{AK(1)} \supset \cdots \\ & || & \cup | & \cup | \\ \mathcal{SH}(\mathbb{C})_{(p)}^{fin} \supset \text{thickid}(c\mathcal{C}_1) \supset \text{thickid}(c\mathcal{C}_2) \supset \cdots . \end{split}$$

Bibliography

- [Ada69] J. F. Adams. "Lectures on generalised cohomology". In: Category Theory, Homology Theory and Their Applications III. Springer Berlin Heidelberg, 1969.
- [Ada74] J. F. Adams. *Stable Homotopy and generalised Homology*. University of Chicago Press, 1974.
- [Ayo07] J. Ayoub. "Les six oprations de Grothendieck et le formalisme des cycles vanescents dans le monde motivique". In: Astrisque 314-315 (2007).
- [Ayo10] J. Ayoub. "Note sur les oprations de Grothendieck et la ralisation de Betti". In: Journal of the Institute of Mathematics of Jussieu 9 issue 2 (2010), pp. 225-263.
- [Bal05] P. Balmer. "The spectrum of prime ideals in tensor triangulated categories". In: J. Reine Angew. Math 588 (2005), pp. 149–168.
- [Bal10] P. Balmer. "Spectra, spectra, spectra tensor triangular spectra versus Zariski spectra of endomorphism rings". In: Algebr. Geom. Topol. 10.3 (2010), pp. 1521–1563.
- [BM13] M. Basterra and M. A. Mandell. "The multiplication on BP". In: Journal of Topology 6 (2013), pp. 285–310.
- [Bor03] S. Borghesi. "Algebraic Morava K-theories". In: Inventiones Mathematicae 151 (2003), pp. 381–413.
- [CS69] P. E. Conner and L. Smith. "On the complex bordism of finite complexes". In: Publ. Math. Inst. Hautes Etudes Sci. 37 (1969), pp. 117–221.
- [DHS88] E. S. Devinatz, M. J. Hopkins, and J. H. Smith. "Nilpotence and stable homotopy theory I". In: Annals of Mathematics 128 (1988), pp. 207–241.
- [DI04] D. Dugger and D. Isaksen. "Topological hypercovers and A1-realizations". In: *Mathematische Zeitschrift* 246 (2004), pp. 667–689.
- [DI05] D. Dugger and D. Isaksen. "Motivic cell structures". In: Algebraic and Geometric Topology 5 (2005), pp. 615–652.
- [DI10] D. Dugger and D. Isaksen. "The motivic Adams spectral sequence". In: Geometry & Topology 14 (2010), pp. 967–1014.

[Dug01]	D. Dugger. "Replacing model categories with simplicial ones". In: <i>Transac-</i> <i>tions of the American Mathematical Society</i> 353 (2001), pp. 5003–5027.
[Dug12]	D. Dugger. Motivic stable homotopy groups of spheres. 2012. URL: http://sma.epfl.ch/~hessb
[Elm+97]	A. D. Elmendorf et al. <i>Rings, modules and algebras in stable homotopy the-</i> ory. American Mathematical Society, 1997.
[FJM01]	H. Fausk, L.G. Lewis Jr., and J.P. May. "The Picard Group of Equivariant Stable Homotopy Theory". In: <i>Advances in Mathematics</i> 163 (2001), pp. 17–33.
[GM95]	J. P. C. Greenlees and J. P. May. <i>Generalized Tate cohomology</i> . Memoirs of the AMS, 1995.
[Gui06]	B. Guillou. A short note on models for equivariant homotopy theory. 2006. URL: http://www.math.uiuc.edu/~bertg/EquivModels.pdf.
[Hat01]	A. Hatcher. Algebraic Topology. Cambridge University Press, 2001.
[HKO11]	P. Hu, I. Kriz, and K. Ormsby. "Convergence of the motivic Adams spectral sequence". In: <i>Journal of K-theory</i> 7, Issue 03 (2011), pp. 573–596.
[HMS94]	M. J. Hopkins, M. Mahowald, and H. Sadofsky. "Constructions of elements in Picard groups". In: <i>Contemporary Mathematics</i> 158 (1994), pp. 89–126.
[HO14]	J. Heller and K. Ormsby. "Galois equivariance and stable motivic homotopy theory". arXiv:1401.4728v2. 2014.
[Hor06]	J. Hornbostel. "Localizations in motivic homotopy theory". In: <i>Math. Proc.</i> <i>Camb. Phil. Soc.</i> 140 (2006), pp. 95–114.
[Hov01]	M. Hovey. "Spectra and symmetric spectra in general homotopy categories". In: Journal of Pure and Applied Algebra 165 (2001), pp. 63–127.
[Hov99]	M. Hovey. Model Categories. AMS, 1999.
[Hoy13]	M. Hoyois. "From algebraic cobordism to motivic cohomology". In: Journal für die reine und angewandte Mathematik (Crelles Journal) (2013).
[HS96]	M. Hovey and H. Sadofsky. "Tate cohomology lowes cromatic Bousfield classes". In: <i>Proc. Amer. Math. Soc.</i> 124 (1996), pp. 3579–3585.
[HS98]	M. J. Hopkins and J. H. Smith. "Nilpotence and stable homotopy theory II". In: Annals of Mathematics 148 (1998), pp. 1–49.
[HSS99]	M. Hovey, B. Shipley, and J. Smith. "Symmetric Spectra". In: J. Amer. Math. Soc. 13 (1999), pp. 149–208.
[Hu01]	P. Hu. "Base change functors in the A1-stable homotopy category". In: <i>Ho-mology, Homotopy and Applications</i> 3 (2001), pp. 417–451.
[Hu03]	P. Hu. "S-modules in the category of schemes". In: Mem. Amer. Math. Soc. 767 (2003), 125pp.
[Hu05]	P. Hu. "On the Picard group of the A^1 -stable homotopy category". In: <i>Topology</i> 44, Issue 3 (2005), pp. 609–640.

[Hul12]	K. Hulek. Elementare Algebraische Geometrie. Springer, 2012.
[Jar00]	J. F. Jardine. "Motivic Symmetric Spectra". In: <i>Documenta Mathematica</i> 5 (2000), pp. 445–552.
[Jou73]	J. P. Jouanolou. "Une Suite exacte de Mayer-Vietoris en K-Theorie algebrique". In: <i>Higher K-Theories.</i> 1973.
[JW75]	D. C. Johnson and W. S. Wilson. "BP Operations and Morava's Extraordinary K-Theories". In: <i>Math. Z.</i> 144 (1975), pp. 55–75.
[JY80]	D. C. Johnson and Z. Yosimura. "Torsion in Brown-Peterson homology and Hurewicz homomorphisms". In: Osaka J. Math. 17 (1980), pp. 117–136.
[Lam05]	T. Y. Lam. <i>Introduction to Quadratic Forms over Fields</i> . Graduate Studies in Mathematics, vol. 67, AMS, 2005.
[Lan73a]	P. S. Landweber. "Annihilator ideals and primitive elements in complex cobordism". In: <i>Illinois J. Math.</i> 17 (1973), pp. 273–284.
[Lan73b]	P. S. Landweber. "Associated prime ideals and Hopf algebras". In: <i>Journal of pure and applied algebra</i> 3 (1973), pp. 43–58.
[Lan76]	P. S. Landweber. "Homological properties of comodules over MU*MU and BP*BP". In: American Journal of Mathematics 98 (1976), pp. 591–610.
[Laz04]	R. K. Lazarsfeld. Positivity in Algebraic Geometry I. Springer, 2004.
[Lev08]	M. Levine. "The homotopy coniveau tower". In: <i>Journal of Topology</i> 1 (2008), pp. 217–267.
[Lev14]	M. Levine. "A comparison of motivic and classical stable homotopy theories". In: <i>Journal of Topology</i> 7 (2014), pp. 327–362.
[LM06]	L. G. Lewis and M. A. Mandell. "Equivariant universal coefficient and Künneth spectral sequences". In: <i>Proc. London Math. Soc.</i> 92 (2006), pp. 505–544.
[LMS86]	L. G. Lewis, J. P. May, and M. Steinberger. <i>Equivariant Stable Homotopy Theory</i> . Springer, 1986.
[Man02]	M. A. Mandell. "Equivariant Symmetric Spectra". In: Contemporary Mathematics 346. AMS, 2002, pp. 399–452.
[May89]	K. H. Mayer. "G-invariante Morse Funktionen". In: manuscripta mathematica 63 (1989), pp. 99–114.
[May96]	J. P. May. <i>Equivariant homotopy and cohomology theory</i> . AMS and CBMS, 1996.
[Mil63]	J. Milnor. Morse Theory. Princeton Univ. Press, 1963.
[Mit85]	S. A. Mitchell. "Finite complexes with A(n)-free cohomology". In: <i>Topology</i> 24 (1985), pp. 227–248.
[MM02]	M. A. Mandell and J. P. May. <i>Equivariant orthogonal spectra and S-modules</i> . Memoirs of the AMS, 2002.

[Mor02]	F. Morel. "On the motivic π_0 of the sphere spectrum". In: Axiomatic, Enriched and Motivic Homotopy Theory. Ed. by J.P.C.Greenlees. NATO Science Series, 2002, pp. 219–260.
[Mor03]	F. Morel. "An introduction to A1-homotopy theory". In: Contemporary developments in algebraic K-theory, I.C.T.P. Lecture Notes 15 (2003), pp. 357–441.
[Mor06]	F. Morel. "A1-algebraic topology". In: Proceedings of the International Con- gress of Mathematicians Madrid (2006).
[Mor99]	F. Morel. "Suite spectrale d'Adams et invariants cohomologiques des formes quadratiques". In: C.R. Acad. Sci. Paris 328, Serie I (1999), pp. 963–968.
[MS06]	J. P. May and J. Sigurdsson. "Parametrized homotopy theory". In: <i>Mathematical Surveys and Monographs. American Mathematical Society</i> 132 (2006).
[MV99]	F. Morel and V. Voevodsky. "A ¹ -homotopy theory of schemes". In: <i>Publica-</i> tions Mathematiques de l'IHS 90 (1999), pp. 45–143.
[Nas95]	C. Nassau. Eine nichtgeometrische Konstruktion der Spektren $P(n)$ und mul- tiplikative Automorphismen von $K(n)$. Diplomarbeit. Goethe Universität Frankfurt am Main. 1995. URL: http://publikationen.ub.uni-frankfurt.de/frontdoor/ir
[Nee01]	A. Neeman. Triangulated Categories. Princeton Univ. Press, 2001.
[Nee96]	A. Neeman. "The Grothendieck duality theorem via Bousfield's techniques and Brown representability". In: <i>Journal of the AMS</i> 9 (1996), pp. 205–236.
[NSØ09]	N. Naumann, M. Spitzweck, and P. A. Østvær. "Motivic Landweber Exactness". In: <i>Documenta Mathematica</i> 14 (2009), pp. 551–593.
[Pel11]	P. Pelaez. "Multiplicative properties of the slice filtration". In: Astrisque 335 (2011), pp. 1–289.
[PPR08]	I. Panin, K. Pimenov, and O. Röndigs. "A universality theorem for Voevod- sky's algebraic cobordism spectrum". In: <i>Homology, Homotopy and Appli-</i> <i>cations</i> 10(2) (2008), pp. 211–226.
[PPR09]	I. Panin, K. Pimenov, and O. Röndigs. "On Voevodsky's Algebraic K-Theory Spectrum". In: <i>Algebraic Topology</i> . Ed. by Nils Baas et al. Vol. 4. Abel Symposia. Springer Berlin Heidelberg, 2009, pp. 279–330.
[PY02]	I. Panin and S. Yagunov. "Rigidity for orientable functors". In: <i>Journal of pure and applied algebra</i> 172 (2002), pp. 49–77.
[Qui71]	D. Quillen. "Elementary Proofs of Some Results of Cobordism Theory Using Steenrod Operations". In: Advances in Mathematics 7 (1971), pp. 29–56.
[Rav84]	D. C. Ravenel. "Localization with Respect to Certain Periodic Homology Theories". In: <i>American Journal of Mathematics</i> 106 (1984), pp. 351–414.
[Rav86]	D. C. Ravenel. Complex Cobordism and Stable Homotopy Groups of Spheres. Academic Press, 1986.

[Rav92]	D. C. Ravenel. <i>Nilpotence and Periodicity in Stable Homotopy Theory</i> . Princeton University Press, 1992.
[Rio05]	J. Riou. "Dualité de Spanier-Whitehead en géométrie algébrique". In: C. R. Acad. Sci. Paris I 340 (2005), pp. 431–436.
[RØ08]	O. Röndigs and P. A. Østvær. "Modules over motivic cohomology". In: Advances in Mathematics 219, Issue 2 (2008), pp. 689–727.
[Sch12]	S. Schwede. Symmetric Spectra. 2012. URL: http://www.math.uni-bonn.de/people/schwede/
[Sch13]	S. Schwede. Lectures on equivariant stable homotopy theory. 2013. URL: http://www.math.uni-bonn.de/people/schwede/equivariant.pdf.
[Seg70]	G. B. Segal. "Equivariant stable homotopy theory". In: Actes, Congrès in- tern. Math. 2 (1970), pp. 59–63.
[Spi10]	M. Spitzweck. "Relations between slices and quotients of the algebraic cobordism spectrum". In: <i>Homology, Homotopy and Applications</i> 12, Issue 2 (2010), pp. 335–351.
[SS00]	S. Schwede and B. Shipley. "Algebras and modules in monoidal model cat- egories". In: <i>Proc. London Math. Soc.</i> 80 (2000), pp. 491–511.
[Sta12]	S. Stahn. Die motivische Adams-Spektralsequenz. 2012. URL: http://www2.math.uni-wupperta
[Str10]	N. P. Strickland. "Thick ideals of finite G-Spectra". Unpublished. 2010.
[Str99]	N. Strickland. "Products on MU-modules". In: <i>Transactions of the American Mathematical Society</i> 351 (1999), pp. 2569–2606.
[Vez01]	G. Vezzosi. "Brown-Peterson spectra in stable A^1 -homotopy theory". In: <i>Rend. Sem. Mat. Univ. di Padova</i> 106 (2001), pp. 47–64.
[Voe03a]	V. Voevodsky. "Motivic cohomology with Z/2 coefficients". In: <i>Publ. Math. Inst. Hautes Etudes Sci.</i> 98 (2003), pp. 1–57.
[Voe03b]	V. Voevodsky. "Reduced power operations in motivic cohomology". In: <i>Publications Mathematiques de l'IHS</i> 98 (2003), pp. 1–57.
[Voe10]	V. Voevodsky. "Motivic Eilenberg-MacLane spaces". In: <i>Publications mathematiques de l'IHES</i> 112 (2010), pp. 1–99.
[Voe96]	V. Voevodsky. "The Milnor Conjecture". Preprint. 1996.
[Voe98]	V. Voevodsky. "A ¹ -Homotopy Theory". In: <i>Documenta Mathematica</i> Extra Volume ICM (1998), pp. 579–604.
[Was69]	A. G. Wasserman. "Equivariant differential topology". In: <i>Topology</i> 8 (1969), pp. 127–150.
[Wür76]	U. Würgler. "Cobordism Theories of Unitary Manifolds with Singularities and Formal Group Laws". In: <i>Math. Z.</i> 150 (1976), pp. 239–260.
[Wür77]	U. Würgler. "On Products in a Family of Cohomology Theories Associated to the Invariant Prime Ideals of $\pi_*(BP)$ ". In: Comment. Math. Helvetici 52 (1977), pp. 457–481.

[Yag05] N. Yagita. "Applications of Atiyah-Hirzebruch spectral sequences for motivic cobordisms". In: Proceedings of the London Mathematical Society 90 (2005), pp. 783–816.

Ruth Joachimi Fachbereich Mathematik und Naturwissenschaften Bergische Universität Wuppertal 42119 Wuppertal Germany ruth.joachimi@d-fine.de